## Gisela Di Venanzio

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8123808/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Multidrug-resistant plasmids repress chromosomally encoded T6SS to enable their dissemination. Proceedings of the National Academy of Sciences of the United States of America, 2019, 116, 1378-1383.	7.1	83
2	Urinary tract colonization is enhanced by a plasmid that regulates uropathogenic Acinetobacter baumannii chromosomal genes. Nature Communications, 2019, 10, 2763.	12.8	80
3	Pathogenic Acinetobacter species have a functional type I secretion system and contact-dependent inhibition systems. Journal of Biological Chemistry, 2017, 292, 9075-9087.	3.4	73
4	Serratia marcescens ShlA Pore-Forming Toxin Is Responsible for Early Induction of Autophagy in Host Cells and Is Transcriptionally Regulated by RcsB. Infection and Immunity, 2014, 82, 3542-3554.	2.2	64
5	Capsule carbohydrate structure determines virulence in Acinetobacter baumannii. PLoS Pathogens, 2021, 17, e1009291.	4.7	59
6	Serratia marcescens Is Able to Survive and Proliferate in Autophagic-Like Vacuoles inside Non-Phagocytic Cells. PLoS ONE, 2011, 6, e24054.	2.5	46
7	Peptidoglycan editing provides immunity to <i>Acinetobacter baumannii</i> during bacterial warfare. Science Advances, 2020, 6, eabb5614.	10.3	44
8	The PhoP/PhoQ System and Its Role in Serratia marcescens Pathogenesis. Journal of Bacteriology, 2012, 194, 2949-2961.	2.2	37
9	The Glycoprotease CpaA Secreted by Medically Relevant Acinetobacter Species Targets Multiple <i>O</i> -Linked Host Glycoproteins. MBio, 2020, 11, .	4.1	31
10	Modern Acinetobacter baumannii clinical isolates replicate inside spacious vacuoles and egress from macrophages. PLoS Pathogens, 2021, 17, e1009802.	4.7	21
11	The Phenylacetic Acid Catabolic Pathway Regulates Antibiotic and Oxidative Stress Responses in Acinetobacter. MBio, 2022, 13, e0186321.	4.1	18
12	A pore-forming toxin enables <i>Serratia</i> a nonlytic egress from host cells. Cellular Microbiology, 2017, 19, e12656.	2.1	16
13	InvL, an Invasin-Like Adhesin, Is a Type II Secretion System Substrate Required for Acinetobacter baumannii Uropathogenesis. MBio, 2022, 13, .	4.1	11
14	Evolutionarily stable gene clusters shed light on the common grounds of pathogenicity in the Acinetobacter calcoaceticus-baumannii complex. PLoS Genetics, 2022, 18, e1010020.	3.5	10
15	Plasmid-Encoded H-NS Controls Extracellular Matrix Composition in a Modern <i>Acinetobacter baumannii</i> Urinary Isolate. Journal of Bacteriology, 2021, 203, e0027721.	2.2	9
16	The fliR gene contributes to the virulence of S. marcescens in a Drosophila intestinal infection model. Scientific Reports, 2022, 12, 3068.	3.3	3