

D Margriet Ouwens

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/817812/publications.pdf>

Version: 2024-02-01

118
papers

6,323
citations

61984

43
h-index

71685

76
g-index

118
all docs

118
docs citations

118
times ranked

9527
citing authors

#	ARTICLE	IF	CITATIONS
1	Postprandial renal haemodynamic effects of the dipeptidyl peptidase-4 inhibitor linagliptin versus the sulphonylurea glimepiride in adults with type 2 diabetes (<scp>RENALIS</scp>): A predefined substudy of a randomized, double-blind trial. <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 115-124.	4.4	7
2	Nudix hydrolase NUDT19 regulates mitochondrial function and ATP production in murine hepatocytes. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2022, 1867, 159153.	2.4	4
3	Hepatic Wnt1 Inducible Signaling Pathway Protein 1 (WISP-1/CCN4) Associates with Markers of Liver Fibrosis in Severe Obesity. <i>Cells</i> , 2021, 10, 1048.	4.1	7
4	Crosstalk of Diabetic Conditions with Static Versus Dynamic Flow Environmentâ€™Impact on Aortic Valve Remodeling. <i>International Journal of Molecular Sciences</i> , 2021, 22, 6976.	4.1	2
5	Divergent dynamics in systemic and tissue-specific metabolic and inflammatory responses during weight loss in subjects with obesity. <i>Cytokine</i> , 2021, 144, 155587.	3.2	3
6	Effects of DPP-4 Inhibitor Linagliptin Versus Sulfonylurea Glimepiride as Add-on to Metformin on Renal Physiology in Overweight Patients With Type 2 Diabetes (RENALIS): A Randomized, Double-Blind Trial. <i>Diabetes Care</i> , 2020, 43, 2889-2893.	8.6	10
7	Physiological Disturbance in Fatty Liver Energy Metabolism Converges on IGFBP2 Abundance and Regulation in Mice and Men. <i>International Journal of Molecular Sciences</i> , 2020, 21, 4144.	4.1	22
8	Effects of dipeptidyl peptidase-4 inhibitor linagliptin versus sulphonylurea glimepiride on systemic haemodynamics in overweight patients with type 2 diabetes: A secondary analysis of an 8-week, randomized, controlled, double-blind trial. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 1847-1856.	4.4	6
9	One-leg inactivity induces a reduction in mitochondrial oxidative capacity, intramyocellular lipid accumulation and reduced insulin signalling upon lipid infusion: a human study with unilateral limb suspension. <i>Diabetologia</i> , 2020, 63, 1211-1222.	6.3	18
10	Rhein, a novel Histone Deacetylase (HDAC) inhibitor with antifibrotic potency in human myocardial fibrosis. <i>Scientific Reports</i> , 2020, 10, 4888.	3.3	22
11	The small chain fatty acid butyrate antagonizes the TCR-stimulation-induced metabolic shift in murine epidermal gamma delta T cells. <i>EXCLI Journal</i> , 2020, 19, 334-350.	0.7	3
12	The adipokine sFRP4 induces insulin resistance and lipogenesis in the liver. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2019, 1865, 2671-2684.	3.8	28
13	Increased Glycolysis and Higher Lactate Production in Hyperglycemic Myotubes. <i>Cells</i> , 2019, 8, 1101.	4.1	30
14	Impact of hyperinsulinemia and hyperglycemia on valvular interstitial cells â€™ A link between aortic heart valve degeneration and type 2 diabetes. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2019, 1865, 2526-2537.	3.8	16
15	Sfrp5 increases glucose-stimulated insulin secretion in the rat pancreatic beta cell line INS-1E. <i>PLoS ONE</i> , 2019, 14, e0213650.	2.5	11
16	Insulin resistance associates with hepatic lobular inflammation in subjects with obesity. <i>Endocrine Connections</i> , 2019, 8, 1294-1301.	1.9	7
17	CDH13 abundance interferes with adipocyte differentiation and is a novel biomarker for adipose tissue health. <i>International Journal of Obesity</i> , 2018, 42, 1039-1050.	3.4	15
18	Deletion of the RabGAP TBC1D1 Leads to Enhanced Insulin Secretion and Fatty Acid Oxidation in Islets From Male Mice. <i>Endocrinology</i> , 2018, 159, 1748-1761.	2.8	9

#	ARTICLE	IF	CITATIONS
19	Assessment of circulating Wnt1 inducible signalling pathway protein 1 (WISP-1)/CCN4 as a novel biomarker of obesity. <i>Journal of Cell Communication and Signaling</i> , 2018, 12, 539-548.	3.4	30
20	Enzymatic Activity Is Not Required for Phospholipase D Mediated TNF- α Regulation and Myocardial Healing. <i>Frontiers in Physiology</i> , 2018, 9, 1698.	2.8	3
21	Increased triacylglycerol - Fatty acid substrate cycling in human skeletal muscle cells exposed to eicosapentaenoic acid. <i>PLoS ONE</i> , 2018, 13, e0208048.	2.5	15
22	The novel adipokine WISP1 associates with insulin resistance and impairs insulin action in human myotubes and mouse hepatocytes. <i>Diabetologia</i> , 2018, 61, 2054-2065.	6.3	34
23	Secretory products from epicardial adipose tissue from patients with type 2 diabetes impair mitochondrial β -oxidation in cardiomyocytes via activation of the cardiac renin-angiotensin system and induction of miR-208a. <i>Basic Research in Cardiology</i> , 2017, 112, 2.	5.9	47
24	Soluble CD14 inhibits contractile function and insulin action in primary adult rat cardiomyocytes. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2017, 1863, 365-374.	3.8	3
25	Effect of the long-acting insulin analogues glargine and degludec on cardiomyocyte cell signalling and function. <i>Cardiovascular Diabetology</i> , 2016, 15, 96.	6.8	6
26	Reduced expression of chemerin in visceral adipose tissue associates with hepatic steatosis in patients with obesity. <i>Obesity</i> , 2016, 24, 2544-2552.	3.0	23
27	Determinants of testosterone levels in human male obesity. <i>Endocrine</i> , 2015, 50, 202-211.	2.3	48
28	Adiponectin may mediate the association between omentin, circulating lipids and insulin sensitivity: results from the KORA F4 study. <i>European Journal of Endocrinology</i> , 2015, 172, 423-432.	3.7	62
29	Over-expression of PRAS40 enhances insulin sensitivity in skeletal muscle. <i>Archives of Physiology and Biochemistry</i> , 2014, 120, 64-72.	2.1	25
30	Does dipeptidyl peptidase-4 inhibition prevent the diabetogenic effects of glucocorticoids in men with the metabolic syndrome? A randomized controlled trial. <i>European Journal of Endocrinology</i> , 2014, 170, 429-439.	3.7	36
31	Secretome profiling of primary human skeletal muscle cells. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2014, 1844, 1011-1017.	2.3	138
32	Adipocyte-derived factors impair insulin signaling in differentiated human vascular smooth muscle cells via the upregulation of miR-143. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2014, 1842, 275-283.	3.8	25
33	Identification of novel adipokines differential regulated in C57BL/Ks and C57BL/6. <i>Archives of Physiology and Biochemistry</i> , 2014, 120, 208-215.	2.1	5
34	Cerebrospinal fluid levels of Alzheimer's disease biomarkers in middle-aged patients with type 1 diabetes. <i>Diabetologia</i> , 2014, 57, 2208-2214.	6.3	40
35	Tissue-Specific Differences in the Development of Insulin Resistance in a Mouse Model for Type 1 Diabetes. <i>Diabetes</i> , 2014, 63, 3856-3867.	0.6	51
36	Effect of Sfrp5 on Cytokine Release and Insulin Action in Primary Human Adipocytes and Skeletal Muscle Cells. <i>PLoS ONE</i> , 2014, 9, e85906.	2.5	36

#	ARTICLE	IF	CITATIONS
37	Knockdown of PRAS40 inhibits insulin action via proteasome-mediated degradation of IRS1 in primary human skeletal muscle cells. <i>Diabetologia</i> , 2013, 56, 1118-1128.	6.3	18
38	The insulin sensitizing effect of topiramate involves K_{ATP} channel activation in the central nervous system. <i>British Journal of Pharmacology</i> , 2013, 170, 908-918.	5.4	18
39	Activin A impairs insulin action in cardiomyocytes via up-regulation of miR-143. <i>Cardiovascular Research</i> , 2013, 100, 201-210.	3.8	57
40	Activin a is associated with impaired myocardial glucose metabolism and left ventricular remodeling in patients with uncomplicated type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2013, 12, 150.	6.8	32
41	Beneficial and Adverse Effects of Testosterone on the Cardiovascular System in Men. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 4300-4310.	3.6	86
42	Prednisolone induces the Wnt signalling pathway in 3T3-L1 adipocytes. <i>Archives of Physiology and Biochemistry</i> , 2013, 119, 52-64.	2.1	8
43	Overexpression of AMP-activated protein kinase or protein kinase D prevents lipid-induced insulin resistance in cardiomyocytes. <i>Journal of Molecular and Cellular Cardiology</i> , 2013, 55, 165-173.	1.9	14
44	Proline-rich Akt substrate of 40-kDa contains a nuclear export signal. <i>Cellular Signalling</i> , 2013, 25, 1762-1768.	3.6	5
45	Glucocorticoid treatment impairs microvascular function in healthy men in association with its adverse effects on glucose metabolism and blood pressure: a randomised controlled trial. <i>Diabetologia</i> , 2013, 56, 2383-2391.	6.3	26
46	VEGF in the Crosstalk between Human Adipocytes and Smooth Muscle Cells: Depot-Specific Release from Visceral and Perivascular Adipose Tissue. <i>Mediators of Inflammation</i> , 2013, 2013, 1-10.	3.0	43
47	Cardioprotective Properties of Omentin-1 in Type 2 Diabetes: Evidence from Clinical and In Vitro Studies. <i>PLoS ONE</i> , 2013, 8, e59697.	2.5	87
48	CD36 inhibition prevents lipid accumulation and contractile dysfunction in rat cardiomyocytes. <i>Biochemical Journal</i> , 2012, 448, 43-53.	3.7	73
49	Identification and Validation of Novel Adipokines Released from Primary Human Adipocytes. <i>Molecular and Cellular Proteomics</i> , 2012, 11, M111.010504.	3.8	187
50	Role of PRAS40 in Akt and mTOR signaling in health and disease. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012, 302, E1453-E1460.	3.5	133
51	Secretory Products From Epicardial Adipose Tissue of Patients With Type 2 Diabetes Mellitus Induce Cardiomyocyte Dysfunction. <i>Circulation</i> , 2012, 126, 2324-2334.	1.6	155
52	Involvement of atypical protein kinase C in the regulation of cardiac glucose and long-chain fatty acid uptake. <i>Frontiers in Physiology</i> , 2012, 3, 361.	2.8	8
53	Sex Steroid-Induced Changes in Circulating Monocyte Chemoattractant Protein-1 Levels May Contribute to Metabolic Dysfunction in Obese Men. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, E1187-E1191.	3.6	20
54	Chemerin as biomarker for insulin sensitivity in males without typical characteristics of metabolic syndrome. <i>Archives of Physiology and Biochemistry</i> , 2012, 118, 135-138.	2.1	38

#	ARTICLE	IF	CITATIONS
55	Heat Shock Protein 60 as a Mediator of Adipose Tissue Inflammation and Insulin Resistance. <i>Diabetes</i> , 2012, 61, 615-625.	0.6	62
56	Endogenous oestradiol and cardiovascular disease in healthy men: a systematic review and meta-analysis of prospective studies. <i>Heart</i> , 2012, 98, 1478-1482.	2.9	20
57	High Oxidative Capacity Due to Chronic Exercise Training Attenuates Lipid-Induced Insulin Resistance. <i>Diabetes</i> , 2012, 61, 2472-2478.	0.6	71
58	Effects of Adding Exercise to a 16-Week Very Low-Calorie Diet in Obese, Insulin-Dependent Type 2 Diabetes Mellitus Patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, 2512-2520.	3.6	57
59	Targeting of mitochondrial reactive oxygen species production does not avert lipid-induced insulin resistance in muscle tissue from mice. <i>Diabetologia</i> , 2012, 55, 2759-2768.	6.3	37
60	Glucose Intolerance and the Amount of Visceral Adipose Tissue Contribute to an Increase in Circulating Triglyceride Concentrations in Caucasian Obese Females. <i>PLoS ONE</i> , 2012, 7, e45145.	2.5	12
61	Secretory products of guinea pig epicardial fat induce insulin resistance and impair primary adult rat cardiomyocyte function. <i>Journal of Cellular and Molecular Medicine</i> , 2011, 15, 2399-2410.	3.6	53
62	Short-term increase of plasma free fatty acids does not interfere with intrinsic mitochondrial function in healthy young men. <i>Metabolism: Clinical and Experimental</i> , 2011, 60, 1398-1405.	3.4	14
63	Prednisolone-induced beta cell dysfunction is associated with impaired endoplasmic reticulum homeostasis in INS-1E cells. <i>Cellular Signalling</i> , 2011, 23, 1708-1715.	3.6	43
64	Subcellular trafficking of the substrate transporters GLUT4 and CD36 in cardiomyocytes. <i>Cellular and Molecular Life Sciences</i> , 2011, 68, 2525-2538.	5.4	66
65	Circulating insulin stimulates fatty acid retention in white adipose tissue via KATP channel activation in the central nervous system only in insulin-sensitive mice. <i>Journal of Lipid Research</i> , 2011, 52, 1712-1722.	4.2	21
66	Dipeptidyl Peptidase 4 Is a Novel Adipokine Potentially Linking Obesity to the Metabolic Syndrome. <i>Diabetes</i> , 2011, 60, 1917-1925.	0.6	506
67	Combined Gene and Protein Expression of Hormone-Sensitive Lipase and Adipose Triglyceride Lipase, Mitochondrial Content, and Adipocyte Size in Subcutaneous and Visceral Adipose Tissue of Morbidly Obese Men. <i>Obesity Facts</i> , 2011, 4, 407-416.	3.4	29
68	Glucagon-Like Peptide-1 Receptor Agonist Treatment Prevents Glucocorticoid-Induced Glucose Intolerance and Islet-Cell Dysfunction in Humans. <i>Diabetes Care</i> , 2011, 34, 412-417.	8.6	117
69	Absence of fatty acid transporter CD36 protects against Western-type diet-related cardiac dysfunction following pressure overload in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2011, 301, E618-E627.	3.5	44
70	High levels of dietary stearate promote adiposity and deteriorate hepatic insulin sensitivity. <i>Nutrition and Metabolism</i> , 2010, 7, 24.	3.0	39
71	The dopamine receptor D2 agonist bromocriptine inhibits glucose-stimulated insulin secretion by direct activation of the β -adrenergic receptors in beta cells. <i>Biochemical Pharmacology</i> , 2010, 79, 1827-1836.	4.4	67
72	Phosphorylation of PRAS40 on Thr246 by PKB/AKT facilitates efficient phosphorylation of Ser183 by mTORC1. <i>Cellular Signalling</i> , 2010, 22, 961-967.	3.6	74

#	ARTICLE	IF	CITATIONS
73	The role of epicardial and perivascular adipose tissue in the pathophysiology of cardiovascular disease. <i>Journal of Cellular and Molecular Medicine</i> , 2010, 14, 2223-2234.	3.6	192
74	Differential regulation of cardiac GLUT4-mediated glucose and CD36-mediated fatty acid uptake by endosomal pH and actin filaments. <i>Chemistry and Physics of Lipids</i> , 2010, 163, S11.	3.2	0
75	Diabetic cardiomyopathy in Zucker diabetic fatty rats: the forgotten right ventricle. <i>Cardiovascular Diabetology</i> , 2010, 9, 25.	6.8	43
76	Sex Steroids Affect Triglyceride Handling, Glucose-Dependent Insulinotropic Polypeptide, and Insulin Sensitivity. <i>Diabetes Care</i> , 2010, 33, 1831-1833.	8.6	31
77	Differential regulation of cardiac glucose and fatty acid uptake by endosomal pH and actin filaments. <i>American Journal of Physiology - Cell Physiology</i> , 2010, 298, C1549-C1559.	4.6	35
78	Permissive action of protein kinase C- α 1 in insulin-induced CD36- and GLUT4 translocation in cardiac myocytes. <i>Journal of Endocrinology</i> , 2009, 201, 199-209.	2.6	32
79	MIF Deficiency Reduces Chronic Inflammation in White Adipose Tissue and Impairs the Development of Insulin Resistance, Glucose Intolerance, and Associated Atherosclerotic Disease. <i>Circulation Research</i> , 2009, 105, 99-107.	4.5	138
80	Altered myocardial substrate metabolism is associated with myocardial dysfunction in early diabetic cardiomyopathy in rats: studies using positron emission tomography. <i>Cardiovascular Diabetology</i> , 2009, 8, 39.	6.8	102
81	Novel insights into glucocorticoid-mediated diabetogenic effects: towards expansion of therapeutic options?. <i>European Journal of Clinical Investigation</i> , 2009, 39, 81-93.	3.4	351
82	The nuclear appearance of ERK1/2 and p38 determines the sequential induction of ATF2-Thr71 and ATF2-Thr69 phosphorylation by serum in JNK-deficient cells. <i>Molecular and Cellular Endocrinology</i> , 2009, 311, 94-100.	3.2	11
83	PRAS40: Target or modulator of mTORC1 signalling and insulin action?. <i>Archives of Physiology and Biochemistry</i> , 2009, 115, 163-175.	2.1	29
84	Loss of 50% of excess weight using a very low energy diet improves insulin-stimulated glucose disposal and skeletal muscle insulin signalling in obese insulin-treated type 2 diabetic patients. <i>Diabetologia</i> , 2008, 51, 309-319.	6.3	63
85	Intracerebroventricular Administration of Neuropeptide Y Induces Hepatic Insulin Resistance via Sympathetic Innervation. <i>Diabetes</i> , 2008, 57, 2304-2310.	0.6	101
86	Dynamics of insulin signalling in liver during hyperinsulinemic euglycaemic clamp conditions in vivo and the effects of high-fat feeding in male mice. <i>Archives of Physiology and Biochemistry</i> , 2007, 113, 173-185.	2.1	5
87	Myocardial insulin action and the contribution of insulin resistance to the pathogenesis of diabetic cardiomyopathy. <i>Archives of Physiology and Biochemistry</i> , 2007, 113, 76-86.	2.1	35
88	Lessons that can be learned from patients with diabetogenic mutations in mitochondrial DNA: implications for common type 2 diabetes. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2007, 10, 693-697.	2.5	19
89	Functioning of oxidative phosphorylation in liver mitochondria of high-fat diet fed rats. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2007, 1772, 307-316.	3.8	47
90	SREBP-1c expression in Schwann cells is affected by diabetes and nutritional status. <i>Molecular and Cellular Neurosciences</i> , 2007, 35, 525-534.	2.2	32

#	ARTICLE	IF	CITATIONS
91	Cardiac contractile dysfunction in insulin-resistant rats fed a high-fat diet is associated with elevated CD36-mediated fatty acid uptake and esterification. <i>Diabetologia</i> , 2007, 50, 1938-1948.	6.3	190
92	Evaluation of proinflammatory cytokines and inflammation markers as biomarkers for the action of thiazolidinediones in Type 2 diabetes mellitus patients and healthy volunteers. <i>British Journal of Clinical Pharmacology</i> , 2006, 62, 391-402.	2.4	27
93	Regulation of Sarcolemmal Transport of Substrates in the Healthy and Diseased Heart. <i>Cardiovascular Drugs and Therapy</i> , 2006, 20, 471-476.	2.6	53
94	Sustained activation of the mammalian target of rapamycin nutrient sensing pathway is associated with hepatic insulin resistance, but not with steatosis, in mice. <i>Diabetologia</i> , 2006, 49, 3049-3057.	6.3	53
95	The Role of c-Jun N-Terminal Kinase, p38, and Extracellular Signal-Regulated Kinase in Insulin-Induced Thr69 and Thr71 Phosphorylation of Activating Transcription Factor 2. <i>Molecular Endocrinology</i> , 2006, 20, 1786-1795.	3.7	25
96	Endogenous Interleukin-10 Protects against Hepatic Steatosis but Does Not Improve Insulin Sensitivity during High-Fat Feeding in Mice. <i>Endocrinology</i> , 2006, 147, 4553-4558.	2.8	76
97	Insulin-Mediated Phosphorylation of the Proline-Rich Akt Substrate PRAS40 Is Impaired in Insulin Target Tissues of High-Fat Diet-Fed Rats. <i>Diabetes</i> , 2006, 55, 3221-3228.	0.6	50
98	Cardiac dysfunction induced by high-fat diet is associated with altered myocardial insulin signalling in rats. <i>Diabetologia</i> , 2005, 48, 1229-1237.	6.3	213
99	Acute hepatic steatosis in mice by blocking β -oxidation does not reduce insulin sensitivity of very-low-density lipoprotein production. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 289, G592-G598.	3.4	56
100	Effect of a 2-day very low-energy diet on skeletal muscle insulin sensitivity in obese type 2 diabetic patients on insulin therapy. <i>Metabolism: Clinical and Experimental</i> , 2005, 54, 1669-1678.	3.4	10
101	IRS-4 mediated mitogenic signalling by insulin and growth hormone in LB cells, a murine T-cell lymphoma devoid of IGF-I receptors. <i>Cellular Signalling</i> , 2003, 15, 385-394.	3.6	9
102	Arsenite stimulated glucose transport in 3T3-L1 adipocytes involves both Glut4 translocation and p38 MAPK activity. <i>FEBS Journal</i> , 2003, 270, 3891-3903.	0.2	57
103	Hepatic VLDL Production in ob/ob Mice Is Not Stimulated by Massive De Novo Lipogenesis but Is Less Sensitive to the Suppressive Effects of Insulin. <i>Diabetes</i> , 2003, 52, 1081-1089.	0.6	80
104	CD36 deficiency increases insulin sensitivity in muscle, but induces insulin resistance in the liver in mice. <i>Journal of Lipid Research</i> , 2003, 44, 2270-2277.	4.2	155
105	Increased Hepatic Insulin Sensitivity Together with Decreased Hepatic Triglyceride Stores in Hormone-Sensitive Lipase-Deficient Mice. <i>Endocrinology</i> , 2003, 144, 3456-3462.	2.8	104
106	Skin fibroblasts of children with idiopathic short stature show an increased mitogenic response to IGF-I and secrete more IGFBP-3. <i>Clinical Endocrinology</i> , 2002, 56, 439-447.	2.4	14
107	Growth factors can activate ATF2 via a two-step mechanism: phosphorylation of Thr71 through the Ras-MEK-ERK pathway and of Thr69 through RalGDS-Src-p38. <i>EMBO Journal</i> , 2002, 21, 3782-3793.	7.8	204
108	Modulation of insulin-stimulated glycogen synthesis by Src Homology Phosphatase 2. <i>Molecular and Cellular Endocrinology</i> , 2001, 175, 131-140.	3.2	19

#	ARTICLE	IF	CITATIONS
109	Hyperosmotic stress activates the insulin receptor in CHO cells. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2001, 1540, 97-106.	4.1	25
110	Molecular mechanisms of contraction-regulated cardiac glucose transport. <i>Biochemical Journal</i> , 2000, 346, 841.	3.7	5
111	Mammalian target of rapamycin is a direct target for protein kinase B: identification of a convergence point for opposing effects of insulin and amino-acid deficiency on protein translation. <i>Biochemical Journal</i> , 1999, 344, 427.	3.7	203
112	Two Naturally Occurring Insulin Receptor Tyrosine Kinase Domain Mutants Provide Evidence That Phosphoinositide 3-Kinase Activation Alone Is Not Sufficient for the Mediation of Insulin's Metabolic and Mitogenic Effects. <i>Journal of Biological Chemistry</i> , 1997, 272, 30208-30214.	3.4	79
113	Expression, Enzyme Activity, and Subcellular Localization of Mammalian Target of Rapamycin in Insulin-Responsive Cells. <i>Biochemical and Biophysical Research Communications</i> , 1997, 241, 704-709.	2.1	69
114	Replacement of the Conserved Tyrosine 1210 by Phenylalanine in the Insulin Receptor Affects Insulin-Induced Dephosphorylation of Focal Adhesion Kinase but Leaves Other Responses Intact. <i>Biochemistry</i> , 1996, 35, 10377-10382.	2.5	10
115	Insulin-induced tyrosine dephosphorylation of paxillin and focal adhesion kinase requires active phosphotyrosine phosphatase 1D. <i>Biochemical Journal</i> , 1996, 318, 609-614.	3.7	43
116	Expression of a dominant-negative Ras mutant does not affect stimulation of glucose uptake and glycogen synthesis by insulin. <i>Diabetologia</i> , 1996, 39, 558-563.	6.3	3
117	Presence of Gonadotropin-Releasing Hormone (GnRH) Binding Sites and Compounds with GnRH-Like Activity in the Ovary of African Catfish, <i>Clarias Gariepinus</i> 1. <i>Biology of Reproduction</i> , 1994, 50, 643-652.	2.7	28
118	Activation of Overexpressed Receptors for Insulin and Epidermal Growth Factor Interferes in Mitogenic Signaling without Affecting the Activation of p21ras. <i>Biochemistry</i> , 1994, 33, 7453-7459.	2.5	10