

# Ãsa Cider

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8205840/publications.pdf>

Version: 2024-02-01

45  
papers

1,295  
citations

361413

20  
h-index

361022

35  
g-index

46  
all docs

46  
docs citations

46  
times ranked

1586  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Exercise as migraine prophylaxis: A randomized study using relaxation and topiramate as controls. Cephalalgia, 2011, 31, 1428-1438.   | 3.9 | 207       |
| 2  | Does moderate-to-high intensity Nordic walking improve functional capacity and pain in fibromyalgia? A prospective randomized controlled trial. Arthritis Research and Therapy, 2010, 12, R189.                         | 3.5 | 92        |
| 3  | Hydrotherapy-a new approach to improve function in the older patient with chronic heart failure. European Journal of Heart Failure, 2003, 5, 527-535.   | 7.1 | 87        |
| 4  | A Study to Evaluate the Feasibility of an Aerobic Exercise Program in Patients With Migraine. Headache, 2009, 49, 563-570.  | 3.9 | 74        |
| 5  | The impact on kinesiophobia (fear of movement) by clinical variables for patients with coronary artery disease. International Journal of Cardiology, 2013, 167, 391-397.  | 1.7 | 68        |
| 6  | Immersion in warm water induces improvement in cardiac function in patients with chronic heart failure. European Journal of Heart Failure, 2006, 8, 308-313.  | 7.1 | 55        |
| 7  | Muscle function in adults with congenital heart disease. International Journal of Cardiology, 2014, 170, 358-363.   | 1.7 | 47        |
| 8  | Validation of a questionnaire to detect kinesiophobia (fear of movement) in patients with coronary artery disease. Journal of Rehabilitation Medicine, 2012, 44, 363-369.   | 1.1 | 45        |
| 9  | Kinesiophobia mediates the influences on attendance at exercise-based cardiac rehabilitation in patients with coronary artery disease. Physiotherapy Theory and Practice, 2016, 32, 571-580.                            | 1.3 | 38        |
| 10 | Cardiorespiratory effects of warm water immersion in elderly patients with chronic heart failure. Clinical Physiology and Functional Imaging, 2005, 25, 313-317.  | 1.2 | 36        |
| 11 | Exercise in Elderly Patients with Chronic Heart Failure in Primary Care: Effects on Physical Capacity and Health-Related Quality of Life. European Journal of Cardiovascular Nursing, 2011, 10, 150-158.                | 0.9 | 36        |
| 12 | The 2011 outcome from the Swedish Health Care Registry on Heart Disease (SWEDEHEART). Scandinavian Cardiovascular Journal, 2013, 47, 1-10.  | 1.2 | 35        |
| 13 | Reliability of Clinical Muscular Endurance Tests in Patients with Chronic Heart Failure. European Journal of Cardiovascular Nursing, 2006, 5, 122-126.  | 0.9 | 34        |
| 14 | Muscular performance in heart failure. Journal of Cardiac Failure, 1998, 4, 97-104.   | 1.7 | 32        |
| 15 | A group-based exercise program did not improve physical activity in patients with chronic heart failure and comorbidity: A randomized controlled trial. Journal of Rehabilitation Medicine, 2014, 46, 461-467.          | 1.1 | 32        |
| 16 | Effects of High Frequency Exercise in Patients before and after Elective Percutaneous Coronary Intervention. European Journal of Cardiovascular Nursing, 2008, 7, 307-313.  | 0.9 | 30        |
| 17 | Physical inactivity and smoking after myocardial infarction as predictors for readmission and survival: results from the SWEDEHEART-registry. Clinical Research in Cardiology, 2019, 108, 324-332.                      | 3.3 | 29        |
| 18 | Participation in exercise-based cardiac rehabilitation is related to reduced total mortality in both men and women: results from the SWEDEHEART registry. European Journal of Preventive Cardiology, 2022, 29, 485-492. | 1.8 | 28        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Benefit of warm water immersion on biventricular function in patients with chronic heart failure. <i>Cardiovascular Ultrasound</i> , 2009, 7, 33.  | 1.6 | 26        |
| 20 | Physical Activity on Prescription (PAP), in patients with metabolic risk factors. A 6-month follow-up study in primary health care. <i>PLoS ONE</i> , 2017, 12, e0175190.  | 2.5 | 26        |
| 21 | SWEDHEART Annual Report 2012. <i>Scandinavian Cardiovascular Journal</i> , 2014, 48, 1-1.  | 1.2 | 25        |
| 22 | Provocation of Migraine after Maximal Exercise: A Test-Retest Study. <i>European Neurology</i> , 2017, 78, 22-27.  | 1.4 | 21        |
| 23 | Test-retest reliability, agreement, and minimal detectable change in the 6-minute walk test in patients with intermittent claudication. <i>Journal of Vascular Surgery</i> , 2020, 71, 197-203.                          | 1.1 | 21        |
| 24 | Effects of Early Bedside Cycle Exercise on Intracranial Pressure and Systemic Hemodynamics in Critically Ill Patients in a Neurointensive Care Unit. <i>Neurocritical Care</i> , 2016, 25, 434-439.                      | 2.4 | 19        |
| 25 | Physical activity in relation to cardiac risk markers in secondary prevention of coronary artery disease. <i>International Journal of Cardiology</i> , 2013, 168, 478-483.   | 1.7 | 18        |
| 26 | Which patients benefit from physical activity on prescription (PAP)? A prospective observational analysis of factors that predict increased physical activity. <i>BMC Public Health</i> , 2019, 19, 482.                 | 2.9 | 18        |
| 27 | Relevance of Kinesiophobia in Relation to Changes Over Time Among Patients After an Acute Coronary Artery Disease Event. <i>Journal of Cardiopulmonary Rehabilitation and Prevention</i> , 2018, 38, 224-230.            | 2.1 | 15        |
| 28 | Peripheral muscle training with resistance exercise bands in patients with chronic heart failure. Long-term effects on walking distance and quality of life; a pilot study. <i>ESC Heart Failure</i> , 2018, 5, 241-248. | 3.1 | 15        |
| 29 | Acute effects of physical exercise on the serum insulin-like growth factor system in women with fibromyalgia. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 37.   | 1.9 | 10        |
| 30 | Exercise-based cardiac rehabilitation improves physical fitness in patients with permanent atrial fibrillation – A randomized controlled study. <i>Translational Sports Medicine</i> , 2020, 3, 415-425.                 | 1.1 | 10        |
| 31 | Test-retest reliability of six-minute walk tests over a one-year period in patients with chronic heart failure. <i>Clinical Physiology and Functional Imaging</i> , 2020, 40, 284-289.                                   | 1.2 | 10        |
| 32 | Cerebrovascular and systemic haemodynamic parameters during passive exercise. <i>Advances in Physiotherapy</i> , 2010, 12, 58-63.  | 0.2 | 9         |
| 33 | Is hydrotherapy an appropriate form of exercise for elderly patients with biventricular systolic heart failure?. <i>Journal of Geriatric Cardiology</i> , 2013, 9, 408-410.  | 0.2 | 9         |
| 34 | Long-term physical activity on prescription intervention for patients with insufficient physical activity level – a randomized controlled trial. <i>Trials</i> , 2020, 21, 793.  | 1.6 | 9         |
| 35 | Blood flow velocity and vascular resistance during passive leg exercise in the critically ill patient. <i>Clinical Physiology and Functional Imaging</i> , 2012, 32, 338-342.  | 1.2 | 7         |
| 36 | Effects of exercise on fatigue and physical capacity in men with chronic widespread pain - a pilot study. <i>BMC Sports Science, Medicine and Rehabilitation</i> , 2016, 8, 29.  | 1.7 | 7         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | High frequency home-based exercise decreases levels of vascular endothelial growth factor in patients with stable angina pectoris. <i>European Journal of Preventive Cardiology</i> , 2015, 22, 575-581.                        | 1.8 | 4         |
| 38 | Reliability of two questionnaires on physical function in patients with stable coronary artery disease. <i>European Journal of Cardiovascular Nursing</i> , 2016, 15, 142-149.  | 0.9 | 3         |
| 39 | Exercise capacity, physical activity, and health-related quality of life in adults with CHD. <i>Cardiology in the Young</i> , 2020, 30, 668-673.  | 0.8 | 3         |
| 40 | Fractures in children and young adults with and without congenital heart disease. <i>International Journal of Cardiology Congenital Heart Disease</i> , 2021, 2, 100072.  | 0.4 | 2         |
| 41 | Nonresponders of Physical Activity on Prescription (PAP) Can Increase Their Exercise Capacity with Enhanced Physiotherapist Support. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 4795. | 2.6 | 1         |
| 42 | Effects of exercise training, with or without supplemental oxygen, in adults with complex congenital heart disease. <i>International Journal of Cardiology Congenital Heart Disease</i> , 2021, 3, 100109.                      | 0.4 | 1         |
| 43 | Effects of 3 months of detraining following cardiac rehabilitation in patients with atrial fibrillation. <i>European Review of Aging and Physical Activity</i> , 2022, 19, 14.  | 2.9 | 1         |
| 44 | Response to "Exercise programmes and quality of life in the elderly: important facts". <i>European Journal of Cardiovascular Nursing</i> , 2012, 11, 128-128.   | 0.9 | 0         |
| 45 | Post-operative musculoskeletal outcomes in patients with coarctation of the aorta following different surgical approaches. <i>International Journal of Cardiology</i> , 2021, 327, 80-85.                                       | 1.7 | 0         |