Ãsa Cider

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8205840/publications.pdf

Version: 2024-02-01

45 papers 1,295 citations

20 h-index 35 g-index

46 all docs

46 docs citations

times ranked

46

1586 citing authors

#	Article	IF	CITATIONS
1	Participation in exercise-based cardiac rehabilitation is related to reduced total mortality in both men and women: results from the SWEDEHEART registry. European Journal of Preventive Cardiology, 2022, 29, 485-492.	1.8	28
2	Effects of 3 months of detraining following cardiac rehabilitation in patients with atrial fibrillation. European Review of Aging and Physical Activity, 2022, 19, 14.	2.9	1
3	Fractures in children and young adults with and without congenital heart disease. International Journal of Cardiology Congenital Heart Disease, 2021, 2, 100072.	0.4	2
4	Post-operative musculoskeletal outcomes in patients with coarctation of the aorta following different surgical approaches. International Journal of Cardiology, 2021, 327, 80-85.	1.7	0
5	Nonresponders of Physical Activity on Prescription (PAP) Can Increase Their Exercise Capacity with Enhanced Physiotherapist Support. International Journal of Environmental Research and Public Health, 2021, 18, 4795.	2.6	1
6	Effects of exercise training, with or without supplemental oxygen, in adults with complex congenital heart disease. International Journal of Cardiology Congenital Heart Disease, 2021, 3, 100109.	0.4	1
7	Test-retest reliability, agreement, and minimal detectable change in the 6-minute walk test in patients with intermittent claudication. Journal of Vascular Surgery, 2020, 71, 197-203.	1.1	21
8	Exercise $\hat{a} \in b$ as ed cardiac rehabilitation improves physical fitness in patients with permanent atrial fibrillation $\hat{a} \in a$ A randomized controlled study. Translational Sports Medicine, 2020, 3, 415-425.	1.1	10
9	Long-term physical activity on prescription intervention for patients with insufficient physical activity levelâ€"a randomized controlled trial. Trials, 2020, 21, 793.	1.6	9
10	Test–retest reliability of sixâ€minute walk tests over a oneâ€year period in patients with chronic heart failure. Clinical Physiology and Functional Imaging, 2020, 40, 284-289.	1.2	10
11	Exercise capacity, physical activity, and health-related quality of life in adults with CHD. Cardiology in the Young, 2020, 30, 668-673.	0.8	3
12	Which patients benefit from physical activity on prescription (PAP)? A prospective observational analysis of factors that predict increased physical activity. BMC Public Health, 2019, 19, 482.	2.9	18
13	Physical inactivity and smoking after myocardial infarction as predictors for readmission and survival: results from the SWEDEHEART-registry. Clinical Research in Cardiology, 2019, 108, 324-332.	3.3	29
14	Relevance of Kinesiophobia in Relation to Changes Over Time Among Patients After an Acute Coronary Artery Disease Event. Journal of Cardiopulmonary Rehabilitation and Prevention, 2018, 38, 224-230.	2.1	15
15	Peripheral muscle training with resistance exercise bands in patients with chronic heart failure. Longâ€ŧerm effects on walking distance and quality of life; a pilot study. ESC Heart Failure, 2018, 5, 241-248.	3.1	15
16	Acute effects of physical exercise on the serum insulin-like growth factor system in women with fibromyalgia. BMC Musculoskeletal Disorders, 2017, 18, 37.	1.9	10
17	Provocation of Migraine after Maximal Exercise: A Test-Retest Study. European Neurology, 2017, 78, 22-27.	1.4	21
18	Physical Activity on Prescription (PAP), in patients with metabolic risk factors. A 6-month follow-up study in primary health care. PLoS ONE, 2017, 12, e0175190.	2.5	26

#	Article	IF	CITATIONS
19	Effects of exercise on fatigue and physical capacity in men with chronic widespread pain - a pilot study. BMC Sports Science, Medicine and Rehabilitation, 2016, 8, 29.	1.7	7
20	Effects of Early Bedside Cycle Exercise on Intracranial Pressure and Systemic Hemodynamics in Critically III Patients in a Neurointensive Care Unit. Neurocritical Care, 2016, 25, 434-439.	2.4	19
21	Kinesiophobia mediates the influences on attendance at exercise-based cardiac rehabilitation in patients with coronary artery disease. Physiotherapy Theory and Practice, 2016, 32, 571-580.	1.3	38
22	Reliability of two questionnaires on physical function in patients with stable coronary artery disease. European Journal of Cardiovascular Nursing, 2016, 15, 142-149.	0.9	3
23	High frequency home-based exercise decreases levels of vascular endothelial growth factor in patients with stable angina pectoris. European Journal of Preventive Cardiology, 2015, 22, 575-581.	1.8	4
24	A group-based exercise program did not improve physical activity in patients with chronic heart failure and comorbidity: A randomized controlled trial. Journal of Rehabilitation Medicine, 2014, 46, 461-467.	1.1	32
25	SWEDEHEART Annual Report 2012. Scandinavian Cardiovascular Journal, 2014, 48, 1-1.	1.2	25
26	Muscle function in adults with congenital heart disease. International Journal of Cardiology, 2014, 170, 358-363.	1.7	47
27	The 2011 outcome from the Swedish Health Care Registry on Heart Disease (SWEDEHEART). Scandinavian Cardiovascular Journal, 2013, 47, 1-10.	1.2	35
28	Physical activity in relation to cardiac risk markers in secondary prevention of coronary artery disease. International Journal of Cardiology, 2013, 168, 478-483.	1.7	18
29	The impact on kinesiophobia (fear of movement) by clinical variables for patients with coronary artery disease. International Journal of Cardiology, 2013, 167, 391-397.	1.7	68
30	Is hydrotherapy an appropriate form of exercise for elderly patients with biventricular systolic heart failure?. Journal of Geriatric Cardiology, 2013, 9, 408-410.	0.2	9
31	Response to â€Exercise programmes and quality of life in the elderly: important facts'. European Journal of Cardiovascular Nursing, 2012, 11, 128-128.	0.9	0
32	Validation of a questionnaire to detect kinesiophobia (fear of movement) in patients with coronary artery disease. Journal of Rehabilitation Medicine, 2012, 44, 363-369.	1.1	45
33	Blood flow velocity and vascular resistance during passive leg exercise in the critically ill patient. Clinical Physiology and Functional Imaging, 2012, 32, 338-342.	1.2	7
34	Exercise as migraine prophylaxis: A randomized study using relaxation and topiramate as controls. Cephalalgia, 2011, 31, 1428-1438.	3.9	207
35	Exercise in Elderly Patients with Chronic Heart Failure in Primary Care: Effects on Physical Capacity and Health-Related Quality of Life. European Journal of Cardiovascular Nursing, 2011, 10, 150-158.	0.9	36
36	Cerebrovascular and systemic haemodynamic parameters during passive exercise. Advances in Physiotherapy, 2010, 12, 58-63.	0.2	9

Ã...sa Cider

#	Article	IF	CITATIONS
37	Does moderate-to-high intensity Nordic walking improve functional capacity and pain in fibromyalgia? A prospective randomized controlled trial. Arthritis Research and Therapy, 2010, 12, R189.	3.5	92
38	Benefit of warm water immersion on biventricular function in patients with chronic heart failure. Cardiovascular Ultrasound, 2009, 7, 33.	1.6	26
39	A Study to Evaluate the Feasibility of an Aerobic Exercise Program in Patients With Migraine. Headache, 2009, 49, 563-570.	3.9	74
40	Effects of High Frequency Exercise in Patients before and after Elective Percutaneous Coronary Intervention. European Journal of Cardiovascular Nursing, 2008, 7, 307-313.	0.9	30
41	Reliability of Clinical Muscular Endurance Tests in Patients with Chronic Heart Failure. European Journal of Cardiovascular Nursing, 2006, 5, 122-126.	0.9	34
42	Immersion in warm water induces improvement in cardiac function in patients with chronic heart failure. European Journal of Heart Failure, 2006, 8, 308-313.	7.1	55
43	Cardiorespiratory effects of warm water immersion in elderly patients with chronic heart failure. Clinical Physiology and Functional Imaging, 2005, 25, 313-317.	1.2	36
44	Hydrotherapy-a new approach to improve function in the older patient with chronic heart failure. European Journal of Heart Failure, 2003, 5, 527-535.	7.1	87
45	Muscular performance in heart failure. Journal of Cardiac Failure, 1998, 4, 97-104.	1.7	32