
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8243487/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	PRIME: A Programme to Reduce the Treatment Gap for Mental Disorders in Five Low- and Middle-Income Countries. PLoS Medicine, 2012, 9, e1001359.	8.4	436
2	Acceptability and feasibility of using non-specialist health workers to deliver mental health care: Stakeholder perceptions from the PRIME district sites in Ethiopia, India, Nepal, South Africa, and Uganda. Social Science and Medicine, 2014, 118, 33-42.	3.8	259
3	Challenges and Opportunities for Implementing Integrated Mental Health Care: A District Level Situation Analysis from Five Low- and Middle-Income Countries. PLoS ONE, 2014, 9, e88437.	2.5	255
4	What happens to patients with treatment-resistant depression? A systematic review of medium to long term outcome studies. Journal of Affective Disorders, 2009, 116, 4-11.	4.1	244
5	A Multidimensional Tool to Quantify Treatment Resistance in Depression. Journal of Clinical Psychiatry, 2009, 70, 177-184.	2.2	211
6	The path to longer and healthier lives for all Africans by 2030: the Lancet Commission on the future of health in sub-Saharan Africa. Lancet, The, 2017, 390, 2803-2859.	13.7	150
7	The impact of childhood adversity on suicidality and clinical course in treatment-resistant depression. Journal of Affective Disorders, 2014, 152-154, 122-130.	4.1	132
8	Validity of brief screening questionnaires to detect depression in primary care in Ethiopia. Journal of Affective Disorders, 2015, 186, 32-39.	4.1	131
9	Excess mortality in severe mental illness: 10-year population-based cohort study in rural Ethiopia. British Journal of Psychiatry, 2015, 206, 289-296.	2.8	130
10	Stigma, explanatory models and unmet needs of caregivers of children with developmental disorders in a low-income African country: a cross-sectional facility-based survey. BMC Health Services Research, 2016, 16, 152.	2.2	119
11	Toxoplasmosis as a cause for behaviour disorders - overview of evidence and mechanisms. Folia Parasitologica, 2010, 57, 105-113.	1.3	117
12	Clinical Course and Outcome of Schizophrenia in a Predominantly Treatment-Naive Cohort in Rural Ethiopia. Schizophrenia Bulletin, 2009, 35, 646-654.	4.3	103
13	The Maudsley Staging Method for Treatment-Resistant Depression. Journal of Clinical Psychiatry, 2009, 70, 952-957.	2.2	99
14	Population level mental distress in rural Ethiopia. BMC Psychiatry, 2014, 14, 194.	2.6	98
15	Development of a scalable mental healthcare plan for a rural district in Ethiopia. British Journal of Psychiatry, 2016, 208, s4-s12.	2.8	93
16	Evaluation of district mental healthcare plans: The PRIME consortium methodology. British Journal of Psychiatry, 2016, 208, s63-s70.	2.8	92
17	Internalized stigma among patients with schizophrenia in Ethiopia: a cross-sectional facility-based study. BMC Psychiatry, 2012, 12, 239.	2.6	89
18	"l cry every day and night, I have my son tied in chains― physical restraint of people with schizophrenia in community settings in Ethiopia. Globalization and Health, 2017, 13, 47.	4.9	78

#	Article	IF	CITATIONS
19	Containment of COVID-19 in Ethiopia and implications for tuberculosis care and research. Infectious Diseases of Poverty, 2020, 9, 131.	3.7	76
20	District mental healthcare plans for five low-and middle-income countries: Commonalities, variations and evidence gaps. British Journal of Psychiatry, 2016, 208, s47-s54.	2.8	73
21	Five-year mortality in a cohort of people with schizophrenia in Ethiopia. BMC Psychiatry, 2011, 11, 165.	2.6	69
22	Planning and evaluating mental health services in low- and middle-income countries using theory of change. British Journal of Psychiatry, 2016, 208, s55-s62.	2.8	69
23	Prediction of longer-term outcome of treatment-resistant depression in tertiary care. British Journal of Psychiatry, 2012, 201, 369-375.	2.8	65
24	Validation of the World Health Organization Disability Assessment Schedule in people with severe mental disorders in rural Ethiopia. Health and Quality of Life Outcomes, 2017, 15, 64.	2.4	64
25	Standardisation framework for the Maudsley staging method for treatment resistance in depression. BMC Psychiatry, 2018, 18, 100.	2.6	64
26	Community-based rehabilitation intervention for people with schizophrenia in Ethiopia (RISE): a 12 month mixed methods pilot study. BMC Psychiatry, 2018, 18, 250.	2.6	63
27	Treatment Contact Coverage for Probable Depressive and Probable Alcohol Use Disorders in Four Low- and Middle-Income Country Districts: The PRIME Cross-Sectional Community Surveys. PLoS ONE, 2016, 11, e0162038.	2.5	62
28	Treatment gap and help-seeking for postpartum depression in a rural African setting. BMC Psychiatry, 2016, 16, 196.	2.6	61
29	Using workshops to develop theories of change in five low and middle income countries: lessons from the programme for improving mental health care (PRIME). International Journal of Mental Health Systems, 2014, 8, 15.	2.7	60
30	Treatment gap, help-seeking, stigma and magnitude of alcohol use disorder in rural Ethiopia. Substance Abuse Treatment, Prevention, and Policy, 2019, 14, 4.	2.2	58
31	Participatory planning of a primary care service for people with severe mental disorders in rural Ethiopia. Health Policy and Planning, 2016, 31, 367-376.	2.7	55
32	The potential use of digital health technologies in the African context: a systematic review of evidence from Ethiopia. Npj Digital Medicine, 2021, 4, 125.	10.9	54
33	Suicide and suicide attempts in people with severe mental disorders in Butajira, Ethiopia: 10Âyear follow-up of a population-based cohort. BMC Psychiatry, 2014, 14, 150.	2.6	51
34	Scaling up integrated primary mental health in six low- and middle-income countries: obstacles, synergies and implications for systems reform. BJPsych Open, 2019, 5, e69.	0.7	51
35	Stakeholder analysis of the Programme for Improving Mental health carE (PRIME): baseline findings. International Journal of Mental Health Systems, 2015, 9, 27.	2.7	50
36	Multidimensional impact of severe mental illness on family members: systematic review. BMJ Open, 2019, 9, e032391.	1.9	49

#	Article	IF	CITATIONS
37	Low-coverage sequencing cost-effectively detects known and novel variation in underrepresented populations. American Journal of Human Genetics, 2021, 108, 656-668.	6.2	49
38	Adapting and pre-testing the World Health Organization's Caregiver Skills Training programme for autism and other developmental disorders in a very low-resource setting: Findings from Ethiopia. Autism, 2020, 24, 51-63.	4.1	48
39	Burden of mental disorders and unmet needs among street homeless people in Addis Ababa, Ethiopia. BMC Medicine, 2014, 12, 138.	5.5	47
40	Antenatal depressive symptoms and maternal health care utilisation: a population-based study of pregnant women in Ethiopia. BMC Pregnancy and Childbirth, 2016, 16, 301.	2.4	46
41	Hazardous alcohol use and associated factors in a rural Ethiopian district: a cross-sectional community survey. BMC Public Health, 2016, 16, 218.	2.9	46
42	Conceptualisation of job-related wellbeing, stress and burnout among healthcare workers in rural Ethiopia: a qualitative study. BMC Health Services Research, 2017, 17, 412.	2.2	46
43	Developing a Theory of Change model of service user and caregiver involvement in mental health system strengthening in primary health care in rural Ethiopia. International Journal of Mental Health Systems, 2020, 14, 51.	2.7	46
44	Development of a Community-Based Rehabilitation Intervention for People with Schizophrenia in Ethiopia. PLoS ONE, 2015, 10, e0143572.	2.5	46
45	Detection and treatment initiation for depression and alcohol use disorders: facility-based cross-sectional studies in five low-income and middle-income country districts. BMJ Open, 2018, 8, e023421.	1.9	45
46	Catastrophic health expenditure and impoverishment in households of persons with depression: a cross-sectional, comparative study in rural Ethiopia. BMC Public Health, 2019, 19, 930.	2.9	44
47	Coping strategies of women with postpartum depression symptoms in rural Ethiopia: a cross-sectional community study. BMC Psychiatry, 2018, 18, 41.	2.6	43
48	Symptomatic and functional outcome of bipolar disorder in Butajira, Ethiopia. Journal of Affective Disorders, 2006, 90, 239-249.	4.1	41
49	Estimating the cost of implementing district mental healthcare plans in five low- and middle-income countries: The PRIME study. British Journal of Psychiatry, 2016, 208, s71-s78.	2.8	40
50	Antenatal depressive symptoms and perinatal complications: a prospective study in rural Ethiopia. BMC Psychiatry, 2017, 17, 301.	2.6	40
51	Explanatory models of depression in sub-Saharan Africa: Synthesis of qualitative evidence. Social Science and Medicine, 2020, 246, 112760.	3.8	40
52	Violence and violent victimization in people with severe mental illness in a rural low-income country setting: A comparative cross-sectional community study. Schizophrenia Research, 2014, 152, 275-282.	2.0	39
53	Task sharing for the care of severe mental disorders in a low-income country (TaSCS): study protocol for a randomised, controlled, non-inferiority trial. Trials, 2016, 17, 76.	1.6	39
54	Homelessness and severe mental illness in low- and middle-income countries: scoping review. BJPsych Open, 2019, 5, e57.	0.7	39

#	Article	IF	CITATIONS
55	Beyond the Biomedical: Community Resources for Mental Health Care in Rural Ethiopia. PLoS ONE, 2015, 10, e0126666.	2.5	39
56	Antibacterial activity of methanol extracts of the leaves ofÂthree medicinal plants against selected bacteria isolated from wounds of lymphoedema patients. BMC Complementary Medicine and Therapies, 2021, 21, 2.	2.7	38
57	Recognition of depression by primary care clinicians in rural Ethiopia. BMC Family Practice, 2017, 18, 56.	2.9	36
58	Longitudinal course of symptom severity and fluctuation in patients with treatment-resistant unipolar and bipolar depression. Psychiatry Research, 2013, 207, 143-149.	3.3	34
59	Global mental health: perspectives from Ethiopia. Global Health Action, 2014, 7, 25447.	1.9	34
60	Community-based Rehabilitation Intervention for people with Schizophrenia in Ethiopia (RISE): study protocol for a cluster randomised controlled trial. Trials, 2016, 17, 299.	1.6	34
61	Health system governance to support scale up of mental health care in Ethiopia: a qualitative study. International Journal of Mental Health Systems, 2017, 11, 38.	2.7	34
62	The fight to end tuberculosis must not be forgotten in the COVID-19 outbreak. Nature Medicine, 2020, 26, 811-812.	30.7	34
63	Trimethoprim as Adjuvant Treatment in Schizophrenia: A Double-Blind, Randomized, Placebo-Controlled Clinical Trial. Schizophrenia Bulletin, 2010, 36, 846-851.	4.3	33
64	Developing a mental health care plan in a low resource setting: the theory of change approach. BMC Health Services Research, 2015, 15, 429.	2.2	33
65	Engaging and staying engaged: a phenomenological study of barriers to equitable access to mental healthcare for people with severe mental disorders in a rural African setting. International Journal for Equity in Health, 2017, 16, 156.	3.5	33
66	Under detection of depression in primary care settings in low and middle-income countries: a systematic review and meta-analysis. Systematic Reviews, 2022, 11, 21.	5.3	32
67	Non-fatal suicidal behaviour in rural Ethiopia: a cross-sectional facility- and population-based study. BMC Psychiatry, 2016, 16, 75.	2.6	31
68	Training needs and perspectives of community health workers in relation to integrating child mental health care into primary health care in a rural setting in sub-Saharan Africa: a mixed methods study. International Journal of Mental Health Systems, 2017, 11, 15.	2.7	31
69	Electronic pillbox-enabled self-administered therapy versus standard directly observed therapy for tuberculosis medication adherence and treatment outcomes in Ethiopia (SELFTB): protocol for a multicenter randomized controlled trial. Trials, 2020, 21, 383.	1.6	30
70	Postpartum depressive symptoms in the context of high social adversity and reproductive health threats: a population-based study. International Journal of Mental Health Systems, 2018, 12, 42.	2.7	29
71	Bipolar disorder among an isolated island community in Ethiopia. Journal of Affective Disorders, 2004, 80, 1-10.	4.1	28
72	Long-term clinical course and outcome of schizophrenia in rural Ethiopia: 10-year follow-up of a population-based cohort. Schizophrenia Research, 2015, 161, 414-420.	2.0	27

#	Article	IF	CITATIONS
73	Scaling-up essential neuropsychiatric services in Ethiopia: a cost-effectiveness analysis. Health Policy and Planning, 2016, 31, 504-513.	2.7	27
74	Global mental health and schizophrenia. Current Opinion in Psychiatry, 2018, 31, 193-199.	6.3	27
75	Ethiopian community health workers' beliefs and attitudes towards children with autism: Impact of a brief training intervention. Autism, 2019, 23, 39-49.	4.1	27
76	Burden of tuberculosis and challenges related to screening and diagnosis in Ethiopia. Journal of Clinical Tuberculosis and Other Mycobacterial Diseases, 2020, 19, 100158.	1.3	27
77	Psychological interventions for bipolar disorder in low- and middle-income countries: systematic review. BJPsych Open, 2018, 4, 375-384.	0.7	26
78	Burnout among primary healthcare workers during implementation of integrated mental healthcare in rural Ethiopia: a cohort study. Human Resources for Health, 2019, 17, 58.	3.1	26
79	Prospective evaluation of specialist inpatient treatment for refractory affective disorders. Journal of Affective Disorders, 2011, 131, 92-103.	4.1	25
80	Equitable access to integrated primary mental healthcare for people with severe mental disorders in Ethiopia: a formative study. International Journal for Equity in Health, 2016, 15, 121.	3.5	24
81	Partnerships in a Global Mental Health Research Programme—the Example of PRIME. Global Social Welfare, 2019, 6, 159-175.	1.9	24
82	Five-year clinical course and outcome of schizophrenia in Ethiopia. Schizophrenia Research, 2012, 136, 137-142.	2.0	23
83	The psychosis treatment gap and its consequences in rural Ethiopia. BMC Psychiatry, 2019, 19, 325.	2.6	23
84	Catastrophic out-of-pocket payments for households of people with severe mental disorder: a comparative study in rural Ethiopia. International Journal of Mental Health Systems, 2019, 13, 39.	2.7	23
85	Building the capacity of policy-makers and planners to strengthen mental health systems in low- and middle-income countries: a systematic review. BMC Health Services Research, 2016, 16, 601.	2.2	22
86	Antenatal depressive symptoms and utilisation of delivery and postnatal care: a prospective study in rural Ethiopia. BMC Pregnancy and Childbirth, 2017, 17, 206.	2.4	22
87	A descriptive analysis of depression and pain complaints among patients with cancer in a low income country. PLoS ONE, 2018, 13, e0193713.	2.5	22
88	Movement disorders in neuroleptic-naÃ ⁻ ve patients with schizophrenia spectrum disorders. BMC Psychiatry, 2014, 14, 280.	2.6	21
89	Household economic costs associated with mental, neurological and substance use disorders: a cross-sectional survey in six low- and middle-income countries. BJPsych Open, 2019, 5, e34.	0.7	21
90	Community-based rehabilitation intervention for people with schizophrenia in Ethiopia (RISE): results of a 12-month cluster-randomised controlled trial. The Lancet Global Health, 2022, 10, e530-e542.	6.3	21

#	Article	IF	CITATIONS
91	Development and validation of a contextual measure of functioning for people living with severe mental disorders in rural Africa. BMC Psychiatry, 2016, 16, 311.	2.6	20
92	Developing a measure of mental health service satisfaction for use in low income countries: a mixed methods study. BMC Health Services Research, 2017, 17, 183.	2.2	20
93	Ethics in global health research: the need for balance. The Lancet Global Health, 2015, 3, e516-e517.	6.3	19
94	Equitable access to mental healthcare integrated in primary care for people with severe mental disorders in rural Ethiopia: a community-based cross-sectional study. International Journal of Mental Health Systems, 2019, 13, 78.	2.7	19
95	Long-term symptomatic and functional outcome following an intensive inpatient multidisciplinary intervention for treatment-resistant affective disorders. Journal of Affective Disorders, 2014, 166, 334-342.	4.1	18
96	Help seeking and suicidality among people with epilepsy in a rural low income country setting: cross-sectional survey. International Journal of Mental Health Systems, 2017, 11, 44.	2.7	18
97	Functional impairment among people with severe and enduring mental disorder in rural Ethiopia: a cross-sectional study. Social Psychiatry and Psychiatric Epidemiology, 2018, 53, 803-814.	3.1	18
98	Mental health problems and socioeconomic disadvantage: a controlled household study in rural Ethiopia. International Journal for Equity in Health, 2019, 18, 121.	3.5	18
99	Antibacterial and Antifungal Activities of Ethiopian Medicinal Plants: A Systematic Review. Frontiers in Pharmacology, 2021, 12, 633921.	3.5	18
100	Utility of the concept of minor depressive disorder: Evidence from a large rural community sample in a developing country setting. Journal of Affective Disorders, 2007, 104, 111-118.	4.1	17
101	Symptom profile and short term outcome of catatonia: an exploratory clinical study. BMC Psychiatry, 2015, 15, 164.	2.6	17
102	What constitutes problematic khat use? An exploratory mixed methods study in Ethiopia. Substance Abuse Treatment, Prevention, and Policy, 2017, 12, 17.	2.2	17
103	Antenatal predictors of incident and persistent postnatal depressive symptoms in rural Ethiopia: a population-based prospective study. Reproductive Health, 2019, 16, 28.	3.1	17
104	Systematic review of Ethiopian medicinal plants used for their anti-inflammatory and wound healing activities. Journal of Ethnopharmacology, 2021, 276, 114179.	4.1	17
105	Challenges and Opportunities for Drug Discovery in Developing Countries: The Example of Cutaneous Leishmaniasis. ACS Medicinal Chemistry Letters, 2020, 11, 2058-2062.	2.8	16
106	Predictors of carer-burden in schizophrenia: a five-year follow-up study in Butajira, Ethiopia. Ethiopian Medical Journal, 2012, 50, 125-33.	0.6	16
107	Long-Term Impact of Residual Symptoms in Treatment-Resistant Depression. Canadian Journal of Psychiatry, 2011, 56, 549-557.	1.9	15
108	Adjuvant therapy with minocycline for schizophrenia (The MINOS Trial): study protocol for a double-blind randomized placebo-controlled trial. Trials, 2013, 14, 406.	1.6	15

#	Article	IF	CITATIONS
109	Problematic khat use as a possible risk factor for harmful use of other psychoactive substances: a mixed method study in Ethiopia. Substance Abuse Treatment, Prevention, and Policy, 2017, 12, 47.	2.2	15
110	Validity of the Concept of Minor Depression in a Developing Country Setting. Journal of Nervous and Mental Disease, 2008, 196, 22-28.	1.0	14
111	Proposed Training Areas for Global Mental Health Researchers. Academic Psychiatry, 2016, 40, 679-685.	0.9	14
112	Implementing integrated services for people with epilepsy in primary care in Ethiopia: a qualitative study. BMC Health Services Research, 2018, 18, 372.	2.2	14
113	A cross-sectional study to evaluate depression and quality of life among patients with lymphoedema due to podoconiosis, lymphatic filariasis and leprosy. Transactions of the Royal Society of Tropical Medicine and Hygiene, 2020, 114, 983-994.	1.8	14
114	Massachusetts General Hospital Global Psychiatric Clinical Research Training Program: A New Fellowship in Global Mental Health. Academic Psychiatry, 2016, 40, 695-697.	0.9	13
115	EnDPoINT: protocol for an implementation research study to integrate a holistic package of physical health, mental health and psychosocial care for podoconiosis, lymphatic filariasis and leprosy into routine health services in Ethiopia. BMJ Open, 2020, 10, e037675.	1.9	13
116	Psychotherapy Knowledge Translation and Interpersonal Psychotherapy: Using Best-Education Practices to Transform Mental Health Care in Canada and Ethiopia. American Journal of Psychotherapy, 2014, 68, 463-488.	1.2	12
117	Care for bipolar disorder in LMICs needs evidence from local settings. Lancet Psychiatry,the, 2015, 2, 772-773.	7.4	12
118	Stakeholder perspectives on antenatal depression and the potential for psychological intervention in rural Ethiopia: a qualitative study. BMC Pregnancy and Childbirth, 2020, 20, 371.	2.4	12
119	Efficacy and cost-effectiveness of task-shared care for people with severe mental disorders in Ethiopia (TaSCS): a single-blind, randomised, controlled, phase 3 non-inferiority trial. Lancet Psychiatry,the, 2022, 9, 59-71.	7.4	11
120	Rural vs urban residence and experience of discrimination among people with severe mental illnesses in Ethiopia. BMC Psychiatry, 2019, 19, 340.	2.6	10
121	Tuberculosis Prevalence and Predictors Among Health Care-Seeking People Screened for Cough of Any Duration in Ethiopia: A Multicenter Cross-Sectional Study. Frontiers in Public Health, 2021, 9, 805726.	2.7	10
122	"He can send her to her parents― The interaction between marriageability, gender and serious mental illness in rural Ethiopia. BMC Psychiatry, 2019, 19, 315.	2.6	9
123	Definition and Validity of the Construct "Problematic Khat Use†A Systematic Review. European Addiction Research, 2019, 25, 161-172.	2.4	9
124	Impact of integrated mental health care on food insecurity of households of people with severe mental illness in a rural African district: a communityâ€based, controlled before–after study. Tropical Medicine and International Health, 2020, 25, 414-423.	2.3	9
125	Development of an integrated, holistic care package for people with lymphoedema for use at the level of the Primary Health Care Unit in Ethiopia. PLoS Neglected Tropical Diseases, 2021, 15, e0009332.	3.0	9
126	The impact of podoconiosis, lymphatic filariasis, and leprosy on disability and mental well-being: A systematic review. PLoS Neglected Tropical Diseases, 2021, 15, e0009492.	3.0	9

#	Article	IF	CITATIONS
127	Community diagnosis of common skin diseases in the Zay community of the Zeway Islands, Ethiopia. Ethiopian Medical Journal, 2005, 43, 189-95.	0.6	9
128	The effect of gender on the long-term course and outcome of schizophrenia in rural Ethiopia: a population-based cohort. Social Psychiatry and Psychiatric Epidemiology, 2020, 55, 1581-1591.	3.1	8
129	International Clinical Trial Day and clinical trials in Ethiopia and Africa. Trials, 2014, 15, 493.	1.6	7
130	Health Gains and Financial Protection Provided by the Ethiopian Mental Health Strategy: an Extended Cost-Effectiveness Analysis. Health Policy and Planning, 2017, 32, czw134.	2.7	7
131	Functional impairment among people diagnosed with depression in primary healthcare in rural Ethiopia: a comparative cross-sectional study. International Journal of Mental Health Systems, 2019, 13, 50.	2.7	7
132	Food insecurity and work impairment in people with severe mental disorders in a rural district of Ethiopia: a cross-sectional survey. Social Psychiatry and Psychiatric Epidemiology, 2019, 54, 1055-1066.	3.1	7
133	Trauma exposure, depression, suicidal ideation, and alcohol use in people with severe mental disorder in Ethiopia. Social Psychiatry and Psychiatric Epidemiology, 2019, 54, 835-842.	3.1	7
134	Gender-specific experiences of serious mental illness in rural Ethiopia: A qualitative study. Global Public Health, 2020, 15, 185-199.	2.0	7
135	Comorbid mental disorders and quality of life of people with epilepsy attending primary health care clinics in rural Ethiopia. PLoS ONE, 2021, 16, e0238137.	2.5	7
136	Understanding the key processes of excellence as a prerequisite to establishing academic centres of excellence in Africa. BMC Medical Education, 2021, 21, 36.	2.4	7
137	Serum level of high sensitive C-reactive protein and IL â^ 6 markers in patients with treatment-resistant schizophrenia in Ethiopia: a comparative study. BMC Psychiatry, 2021, 21, 428.	2.6	7
138	Antimicrobial susceptibility of bacteria isolated from the infected wounds of patients with lymphoedema in East Wollega, Ethiopia. Transactions of the Royal Society of Tropical Medicine and Hygiene, 2020, 114, 962-973.	1.8	7
139	Economic assessment of a community-based care package for people with lower limb disorder caused by lymphatic filariasis, podoconiosis and leprosy in Ethiopia. Transactions of the Royal Society of Tropical Medicine and Hygiene, 2020, 114, 1021-1034.	1.8	7
140	Prevalence of metabolic syndrome among patients with schizophrenia in Ethiopia. BMC Psychiatry, 2021, 21, 620.	2.6	7
141	Short-term impact of celebrating the international clinical trial day: experience from Ethiopia. Trials, 2017, 18, 332.	1.6	6
142	Health service costs and their association with functional impairment among adults receiving integrated mental health care in five low- and middle-income countries: the PRIME cohort study. Health Policy and Planning, 2020, 35, 567-576.	2.7	6
143	Knowledge, attitudes and practices of health professionals towards people living with lymphoedema caused by lymphatic filariasis, podoconiosis and leprosy in northern Ethiopia. International Health, 2022, 14, 530-536.	2.0	6
144	Comparison of hypothalamo-pituitary-adrenal function in treatment resistant unipolar and bipolar depression. Translational Psychiatry, 2021, 11, 244.	4.8	6

#	Article	IF	CITATIONS
145	"We improved our life because I cut my drinkingâ€: Qualitative analysis of a brief intervention for people with alcohol use disorder in Ethiopian primary health care. Journal of Substance Abuse Treatment, 2022, 132, 108636.	2.8	6
146	The distribution of mental disorders among an isolated island community in southern Ethiopia. Ethiopian Medical Journal, 2005, 43, 71-7.	0.6	6
147	Neuroleptic Malignant Syndrome: Diagnostic and Therapeutic Dilemmas. Behavioural Neurology, 2005, 16, 9-13.	2.1	5
148	Traumatic events and posttraumatic stress disorder in individuals with severe mental illness in a non-western setting: Data from rural Ethiopia Psychological Trauma: Theory, Research, Practice, and Policy, 2021, 13, 684-693.	2.1	5
149	"Medical treatments are also part of God's gift†Holy water attendants' perspectives on a collaboration between spiritual and psychiatric treatment for mental illness in Ethiopia. Transcultural Psychiatry, 2021, 58, 585-599.	1.6	5
150	Multidimensional and intergenerational impact of Severe Mental Disorders. EClinicalMedicine, 2021, 41, 101151.	7.1	5
151	Effect of a Community-Based Holistic Care Package on Physical and Psychosocial Outcomes in People with Lower Limb Disorder Caused by Lymphatic Filariasis, Podoconiosis, and Leprosy in Ethiopia: Results from the EnDPoINT Pilot Cohort Study. American Journal of Tropical Medicine and Hygiene, 2022, 107, 624-631.	1.4	5
152	Conceptualising centres of excellence: a scoping review of global evidence. BMJ Open, 2022, 12, e050419.	1.9	4
153	Understandings of depression among community members and primary healthcare attendees in rural Ethiopia: A qualitative study. Transcultural Psychiatry, 2023, 60, 412-427.	1.6	4
154	The 10-year functional outcome of schizophrenia in Butajira, Ethiopia. Heliyon, 2019, 5, e01272.	3.2	3
155	Development of a psychological intervention for people with bipolar disorder in rural Ethiopia. BJPsych Open, 2021, 7, .	0.7	3
156	Ethiopian medicinal plants used for their anti-inflammatory, wound healing or anti-infective activities: protocol for systematic literature review and meta-analysisEthiopian medicinal plants used for their anti-inflammatory, wound healing or anti-infective activities: protocol for systematic literature review and interative activities protocol for systematic literature seture review and seture activities activities in the systematic literature review and meta-analysisEthiopian medicinal plants used for their anti-inflammatory, wound healing or anti-infective activities: protocol for systematic literature seture review and meta-analysis. BMJ Open Science, 2020, 44, e100064.	1.7	3
157	Towards effective and sustainable global academic partnerships through a maturity model informed by the capability approach. Globalization and Health, 2021, 17, 131.	4.9	3
158	The 11-year functional outcome of bipolar and major depressive disorders in Butajira, Ethiopia. Neurology Psychiatry and Brain Research, 2019, 32, 68-76.	2.0	2
159	Why doesn't God say "enough� Experiences of living with bipolar disorder in rural Ethiopia. Social Science and Medicine, 2021, 270, 113625.	3.8	2
160	Posttraumatic stress disorder intervention for people with severe mental illness in a low-income country primary care setting: a randomized feasibility trial protocol. Pilot and Feasibility Studies, 2021, 7, 149.	1.2	2
161	A qualitative study on the implementation of a holistic care package for control and management of lymphoedema: experience from a pilot intervention in northern Ethiopia. BMC Health Services Research, 2021, 21, 1065.	2.2	2
162	Performance of primary health care workers in detection of mental disorders comorbid with epilepsy in rural Ethiopia. BMC Family Practice, 2021, 22, 204.	2.9	2

#	Article	IF	CITATIONS
163	Exploring the concept of problematic khat use in the Gurage community, South Central Ethiopia: a qualitative study. BMJ Open, 2020, 10, e037907.	1.9	2
164	Psychometric properties of screening questionnaires to detect depression in primary healthcare setting in rural Ethiopia. , 2022, 23, .		2
165	Pathways into and out of homelessness among people with severe mental illness in rural Ethiopia: a qualitative study. BMC Public Health, 2021, 21, 568.	2.9	1
166	Early detection and management of major non-communicable diseases in urban primary healthcare facilities in Ethiopia: a study protocol for a type-3 hybrid implementation-effectiveness design. BMJ Open, 2021, 11, e040564.	1.9	1
167	Unipolar depression. , 2012, , 287-311.		1
168	Validation of the Problematic Khat Use Screening Test: A Cross-Sectional Study. European Addiction Research, 2022, 28, 275-286.	2.4	1
169	Exploring the concept of problematic khat use in the Gurage community, South Central Ethiopia: a qualitative study. BMJ Open, 2020, 10, e037907.	1.9	0