## Abebaw Fekadu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8243487/publications.pdf

Version: 2024-02-01

169 papers 6,478 citations

76326 40 h-index 91884 69 g-index

179 all docs

179 docs citations

179 times ranked

6326 citing authors

#	Article	IF	CITATIONS
1	Understandings of depression among community members and primary healthcare attendees in rural Ethiopia: A qualitative study. Transcultural Psychiatry, 2023, 60, 412-427.	1.6	4
2	Knowledge, attitudes and practices of health professionals towards people living with lymphoedema caused by lymphatic filariasis, podoconiosis and leprosy in northern Ethiopia. International Health, 2022, 14, 530-536.	2.0	6
3	"We improved our life because I cut my drinking― Qualitative analysis of a brief intervention for people with alcohol use disorder in Ethiopian primary health care. Journal of Substance Abuse Treatment, 2022, 132, 108636.	2.8	6
4	Efficacy and cost-effectiveness of task-shared care for people with severe mental disorders in Ethiopia (TaSCS): a single-blind, randomised, controlled, phase 3 non-inferiority trial. Lancet Psychiatry,the, 2022, 9, 59-71.	7.4	11
5	Conceptualising centres of excellence: a scoping review of global evidence. BMJ Open, 2022, 12, e050419.	1.9	4
6	Under detection of depression in primary care settings in low and middle-income countries: a systematic review and meta-analysis. Systematic Reviews, 2022, 11, 21.	5.3	32
7	Validation of the Problematic Khat Use Screening Test: A Cross-Sectional Study. European Addiction Research, 2022, 28, 275-286.	2.4	1
8	Community-based rehabilitation intervention for people with schizophrenia in Ethiopia (RISE): results of a 12-month cluster-randomised controlled trial. The Lancet Global Health, 2022, 10, e530-e542.	6.3	21
9	Psychometric properties of screening questionnaires to detect depression in primary healthcare setting in rural Ethiopia. , 2022, 23, .		2
10	Effect of a Community-Based Holistic Care Package on Physical and Psychosocial Outcomes in People with Lower Limb Disorder Caused by Lymphatic Filariasis, Podoconiosis, and Leprosy in Ethiopia: Results from the EnDPoINT Pilot Cohort Study. American Journal of Tropical Medicine and Hygiene, 2022, 107, 624-631.	1.4	5
11	Why doesn't God say "enough� Experiences of living with bipolar disorder in rural Ethiopia. Social Science and Medicine, 2021, 270, 113625.	3.8	2
12	Comorbid mental disorders and quality of life of people with epilepsy attending primary health care clinics in rural Ethiopia. PLoS ONE, 2021, 16, e0238137.	2.5	7
13	Understanding the key processes of excellence as a prerequisite to establishing academic centres of excellence in Africa. BMC Medical Education, 2021, 21, 36.	2.4	7
14	Antibacterial activity of methanol extracts of the leaves ofÂthree medicinal plants against selected bacteria isolated from wounds of lymphoedema patients. BMC Complementary Medicine and Therapies, 2021, 21, 2.	2.7	38
15	Traumatic events and posttraumatic stress disorder in individuals with severe mental illness in a non-western setting: Data from rural Ethiopia Psychological Trauma: Theory, Research, Practice, and Policy, 2021, 13, 684-693.	2.1	5
16	Pathways into and out of homelessness among people with severe mental illness in rural Ethiopia: a qualitative study. BMC Public Health, 2021, 21, 568.	2.9	1
17	Comparison of hypothalamo-pituitary-adrenal function in treatment resistant unipolar and bipolar depression. Translational Psychiatry, 2021, 11, 244.	4.8	6
18	Development of an integrated, holistic care package for people with lymphoedema for use at the level of the Primary Health Care Unit in Ethiopia. PLoS Neglected Tropical Diseases, 2021, 15, e0009332.	3.0	9

#	Article	IF	CITATIONS
19	Low-coverage sequencing cost-effectively detects known and novel variation in underrepresented populations. American Journal of Human Genetics, 2021, 108, 656-668.	6.2	49
20	"Medical treatments are also part of God's gift― Holy water attendants' perspectives on a collaboration between spiritual and psychiatric treatment for mental illness in Ethiopia. Transcultural Psychiatry, 2021, 58, 585-599.	1.6	5
21	Antibacterial and Antifungal Activities of Ethiopian Medicinal Plants: A Systematic Review. Frontiers in Pharmacology, 2021, 12, 633921.	3.5	18
22	The impact of podoconiosis, lymphatic filariasis, and leprosy on disability and mental well-being: A systematic review. PLoS Neglected Tropical Diseases, 2021, 15, e0009492.	3.0	9
23	Posttraumatic stress disorder intervention for people with severe mental illness in a low-income country primary care setting: a randomized feasibility trial protocol. Pilot and Feasibility Studies, 2021, 7, 149.	1.2	2
24	Systematic review of Ethiopian medicinal plants used for their anti-inflammatory and wound healing activities. Journal of Ethnopharmacology, 2021, 276, 114179.	4.1	17
25	The potential use of digital health technologies in the African context: a systematic review of evidence from Ethiopia. Npj Digital Medicine, 2021, 4, 125.	10.9	54
26	Serum level of high sensitive C-reactive protein and IL â^ 6 markers in patients with treatment-resistant schizophrenia in Ethiopia: a comparative study. BMC Psychiatry, 2021, 21, 428.	2.6	7
27	Development of a psychological intervention for people with bipolar disorder in rural Ethiopia. BJPsych Open, 2021, 7, .	0.7	3
28	Multidimensional and intergenerational impact of Severe Mental Disorders. EClinicalMedicine, 2021, 41, 101151.	7.1	5
29	Early detection and management of major non-communicable diseases in urban primary healthcare facilities in Ethiopia: a study protocol for a type-3 hybrid implementation-effectiveness design. BMJ Open, 2021, 11, e040564.	1.9	1
30	A qualitative study on the implementation of a holistic care package for control and management of lymphoedema: experience from a pilot intervention in northern Ethiopia. BMC Health Services Research, 2021, 21, 1065.	2.2	2
31	Performance of primary health care workers in detection of mental disorders comorbid with epilepsy in rural Ethiopia. BMC Family Practice, 2021, 22, 204.	2.9	2
32	Towards effective and sustainable global academic partnerships through a maturity model informed by the capability approach. Globalization and Health, 2021, 17, 131.	4.9	3
33	Prevalence of metabolic syndrome among patients with schizophrenia in Ethiopia. BMC Psychiatry, 2021, 21, 620.	2.6	7
34	Tuberculosis Prevalence and Predictors Among Health Care-Seeking People Screened for Cough of Any Duration in Ethiopia: A Multicenter Cross-Sectional Study. Frontiers in Public Health, 2021, 9, 805726.	2.7	10
35	Adapting and pre-testing the World Health Organization's Caregiver Skills Training programme for autism and other developmental disorders in a very low-resource setting: Findings from Ethiopia. Autism, 2020, 24, 51-63.	4.1	48
36	Gender-specific experiences of serious mental illness in rural Ethiopia: A qualitative study. Global Public Health, 2020, 15, 185-199.	2.0	7

#	Article	IF	CITATIONS
37	Explanatory models of depression in sub-Saharan Africa: Synthesis of qualitative evidence. Social Science and Medicine, 2020, 246, 112760.	3.8	40
38	Developing a Theory of Change model of service user and caregiver involvement in mental health system strengthening in primary health care in rural Ethiopia. International Journal of Mental Health Systems, 2020, 14, 51.	2.7	46
39	Challenges and Opportunities for Drug Discovery in Developing Countries: The Example of Cutaneous Leishmaniasis. ACS Medicinal Chemistry Letters, 2020, 11, 2058-2062.	2.8	16
40	Containment of COVID-19 in Ethiopia and implications for tuberculosis care and research. Infectious Diseases of Poverty, 2020, 9, 131.	3.7	76
41	Electronic pillbox-enabled self-administered therapy versus standard directly observed therapy for tuberculosis medication adherence and treatment outcomes in Ethiopia (SELFTB): protocol for a multicenter randomized controlled trial. Trials, 2020, 21, 383.	1.6	30
42	The fight to end tuberculosis must not be forgotten in the COVID-19 outbreak. Nature Medicine, 2020, 26, 811-812.	30.7	34
43	Burden of tuberculosis and challenges related to screening and diagnosis in Ethiopia. Journal of Clinical Tuberculosis and Other Mycobacterial Diseases, 2020, 19, 100158.	1.3	27
44	Health service costs and their association with functional impairment among adults receiving integrated mental health care in five low- and middle-income countries: the PRIME cohort study. Health Policy and Planning, 2020, 35, 567-576.	2.7	6
45	Stakeholder perspectives on antenatal depression and the potential for psychological intervention in rural Ethiopia: a qualitative study. BMC Pregnancy and Childbirth, 2020, 20, 371.	2.4	12
46	Impact of integrated mental health care on food insecurity of households of people with severe mental illness in a rural African district: a communityâ€based, controlled before–after study. Tropical Medicine and International Health, 2020, 25, 414-423.	2.3	9
47	The effect of gender on the long-term course and outcome of schizophrenia in rural Ethiopia: a population-based cohort. Social Psychiatry and Psychiatric Epidemiology, 2020, 55, 1581-1591.	3.1	8
48	Antimicrobial susceptibility of bacteria isolated from the infected wounds of patients with lymphoedema in East Wollega, Ethiopia. Transactions of the Royal Society of Tropical Medicine and Hygiene, 2020, 114, 962-973.	1.8	7
49	EnDPoINT: protocol for an implementation research study to integrate a holistic package of physical health, mental health and psychosocial care for podoconiosis, lymphatic filariasis and leprosy into routine health services in Ethiopia. BMJ Open, 2020, 10, e037675.	1.9	13
50	Ethiopian medicinal plants used for their anti-inflammatory, wound healing or anti-infective activities: protocol for systematic literature review and meta-analysisEthiopian medicinal plants used for their anti-inflammatory, wound healing or anti-infective activities: protocol for systematic literature review and meta-analysis. BMJ Open Science, 2020, 44, e100064.	1.7	3
51	Economic assessment of a community-based care package for people with lower limb disorder caused by lymphatic filariasis, podoconiosis and leprosy in Ethiopia. Transactions of the Royal Society of Tropical Medicine and Hygiene, 2020, 114, 1021-1034.	1.8	7
52	A cross-sectional study to evaluate depression and quality of life among patients with lymphoedema due to podoconiosis, lymphatic filariasis and leprosy. Transactions of the Royal Society of Tropical Medicine and Hygiene, 2020, 114, 983-994.	1.8	14
53	Exploring the concept of problematic khat use in the Gurage community, South Central Ethiopia: a qualitative study. BMJ Open, 2020, 10, e037907.	1.9	0
54	Exploring the concept of problematic khat use in the Gurage community, South Central Ethiopia: a qualitative study. BMJ Open, 2020, 10, e037907.	1.9	2

#	Article	IF	CITATIONS
55	Mental health problems and socioeconomic disadvantage: a controlled household study in rural Ethiopia. International Journal for Equity in Health, 2019, 18, 121.	3.5	18
56	Catastrophic health expenditure and impoverishment in households of persons with depression: a cross-sectional, comparative study in rural Ethiopia. BMC Public Health, 2019, 19, 930.	2.9	44
57	Functional impairment among people diagnosed with depression in primary healthcare in rural Ethiopia: a comparative cross-sectional study. International Journal of Mental Health Systems, 2019, 13, 50.	2.7	7
58	Burnout among primary healthcare workers during implementation of integrated mental healthcare in rural Ethiopia: a cohort study. Human Resources for Health, 2019, 17, 58.	3.1	26
59	Homelessness and severe mental illness in low- and middle-income countries: scoping review. BJPsych Open, 2019, 5, e57.	0.7	39
60	The psychosis treatment gap and its consequences in rural Ethiopia. BMC Psychiatry, 2019, 19, 325.	2.6	23
61	"He can send her to her parents†The interaction between marriageability, gender and serious mental illness in rural Ethiopia. BMC Psychiatry, 2019, 19, 315.	2.6	9
62	Household economic costs associated with mental, neurological and substance use disorders: a cross-sectional survey in six low- and middle-income countries. BJPsych Open, 2019, 5, e34.	0.7	21
63	Scaling up integrated primary mental health in six low- and middle-income countries: obstacles, synergies and implications for systems reform. BJPsych Open, 2019, 5, e69.	0.7	51
64	Treatment gap, help-seeking, stigma and magnitude of alcohol use disorder in rural Ethiopia. Substance Abuse Treatment, Prevention, and Policy, 2019, 14, 4.	2.2	58
65	Catastrophic out-of-pocket payments for households of people with severe mental disorder: a comparative study in rural Ethiopia. International Journal of Mental Health Systems, 2019, 13, 39.	2.7	23
66	The 11-year functional outcome of bipolar and major depressive disorders in Butajira, Ethiopia. Neurology Psychiatry and Brain Research, 2019, 32, 68-76.	2.0	2
67	The 10-year functional outcome of schizophrenia in Butajira, Ethiopia. Heliyon, 2019, 5, e01272.	3.2	3
68	Food insecurity and work impairment in people with severe mental disorders in a rural district of Ethiopia: a cross-sectional survey. Social Psychiatry and Psychiatric Epidemiology, 2019, 54, 1055-1066.	3.1	7
69	Definition and Validity of the Construct "Problematic Khat Use― A Systematic Review. European Addiction Research, 2019, 25, 161-172.	2.4	9
70	Antenatal predictors of incident and persistent postnatal depressive symptoms in rural Ethiopia: a population-based prospective study. Reproductive Health, 2019, 16, 28.	3.1	17
71	Trauma exposure, depression, suicidal ideation, and alcohol use in people with severe mental disorder in Ethiopia. Social Psychiatry and Psychiatric Epidemiology, 2019, 54, 835-842.	3.1	7
72	Equitable access to mental healthcare integrated in primary care for people with severe mental disorders in rural Ethiopia: a community-based cross-sectional study. International Journal of Mental Health Systems, 2019, 13, 78.	2.7	19

#	Article	IF	CITATIONS
73	Multidimensional impact of severe mental illness on family members: systematic review. BMJ Open, 2019, 9, e032391.	1.9	49
74	Rural vs urban residence and experience of discrimination among people with severe mental illnesses in Ethiopia. BMC Psychiatry, 2019, 19, 340.	2.6	10
75	Partnerships in a Global Mental Health Research Programme—the Example of PRIME. Global Social Welfare, 2019, 6, 159-175.	1.9	24
76	Ethiopian community health workers' beliefs and attitudes towards children with autism: Impact of a brief training intervention. Autism, 2019, 23, 39-49.	4.1	27
77	Coping strategies of women with postpartum depression symptoms in rural Ethiopia: a cross-sectional community study. BMC Psychiatry, 2018, 18, 41.	2.6	43
78	Global mental health and schizophrenia. Current Opinion in Psychiatry, 2018, 31, 193-199.	6.3	27
79	Standardisation framework for the Maudsley staging method for treatment resistance in depression. BMC Psychiatry, 2018, 18, 100.	2.6	64
80	Detection and treatment initiation for depression and alcohol use disorders: facility-based cross-sectional studies in five low-income and middle-income country districts. BMJ Open, 2018, 8, e023421.	1.9	45
81	Psychological interventions for bipolar disorder in low- and middle-income countries: systematic review. BJPsych Open, 2018, 4, 375-384.	0.7	26
82	Functional impairment among people with severe and enduring mental disorder in rural Ethiopia: a cross-sectional study. Social Psychiatry and Psychiatric Epidemiology, 2018, 53, 803-814.	3.1	18
83	Implementing integrated services for people with epilepsy in primary care in Ethiopia: a qualitative study. BMC Health Services Research, 2018, 18, 372.	2.2	14
84	Community-based rehabilitation intervention for people with schizophrenia in Ethiopia (RISE): a 12 month mixed methods pilot study. BMC Psychiatry, 2018, 18, 250.	2.6	63
85	Postpartum depressive symptoms in the context of high social adversity and reproductive health threats: a population-based study. International Journal of Mental Health Systems, 2018, 12, 42.	2.7	29
86	A descriptive analysis of depression and pain complaints among patients with cancer in a low income country. PLoS ONE, 2018, 13, e0193713.	2.5	22
87	Health Gains and Financial Protection Provided by the Ethiopian Mental Health Strategy: an Extended Cost-Effectiveness Analysis. Health Policy and Planning, 2017, 32, czw134.	2.7	7
88	Developing a measure of mental health service satisfaction for use in low income countries: a mixed methods study. BMC Health Services Research, 2017, 17, 183.	2.2	20
89	Training needs and perspectives of community health workers in relation to integrating child mental health care into primary health care in a rural setting in sub-Saharan Africa: a mixed methods study. International Journal of Mental Health Systems, 2017, 11, 15.	2.7	31
90	What constitutes problematic khat use? An exploratory mixed methods study in Ethiopia. Substance Abuse Treatment, Prevention, and Policy, 2017, 12, 17.	2.2	17

#	Article	IF	CITATIONS
91	The path to longer and healthier lives for all Africans by 2030: the Lancet Commission on the future of health in sub-Saharan Africa. Lancet, The, 2017, 390, 2803-2859.	13.7	150
92	Antenatal depressive symptoms and utilisation of delivery and postnatal care: a prospective study in rural Ethiopia. BMC Pregnancy and Childbirth, 2017, 17, 206.	2.4	22
93	Recognition of depression by primary care clinicians in rural Ethiopia. BMC Family Practice, 2017, 18, 56.	2.9	36
94	Conceptualisation of job-related wellbeing, stress and burnout among healthcare workers in rural Ethiopia: a qualitative study. BMC Health Services Research, 2017, 17, 412.	2.2	46
95	Validation of the World Health Organization Disability Assessment Schedule in people with severe mental disorders in rural Ethiopia. Health and Quality of Life Outcomes, 2017, 15, 64.	2.4	64
96	Health system governance to support scale up of mental health care in Ethiopia: a qualitative study. International Journal of Mental Health Systems, 2017, 11, 38.	2.7	34
97	Engaging and staying engaged: a phenomenological study of barriers to equitable access to mental healthcare for people with severe mental disorders in a rural African setting. International Journal for Equity in Health, 2017, 16, 156.	3.5	33
98	"l cry every day and night, I have my son tied in chains†physical restraint of people with schizophrenia in community settings in Ethiopia. Globalization and Health, 2017, 13, 47.	4.9	78
99	Problematic khat use as a possible risk factor for harmful use of other psychoactive substances: a mixed method study in Ethiopia. Substance Abuse Treatment, Prevention, and Policy, 2017, 12, 47.	2.2	15
100	Help seeking and suicidality among people with epilepsy in a rural low income country setting: cross-sectional survey. International Journal of Mental Health Systems, 2017, 11, 44.	2.7	18
101	Short-term impact of celebrating the international clinical trial day: experience from Ethiopia. Trials, 2017, 18, 332.	1.6	6
102	Antenatal depressive symptoms and perinatal complications: a prospective study in rural Ethiopia. BMC Psychiatry, 2017, 17, 301.	2.6	40
103	Treatment gap and help-seeking for postpartum depression in a rural African setting. BMC Psychiatry, 2016, 16, 196.	2.6	61
104	Community-based Rehabilitation Intervention for people with Schizophrenia in Ethiopia (RISE): study protocol for a cluster randomised controlled trial. Trials, 2016, 17, 299.	1.6	34
105	Development and validation of a contextual measure of functioning for people living with severe mental disorders in rural Africa. BMC Psychiatry, 2016, 16, 311.	2.6	20
106	Antenatal depressive symptoms and maternal health care utilisation: a population-based study of pregnant women in Ethiopia. BMC Pregnancy and Childbirth, 2016, 16, 301.	2.4	46
107	Equitable access to integrated primary mental healthcare for people with severe mental disorders in Ethiopia: a formative study. International Journal for Equity in Health, 2016, 15, 121.	3.5	24
108	Development of a scalable mental healthcare plan for a rural district in Ethiopia. British Journal of Psychiatry, 2016, 208, s4-s12.	2.8	93

#	Article	IF	Citations
109	District mental healthcare plans for five low-and middle-income countries: Commonalities, variations and evidence gaps. British Journal of Psychiatry, 2016, 208, s47-s54.	2.8	73
110	Planning and evaluating mental health services in low- and middle-income countries using theory of change. British Journal of Psychiatry, 2016, 208, s55-s62.	2.8	69
111	Estimating the cost of implementing district mental healthcare plans in five low- and middle-income countries: The PRIME study. British Journal of Psychiatry, 2016, 208, s71-s78.	2.8	40
112	Building the capacity of policy-makers and planners to strengthen mental health systems in low- and middle-income countries: a systematic review. BMC Health Services Research, 2016, 16, 601.	2.2	22
113	Task sharing for the care of severe mental disorders in a low-income country (TaSCS): study protocol for a randomised, controlled, non-inferiority trial. Trials, 2016, 17, 76.	1.6	39
114	Hazardous alcohol use and associated factors in a rural Ethiopian district: a cross-sectional community survey. BMC Public Health, 2016, 16, 218.	2.9	46
115	Proposed Training Areas for Global Mental Health Researchers. Academic Psychiatry, 2016, 40, 679-685.	0.9	14
116	Evaluation of district mental healthcare plans: The PRIME consortium methodology. British Journal of Psychiatry, 2016, 208, s63-s70.	2.8	92
117	Non-fatal suicidal behaviour in rural Ethiopia: a cross-sectional facility- and population-based study. BMC Psychiatry, 2016, 16, 75.	2.6	31
118	Stigma, explanatory models and unmet needs of caregivers of children with developmental disorders in a low-income African country: a cross-sectional facility-based survey. BMC Health Services Research, 2016, 16, 152.	2.2	119
119	Scaling-up essential neuropsychiatric services in Ethiopia: a cost-effectiveness analysis. Health Policy and Planning, 2016, 31, 504-513.	2.7	27
120	Massachusetts General Hospital Global Psychiatric Clinical Research Training Program: A New Fellowship in Global Mental Health. Academic Psychiatry, 2016, 40, 695-697.	0.9	13
121	Participatory planning of a primary care service for people with severe mental disorders in rural Ethiopia. Health Policy and Planning, 2016, 31, 367-376.	2.7	55
122	Treatment Contact Coverage for Probable Depressive and Probable Alcohol Use Disorders in Four Low- and Middle-Income Country Districts: The PRIME Cross-Sectional Community Surveys. PLoS ONE, 2016, 11, e0162038.	2.5	62
123	Care for bipolar disorder in LMICs needs evidence from local settings. Lancet Psychiatry,the, 2015, 2, 772-773.	7.4	12
124	Developing a mental health care plan in a low resource setting: the theory of change approach. BMC Health Services Research, 2015, 15, 429.	2.2	33
125	Stakeholder analysis of the Programme for Improving Mental health carE (PRIME): baseline findings. International Journal of Mental Health Systems, 2015, 9, 27.	2.7	50
126	Ethics in global health research: the need for balance. The Lancet Global Health, 2015, 3, e516-e517.	6.3	19

#	Article	IF	Citations
127	Long-term clinical course and outcome of schizophrenia in rural Ethiopia: 10-year follow-up of a population-based cohort. Schizophrenia Research, 2015, 161, 414-420.	2.0	27
128	Excess mortality in severe mental illness: 10-year population-based cohort study in rural Ethiopia. British Journal of Psychiatry, 2015, 206, 289-296.	2.8	130
129	Validity of brief screening questionnaires to detect depression in primary care in Ethiopia. Journal of Affective Disorders, 2015, 186, 32-39.	4.1	131
130	Symptom profile and short term outcome of catatonia: an exploratory clinical study. BMC Psychiatry, 2015, 15, 164.	2.6	17
131	Beyond the Biomedical: Community Resources for Mental Health Care in Rural Ethiopia. PLoS ONE, 2015, 10, e0126666.	2.5	39
132	Development of a Community-Based Rehabilitation Intervention for People with Schizophrenia in Ethiopia. PLoS ONE, 2015, 10, e0143572.	2.5	46
133	Psychotherapy Knowledge Translation and Interpersonal Psychotherapy: Using Best-Education Practices to Transform Mental Health Care in Canada and Ethiopia. American Journal of Psychotherapy, 2014, 68, 463-488.	1.2	12
134	Movement disorders in neuroleptic-naÃ-ve patients with schizophrenia spectrum disorders. BMC Psychiatry, 2014, 14, 280.	2.6	21
135	International Clinical Trial Day and clinical trials in Ethiopia and Africa. Trials, 2014, 15, 493.	1.6	7
136	Population level mental distress in rural Ethiopia. BMC Psychiatry, 2014, 14, 194.	2.6	98
137	Violence and violent victimization in people with severe mental illness in a rural low-income country setting: A comparative cross-sectional community study. Schizophrenia Research, 2014, 152, 275-282.	2.0	39
138	Acceptability and feasibility of using non-specialist health workers to deliver mental health care: Stakeholder perceptions from the PRIME district sites in Ethiopia, India, Nepal, South Africa, and Uganda. Social Science and Medicine, 2014, 118, 33-42.	3.8	259
139	Burden of mental disorders and unmet needs among street homeless people in Addis Ababa, Ethiopia. BMC Medicine, 2014, 12, 138.	5 <b>.</b> 5	47
140	Suicide and suicide attempts in people with severe mental disorders in Butajira, Ethiopia: 10Âyear follow-up of a population-based cohort. BMC Psychiatry, 2014, 14, 150.	2.6	51
141	Using workshops to develop theories of change in five low and middle income countries: lessons from the programme for improving mental health care (PRIME). International Journal of Mental Health Systems, 2014, 8, 15.	2.7	60
142	The impact of childhood adversity on suicidality and clinical course in treatment-resistant depression. Journal of Affective Disorders, 2014, 152-154, 122-130.	4.1	132
143	Long-term symptomatic and functional outcome following an intensive inpatient multidisciplinary intervention for treatment-resistant affective disorders. Journal of Affective Disorders, 2014, 166, 334-342.	4.1	18
144	Global mental health: perspectives from Ethiopia. Global Health Action, 2014, 7, 25447.	1.9	34

#	Article	IF	Citations
145	Challenges and Opportunities for Implementing Integrated Mental Health Care: A District Level Situation Analysis from Five Low- and Middle-Income Countries. PLoS ONE, 2014, 9, e88437.	2.5	255
146	Longitudinal course of symptom severity and fluctuation in patients with treatment-resistant unipolar and bipolar depression. Psychiatry Research, 2013, 207, 143-149.	3.3	34
147	Adjuvant therapy with minocycline for schizophrenia (The MINOS Trial): study protocol for a double-blind randomized placebo-controlled trial. Trials, 2013, 14, 406.	1.6	15
148	PRIME: A Programme to Reduce the Treatment Gap for Mental Disorders in Five Low- and Middle-Income Countries. PLoS Medicine, 2012, 9, e1001359.	8.4	436
149	Prediction of longer-term outcome of treatment-resistant depression in tertiary care. British Journal of Psychiatry, 2012, 201, 369-375.	2.8	65
150	Five-year clinical course and outcome of schizophrenia in Ethiopia. Schizophrenia Research, 2012, 136, 137-142.	2.0	23
151	Internalized stigma among patients with schizophrenia in Ethiopia: a cross-sectional facility-based study. BMC Psychiatry, 2012, 12, 239.	2.6	89
152	Unipolar depression., 2012,, 287-311.		1
153	Predictors of carer-burden in schizophrenia: a five-year follow-up study in Butajira, Ethiopia. Ethiopian Medical Journal, 2012, 50, 125-33.	0.6	16
154	Long-Term Impact of Residual Symptoms in Treatment-Resistant Depression. Canadian Journal of Psychiatry, 2011, 56, 549-557.	1.9	15
155	Prospective evaluation of specialist inpatient treatment for refractory affective disorders. Journal of Affective Disorders, 2011, 131, 92-103.	4.1	25
156	Five-year mortality in a cohort of people with schizophrenia in Ethiopia. BMC Psychiatry, 2011, 11, 165.	2.6	69
157	Trimethoprim as Adjuvant Treatment in Schizophrenia: A Double-Blind, Randomized, Placebo-Controlled Clinical Trial. Schizophrenia Bulletin, 2010, 36, 846-851.	4.3	33
158	Toxoplasmosis as a cause for behaviour disorders - overview of evidence and mechanisms. Folia Parasitologica, 2010, 57, 105-113.	1.3	117
159	Clinical Course and Outcome of Schizophrenia in a Predominantly Treatment-Naive Cohort in Rural Ethiopia. Schizophrenia Bulletin, 2009, 35, 646-654.	4.3	103
160	What happens to patients with treatment-resistant depression? A systematic review of medium to long term outcome studies. Journal of Affective Disorders, 2009, 116, 4-11.	4.1	244
161	A Multidimensional Tool to Quantify Treatment Resistance in Depression. Journal of Clinical Psychiatry, 2009, 70, 177-184.	2.2	211
162	The Maudsley Staging Method for Treatment-Resistant Depression. Journal of Clinical Psychiatry, 2009, 70, 952-957.	2.2	99

#	Article	IF	CITATION
163	Validity of the Concept of Minor Depression in a Developing Country Setting. Journal of Nervous and Mental Disease, 2008, 196, 22-28.	1.0	14
164	Utility of the concept of minor depressive disorder: Evidence from a large rural community sample in a developing country setting. Journal of Affective Disorders, 2007, 104, 111-118.	4.1	17
165	Symptomatic and functional outcome of bipolar disorder in Butajira, Ethiopia. Journal of Affective Disorders, 2006, 90, 239-249.	4.1	41
166	Neuroleptic Malignant Syndrome: Diagnostic and Therapeutic Dilemmas. Behavioural Neurology, 2005, 16, 9-13.	2.1	5
167	The distribution of mental disorders among an isolated island community in southern Ethiopia. Ethiopian Medical Journal, 2005, 43, 71-7.	0.6	6
168	Community diagnosis of common skin diseases in the Zay community of the Zeway Islands, Ethiopia. Ethiopian Medical Journal, 2005, 43, 189-95.	0.6	9
169	Bipolar disorder among an isolated island community in Ethiopia. Journal of Affective Disorders, 2004, 80, 1-10.	4.1	28