

Karin Jirström

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8261944/publications.pdf>

Version: 2024-02-01

222
papers

12,878
citations

26630

56
h-index

32842

100
g-index

226
all docs

226
docs citations

226
times ranked

22048
citing authors

#	ARTICLE	IF	CITATIONS
1	Leukocyte Complexity Predicts Breast Cancer Survival and Functionally Regulates Response to Chemotherapy. <i>Cancer Discovery</i> , 2011, 1, 54-67.	9.4	1,486
2	Tertiary lymphoid structures improve immunotherapy and survival in melanoma. <i>Nature</i> , 2020, 577, 561-565.	27.8	1,209
3	Interleukin-6 as a Therapeutic Target in Human Ovarian Cancer. <i>Clinical Cancer Research</i> , 2011, 17, 6083-6096.	7.0	330
4	Isolation and Characterization of Progenitor-Like Cells from Human Renal Proximal Tubules. <i>American Journal of Pathology</i> , 2011, 178, 828-837.	3.8	231
5	Prognostic Significance of Stromal Platelet-Derived Growth Factor β 2-Receptor Expression in Human Breast Cancer. <i>American Journal of Pathology</i> , 2009, 175, 334-341.	3.8	215
6	SATB2 in Combination With Cytokeratin 20 Identifies Over 95% of all Colorectal Carcinomas. <i>American Journal of Surgical Pathology</i> , 2011, 35, 937-948.	3.7	209
7	Prognostic impact of tumour-infiltrating B cells and plasma cells in colorectal cancer. <i>International Journal of Cancer</i> , 2016, 139, 1129-1139.	5.1	192
8	Nuclear expression of the non-B-cell lineage Sox11 transcription factor identifies mantle cell lymphoma. <i>Blood</i> , 2008, 111, 800-805.	1.4	185
9	Effect of Radiotherapy After Breast-Conserving Surgery for Ductal Carcinoma in Situ: 20 Years Follow-Up in the Randomized SweDCIS Trial. <i>Journal of Clinical Oncology</i> , 2014, 32, 3613-3618.	1.6	184
10	Targeting HMG-CoA reductase with statins in a window-of-opportunity breast cancer trial. <i>Breast Cancer Research and Treatment</i> , 2013, 138, 499-508.	2.5	183
11	Association Between Pak1 Expression and Subcellular Localization and Tamoxifen Resistance in Breast Cancer Patients. <i>Journal of the National Cancer Institute</i> , 2006, 98, 671-680.	6.3	177
12	Human tumors instigate granulysin-expressing hematopoietic cells that promote malignancy by activating stromal fibroblasts in mice. <i>Journal of Clinical Investigation</i> , 2011, 121, 784-799.	8.2	177
13	Molecular stratification of metastatic melanoma using gene expression profiling : Prediction of survival outcome and benefit from molecular targeted therapy. <i>Oncotarget</i> , 2015, 6, 12297-12309.	1.8	148
14	STC1 Expression By Cancer-Associated Fibroblasts Drives Metastasis of Colorectal Cancer. <i>Cancer Research</i> , 2013, 73, 1287-1297.	0.9	144
15	Integration of genomic, transcriptomic and proteomic data identifies two biologically distinct subtypes of invasive lobular breast cancer. <i>Scientific Reports</i> , 2016, 6, 18517.	3.3	143
16	Cancer-associated fibroblast-secreted CXCL16 attracts monocytes to promote stroma activation in triple-negative breast cancers. <i>Nature Communications</i> , 2016, 7, 13050.	12.8	135
17	Hypoxia-Inducible Factor-2 β Correlates to Distant Recurrence and Poor Outcome in Invasive Breast Cancer. <i>Cancer Research</i> , 2008, 68, 9212-9220.	0.9	130
18	Increased claudin-4 expression is associated with poor prognosis and high tumour grade in breast cancer. <i>International Journal of Cancer</i> , 2009, 124, 2088-2097.	5.1	128

#	ARTICLE	IF	CITATIONS
19	CA IX is an Independent Prognostic Marker in Premenopausal Breast Cancer Patients with One to Three Positive Lymph Nodes and a Putative Marker of Radiation Resistance. <i>Clinical Cancer Research</i> , 2006, 12, 6421-6431.	7.0	123
20	High Progesterone Receptor Expression Correlates to the Effect of Adjuvant Tamoxifen in Premenopausal Breast Cancer Patients. <i>Clinical Cancer Research</i> , 2006, 12, 4614-4618.	7.0	121
21	Expression of the cytoskeleton linker protein ezrin in human cancers. <i>Clinical and Experimental Metastasis</i> , 2007, 24, 69-78.	3.3	118
22	Adverse Effect of Adjuvant Tamoxifen in Premenopausal Breast Cancer with Cyclin D1 Gene Amplification. <i>Cancer Research</i> , 2005, 65, 8009-8016.	0.9	117
23	JAM α expression positively correlates with poor prognosis in breast cancer patients. <i>International Journal of Cancer</i> , 2009, 125, 1343-1351.	5.1	115
24	ERK phosphorylation is linked to VEGFR2 expression and Ets-2 phosphorylation in breast cancer and is associated with tamoxifen treatment resistance and small tumours with good prognosis. <i>Oncogene</i> , 2005, 24, 4370-4379.	5.9	106
25	Classification of Breast Cancer Using Genetic Algorithms and Tissue Microarrays. <i>Clinical Cancer Research</i> , 2006, 12, 6459-6468.	7.0	100
26	Wnt5a Induces a Tolerogenic Phenotype of Macrophages in Sepsis and Breast Cancer Patients. <i>Journal of Immunology</i> , 2012, 188, 5448-5458.	0.8	100
27	Heterogeneity of Colorectal Cancer Risk Factors by Anatomical Subsite in 10 European Countries: A Multinational Cohort Study. <i>Clinical Gastroenterology and Hepatology</i> , 2019, 17, 1323-1331.e6.	4.4	99
28	CENP-F expression is associated with poor prognosis and chromosomal instability in patients with primary breast cancer. <i>International Journal of Cancer</i> , 2007, 120, 1434-1443.	5.1	98
29	Molecular Profiling Reveals Low- and High-Grade Forms of Primary Melanoma. <i>Clinical Cancer Research</i> , 2012, 18, 4026-4036.	7.0	96
30	A Biological Signature for Breast Ductal Carcinoma <i>In Situ</i> to Predict Radiotherapy Benefit and Assess Recurrence Risk. <i>Clinical Cancer Research</i> , 2018, 24, 5895-5901.	7.0	88
31	Forkhead Box F1 Regulates Tumor-Promoting Properties of Cancer-Associated Fibroblasts in Lung Cancer. <i>Cancer Research</i> , 2010, 70, 2644-2654.	0.9	84
32	Altered Cytoplasmic-to-Nuclear Ratio of Survivin Is a Prognostic Indicator in Breast Cancer. <i>Clinical Cancer Research</i> , 2008, 14, 2681-2689.	7.0	83
33	Aberrantly activated claudin 6 and 18.2 as potential therapy targets in non-small cell lung cancer. <i>International Journal of Cancer</i> , 2014, 135, 2206-2214.	5.1	82
34	ANLN is a prognostic biomarker independent of Ki-67 and essential for cell cycle progression in primary breast cancer. <i>BMC Cancer</i> , 2016, 16, 904.	2.6	82
35	A Prospective Evaluation of Early Detection Biomarkers for Ovarian Cancer in the European EPIC Cohort. <i>Clinical Cancer Research</i> , 2016, 22, 4664-4675.	7.0	80
36	Therapeutic Rationale to Target Highly Expressed CDK7 Conferring Poor Outcomes in Triple-Negative Breast Cancer. <i>Cancer Research</i> , 2017, 77, 3834-3845.	0.9	79

#	ARTICLE	IF	CITATIONS
37	High expression of cyclin D1 is associated to high proliferation rate and increased risk of mortality in women with ER-positive but not in ER-negative breast cancers. <i>Breast Cancer Research and Treatment</i> , 2017, 164, 667-678.	2.5	79
38	Polymorphisms in fatty acid metabolism-related genes are associated with colorectal cancer risk. <i>Carcinogenesis</i> , 2010, 31, 466-472.	2.8	77
39	Downregulation of miR-92a Is Associated with Aggressive Breast Cancer Features and Increased Tumour Macrophage Infiltration. <i>PLoS ONE</i> , 2012, 7, e36051.	2.5	76
40	Prognostic impact of tumour-associated B cells and plasma cells in epithelial ovarian cancer. <i>Journal of Ovarian Research</i> , 2016, 9, 21.	3.0	76
41	miR-187 Is an Independent Prognostic Factor in Breast Cancer and Confers Increased Invasive Potential <i>In Vitro</i> . <i>Clinical Cancer Research</i> , 2012, 18, 6702-6713.	7.0	75
42	Tumor characteristics and prognosis in women with pregnancy-associated breast cancer. <i>International Journal of Cancer</i> , 2018, 142, 1343-1354.	5.1	75
43	The Prognostic Significance of Tryptophanyl-tRNA Synthetase in Colorectal Cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009, 18, 2949-2956.	2.5	71
44	Hypoxia inducible factor-1 α is a prognostic marker in premenopausal patients with intermediate to highly differentiated breast cancer but not a predictive marker for tamoxifen response. <i>International Journal of Cancer</i> , 2006, 118, 2609-2616.	5.1	70
45	The Impact of the Fourth Edition of the WHO Classification of Lung Tumours on Histological Classification of Resected Pulmonary NSCCs. <i>Journal of Thoracic Oncology</i> , 2016, 11, 862-872.	1.1	70
46	Nuclear expression of the RNA-binding protein RBM3 is associated with an improved clinical outcome in breast cancer. <i>Modern Pathology</i> , 2009, 22, 1564-1574.	5.5	69
47	Epidermal growth factor receptor and vascular endothelial growth factor receptor 2 are specific biomarkers in triple-negative breast cancer. Results from a controlled randomized trial with long-term follow-up. <i>Breast Cancer Research and Treatment</i> , 2010, 120, 491-498.	2.5	69
48	Anthropometric measures and epithelial ovarian cancer risk in the European Prospective Investigation into Cancer and Nutrition. <i>International Journal of Cancer</i> , 2010, 126, 2404-2415.	5.1	68
49	Pre-diagnostic concordance with the WCRF/AICR guidelines and survival in European colorectal cancer patients: a cohort study. <i>BMC Medicine</i> , 2015, 13, 107.	5.5	66
50	Immune effector monocyte-neutrophil cooperation induced by the primary tumor prevents metastatic progression of breast cancer. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 21704-21714.	7.1	66
51	The clinical impact of tumour-infiltrating lymphocytes in colorectal cancer differs by anatomical subsite: A cohort study. <i>International Journal of Cancer</i> , 2017, 141, 1654-1666.	5.1	65
52	Combined Androgen and Estrogen Receptor Status in Breast Cancer: Treatment Prediction and Prognosis in a Population-Based Prospective Cohort. <i>Clinical Cancer Research</i> , 2015, 21, 3640-3650.	7.0	64
53	Prognostic impact of tumour-associated B cells and plasma cells in oesophageal and gastric adenocarcinoma. <i>Journal of Gastrointestinal Oncology</i> , 2016, 7, 848-859.	1.4	64
54	CD99 is a novel prognostic stromal marker in non-small cell lung cancer. <i>International Journal of Cancer</i> , 2012, 131, 2264-2273.	5.1	63

#	ARTICLE	IF	CITATIONS
55	Comprehensive DNA methylation study identifies novel progression-related and prognostic markers for cutaneous melanoma. <i>BMC Medicine</i> , 2017, 15, 101.	5.5	62
56	CDK-mediated activation of the SCF ^{FBXO28} ubiquitin ligase promotes MYC-driven transcription and tumourigenesis and predicts poor survival in breast cancer. <i>EMBO Molecular Medicine</i> , 2013, 5, 1067-1086.	6.9	61
57	Mutational and gene fusion analyses of primary large cell and large cell neuroendocrine lung cancer. <i>Oncotarget</i> , 2015, 6, 22028-22037.	1.8	61
58	Expression of programmed cell death protein 1 (PD-1) and its ligand PD-L1 in colorectal cancer: Relationship with sidedness and prognosis. <i>Oncolmmunology</i> , 2018, 7, e1465165.	4.6	59
59	Down-Regulation of the Oncogene Cyclin D1 Increases Migratory Capacity in Breast Cancer and Is Linked to Unfavorable Prognostic Features. <i>American Journal of Pathology</i> , 2010, 177, 2886-2897.	3.8	58
60	LGR5 in breast cancer and ductal carcinoma in situ: a diagnostic and prognostic biomarker and a therapeutic target. <i>BMC Cancer</i> , 2020, 20, 542.	2.6	58
61	Expression of the chemokine CXCL14 in the tumour stroma is an independent marker of survival in breast cancer. <i>British Journal of Cancer</i> , 2016, 114, 1117-1124.	6.4	57
62	Coffee and tea consumption and the risk of ovarian cancer: a prospective cohort study and updated meta-analysis. <i>American Journal of Clinical Nutrition</i> , 2012, 95, 1172-1181.	4.7	56
63	Immunohistochemistry in the Differential Diagnostics of Primary Lung Cancer. <i>American Journal of Clinical Pathology</i> , 2013, 140, 37-46.	0.7	56
64	HMG-CoA reductase expression in breast cancer is associated with a less aggressive phenotype and influenced by anthropometric factors. <i>International Journal of Cancer</i> , 2008, 123, 1146-1153.	5.1	55
65	RBM3-Regulated Genes Promote DNA Integrity and Affect Clinical Outcome in Epithelial Ovarian Cancer. <i>Translational Oncology</i> , 2011, 4, 212-IN1.	3.7	54
66	BRCA1-like signature in triple negative breast cancer: Molecular and clinical characterization reveals subgroups with therapeutic potential. <i>Molecular Oncology</i> , 2015, 9, 1528-1538.	4.6	54
67	Prognostic Impact of Tumor Cell Programmed Death Ligand 1 Expression and Immune Cell Infiltration in NSCLC. <i>Journal of Thoracic Oncology</i> , 2019, 14, 628-640.	1.1	54
68	Statin-induced anti-proliferative effects via cyclin D1 and p27 in a window-of-opportunity breast cancer trial. <i>Journal of Translational Medicine</i> , 2015, 13, 133.	4.4	53
69	The integrative clinical impact of tumor-infiltrating T lymphocytes and NK cells in relation to B lymphocyte and plasma cell density in esophageal and gastric adenocarcinoma. <i>Oncotarget</i> , 2017, 8, 72108-72126.	1.8	53
70	Breast tumours following combined hormone replacement therapy express favourable prognostic factors. <i>International Journal of Cancer</i> , 2007, 120, 2202-2207.	5.1	51
71	Low RBM3 protein expression correlates with tumour progression and poor prognosis in malignant melanoma: An analysis of 215 cases from the Malmö Diet and Cancer Study. <i>Journal of Translational Medicine</i> , 2011, 9, 114.	4.4	51
72	Prognostic and treatment predictive significance of SATB1 and SATB2 expression in pancreatic and periampullary adenocarcinoma. <i>Journal of Translational Medicine</i> , 2014, 12, 289.	4.4	49

#	ARTICLE	IF	CITATIONS
73	Contribution of Antibody-based Protein Profiling to the Human Chromosome-centric Proteome Project (C-HPP). <i>Journal of Proteome Research</i> , 2013, 12, 2439-2448.	3.7	48
74	Stage at diagnosis and mortality in women with pregnancy-associated breast cancer (PABC). <i>Breast Cancer Research and Treatment</i> , 2013, 139, 183-192.	2.5	47
75	Cysteinyl leukotriene receptor expression pattern affects migration of breast cancer cells and survival of breast cancer patients. <i>International Journal of Cancer</i> , 2011, 129, 9-22.	5.1	46
76	Genome-Wide DNA Methylation Analysis in Melanoma Reveals the Importance of CpG Methylation in MITF Regulation. <i>Journal of Investigative Dermatology</i> , 2015, 135, 1820-1828.	0.7	46
77	Subtypes of fruit and vegetables, variety in consumption and risk of colon and rectal cancer in the European Prospective Investigation into Cancer and Nutrition. <i>International Journal of Cancer</i> , 2015, 137, 2705-2714.	5.1	45
78	Clinical framework for next generation sequencing based analysis of treatment predictive mutations and multiplexed gene fusion detection in non-small cell lung cancer. <i>Oncotarget</i> , 2017, 8, 34796-34810.	1.8	45
79	The prognostic role of HER2 expression in ductal breast carcinoma in situ (DCIS); a population-based cohort study. <i>BMC Cancer</i> , 2015, 15, 468.	2.6	44
80	Prognostic and predictive significance of podocalyxin-like protein expression in pancreatic and periampullary adenocarcinoma. <i>BMC Clinical Pathology</i> , 2015, 15, 10.	1.8	43
81	Pre-diagnostic anthropometry and survival after colorectal cancer diagnosis in Western European populations. <i>International Journal of Cancer</i> , 2014, 135, 1949-1960.	5.1	42
82	WNT5A-mediated β -catenin-independent signalling is a novel regulator of cancer cell metabolism. <i>Carcinogenesis</i> , 2014, 35, 784-794.	2.8	42
83	A non-functional retinoblastoma tumor suppressor (RB) pathway in premenopausal breast cancer is associated with resistance to tamoxifen. <i>Cell Cycle</i> , 2011, 10, 956-962.	2.6	41
84	Low PIP4K2B Expression in Human Breast Tumors Correlates with Reduced Patient Survival: A Role for PIP4K2B in the Regulation of E-Cadherin Expression. <i>Cancer Research</i> , 2013, 73, 6913-6925.	0.9	41
85	Expression of functional toll like receptor 4 in estrogen receptor/progesterone receptor-negative breast cancer. <i>Breast Cancer Research</i> , 2015, 17, 130.	5.0	41
86	Cigarette smoking and risk of histological subtypes of epithelial ovarian cancer in the EPIC cohort study. <i>International Journal of Cancer</i> , 2012, 130, 2204-2210.	5.1	40
87	Coffee, tea and melanoma risk: findings from the European Prospective Investigation into Cancer and Nutrition. <i>International Journal of Cancer</i> , 2017, 140, 2246-2255.	5.1	39
88	Immunohistochemical profiles in primary lung cancers and epithelial pulmonary metastases. <i>Human Pathology</i> , 2019, 84, 221-230.	2.0	39
89	Quantitative, qualitative and spatial analysis of lymphocyte infiltration in periampullary and pancreatic adenocarcinoma. <i>International Journal of Cancer</i> , 2020, 146, 3461-3473.	5.1	39
90	Diagnostic Value of Insulinoma-Associated Protein 1 (INSM1) and Comparison With Established Neuroendocrine Markers in Pulmonary Cancers. <i>Archives of Pathology and Laboratory Medicine</i> , 2020, 144, 1075-1085.	2.5	38

#	ARTICLE	IF	CITATIONS
91	Increased androgen receptor expression in serous carcinoma of the ovary is associated with an improved survival. <i>Journal of Ovarian Research</i> , 2010, 3, 14.	3.0	37
92	Anthropometric factors in relation to different tumor biological subgroups of postmenopausal breast cancer. <i>International Journal of Cancer</i> , 2009, 124, 402-411.	5.1	36
93	Expression of PD-L1 and PD-1 in Chemoradiotherapy-Naïve Esophageal and Gastric Adenocarcinoma: Relationship With Mismatch Repair Status and Survival. <i>Frontiers in Oncology</i> , 2019, 9, 136.	2.8	36
94	Plasma Folate Concentrations Are Positively Associated with Risk of Estrogen Receptor $\hat{2}$ Negative Breast Cancer in a Swedish Nested Case Control Study. <i>Journal of Nutrition</i> , 2010, 140, 1661-1668.	2.9	35
95	The $\hat{1}$ Np63 Proteins Are Key Allies of BRCA1 in the Prevention of Basal-Like Breast Cancer. <i>Cancer Research</i> , 2011, 71, 1933-1944.	0.9	35
96	Coffee and tea consumption, genotype-based <i>CYP1A2</i> and <i>NAT2</i> activity and colorectal cancer risk-Results from the EPIC cohort study. <i>International Journal of Cancer</i> , 2014, 135, 401-412.	5.1	35
97	S100A9 expressed in ER $\hat{+}$ PgR $\hat{+}$ breast cancers induces inflammatory cytokines and is associated with an impaired overall survival. <i>British Journal of Cancer</i> , 2015, 113, 1234-1243.	6.4	35
98	Prediagnostic Intake of Dairy Products and Dietary Calcium and Colorectal Cancer Survival—Results from the EPIC Cohort Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2014, 23, 1813-1823.	2.5	34
99	Clinical impact of T cells, B cells and the PD-1/PD-L1 pathway in muscle invasive bladder cancer: a comparative study of transurethral resection and cystectomy specimens. <i>Oncolmmunology</i> , 2019, 8, e1644108.	4.6	34
100	Complement inhibitor CSMD1 acts as tumor suppressor in human breast cancer. <i>Oncotarget</i> , 2016, 7, 76920-76933.	1.8	34
101	G Protein–Coupled Estrogen Receptor Is Apoptotic and Correlates with Increased Distant Disease-Free Survival of Estrogen Receptor–Positive Breast Cancer Patients. <i>Clinical Cancer Research</i> , 2013, 19, 1681-1692.	7.0	33
102	Weight change later in life and colon and rectal cancer risk in participants in the EPIC-PANACEA study. <i>American Journal of Clinical Nutrition</i> , 2014, 99, 139-147.	4.7	33
103	Inconsistent results in the analysis of ALK rearrangements in non-small cell lung cancer. <i>BMC Cancer</i> , 2016, 16, 603.	2.6	33
104	The clinical importance of tumour-infiltrating macrophages and dendritic cells in periampullary adenocarcinoma differs by morphological subtype. <i>Journal of Translational Medicine</i> , 2017, 15, 152.	4.4	33
105	The emerging role of FK506-binding proteins as cancer biomarkers: a focus on FKBPL. <i>Biochemical Society Transactions</i> , 2011, 39, 663-668.	3.4	32
106	Associations of Anthropometric Factors with KRAS and BRAF Mutation Status of Primary Colorectal Cancer in Men and Women: A Cohort Study. <i>PLoS ONE</i> , 2014, 9, e98964.	2.5	32
107	Expression and Prognostic Significance of Human Epidermal Growth Factor Receptors 1, 2 and 3 in Periampullary Adenocarcinoma. <i>PLoS ONE</i> , 2016, 11, e0153533.	2.5	32
108	The Prognostic Impact of NK/NKT Cell Density in Periampullary Adenocarcinoma Differs by Morphological Type and Adjuvant Treatment. <i>PLoS ONE</i> , 2016, 11, e0156497.	2.5	32

#	ARTICLE	IF	CITATIONS
109	15-Prostaglandin Dehydrogenase Expression Alone or in Combination with ACSM1 Defines a Subgroup of the Apocrine Molecular Subtype of Breast Carcinoma. <i>Molecular and Cellular Proteomics</i> , 2008, 7, 1795-1809.	3.8	31
110	Evaluation of the prognostic significance of MSMB and CRISP3 in prostate cancer using automated image analysis. <i>Modern Pathology</i> , 2011, 24, 708-719.	5.5	31
111	Sushi domain-containing protein 4 (SUSD4) inhibits complement by disrupting the formation of the classical C3 convertase. <i>FASEB Journal</i> , 2013, 27, 2355-2366.	0.5	31
112	High expression of RNA-binding motif protein 3 in esophageal and gastric adenocarcinoma correlates with intestinal metaplasia-associated tumours and independently predicts a reduced risk of recurrence and death. <i>Biomarker Research</i> , 2014, 2, 11.	6.8	31
113	Pre-diagnostic meat and fibre intakes in relation to colorectal cancer survival in the European Prospective Investigation into Cancer and Nutrition. <i>British Journal of Nutrition</i> , 2016, 116, 316-325.	2.3	30
114	Hormonal factors and pancreatic cancer risk in women: The Malmö Diet and Cancer Study. <i>International Journal of Cancer</i> , 2018, 143, 52-62.	5.1	30
115	Immunohistochemical detection of tyrosine phosphatase SHP1 predicts outcome after radical prostatectomy for localized prostate cancer. <i>International Journal of Cancer</i> , 2010, 126, 2296-2307.	5.1	28
116	Use of a standardized diagnostic approach improves the prognostic information of histopathologic factors in pancreatic and periampullary adenocarcinoma. <i>Diagnostic Pathology</i> , 2014, 9, 80.	2.0	28
117	Prognostic impact of COX-2 in non-small cell lung cancer: A comprehensive compartment-specific evaluation of tumor and stromal cell expression. <i>Cancer Letters</i> , 2015, 356, 837-845.	7.2	28
118	p27 ^{Kip1} is a predictive factor for tamoxifen treatment response but not a prognostic marker in premenopausal breast cancer patients. <i>International Journal of Cancer</i> , 2010, 127, 2851-2858.	5.1	27
119	LRIG1 is a prognostic biomarker in non-small cell lung cancer. <i>Acta Oncologica</i> , 2015, 54, 1113-1119.	1.8	27
120	Comparison of Three Different TTF-1 Clones in Resected Primary Lung Cancer and Epithelial Pulmonary Metastases. <i>American Journal of Clinical Pathology</i> , 2018, 150, 533-544.	0.7	27
121	High RBM3 expression is associated with an improved survival and oxaliplatin response in patients with metastatic colorectal cancer. <i>PLoS ONE</i> , 2017, 12, e0182512.	2.5	27
122	SATB1 is an independent prognostic factor in radically resected upper gastrointestinal tract adenocarcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2014, 465, 649-659.	2.8	26
123	Tumor-specific expression of HMG-CoA reductase in a population-based cohort of breast cancer patients. <i>BMC Clinical Pathology</i> , 2015, 15, 8.	1.8	26
124	Discovery of KIRREL as a biomarker for prognostic stratification of patients with thin melanoma. <i>Biomarker Research</i> , 2019, 7, 1.	6.8	26
125	Cyclin E Overexpression Obstructs Infiltrative Behavior in Breast Cancer: A Novel Role Reflected in the Growth Pattern of Medullary Breast Cancers. <i>Cancer Research</i> , 2005, 65, 9727-9734.	0.9	25
126	Receptor Tyrosine Kinase Signaling Favors a Protumorigenic State in Breast Cancer Cells by Inhibiting the Adaptive Immune Response. <i>Cancer Research</i> , 2010, 70, 7776-7787.	0.9	25

#	ARTICLE	IF	CITATIONS
127	Tumors with Nonfunctional Retinoblastoma Protein Are Killed by Reduced β -Tubulin Levels. <i>Journal of Biological Chemistry</i> , 2012, 287, 17241-17247.	3.4	25
128	Local expression of complement factor I in breast cancer cells correlates with poor survival and recurrence. <i>Cancer Immunology, Immunotherapy</i> , 2015, 64, 467-478.	4.2	25
129	Expression of podocalyxin-like protein is an independent prognostic biomarker in resected esophageal and gastric adenocarcinoma. <i>BMC Clinical Pathology</i> , 2016, 16, 13.	1.8	25
130	Expression of IFITM1 as a prognostic biomarker in resected gastric and esophageal adenocarcinoma. <i>Biomarker Research</i> , 2016, 4, 10.	6.8	25
131	The prognostic impact of the tumour stroma fraction: A machine learning-based analysis in 16 human solid tumour types. <i>EBioMedicine</i> , 2021, 65, 103269.	6.1	25
132	Expression and Prognostic Significance of Human Epidermal Growth Factor Receptors 1 and 3 in Gastric and Esophageal Adenocarcinoma. <i>PLoS ONE</i> , 2016, 11, e0148101.	2.5	25
133	Association of the oestrogen receptor beta with hormone status and prognosis in a cohort of female patients with colorectal cancer. <i>European Journal of Cancer</i> , 2017, 83, 279-289.	2.8	24
134	Ovarian cancer early detection by circulating CA125 in the context of anti-CA125 autoantibody levels: Results from the EPIC cohort. <i>International Journal of Cancer</i> , 2018, 142, 1355-1360.	5.1	24
135	Cyclin A Is a Proliferative Marker with Good Prognostic Value in Node-Negative Breast Cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009, 18, 2501-2506.	2.5	23
136	Stromal Expression of β -Arrestin-1 Predicts Clinical Outcome and Tamoxifen Response in Breast Cancer. <i>Journal of Molecular Diagnostics</i> , 2011, 13, 340-351.	2.8	23
137	Reduced Expression of the Polymeric Immunoglobulin Receptor in Pancreatic and Periampullary Adenocarcinoma Signifies Tumour Progression and Poor Prognosis. <i>PLoS ONE</i> , 2014, 9, e112728.	2.5	23
138	HMG-CoA reductase expression in primary colorectal cancer correlates with favourable clinicopathological characteristics and an improved clinical outcome. <i>Diagnostic Pathology</i> , 2014, 9, 78.	2.0	23
139	Fibre intake and incident colorectal cancer depending on fibre source, sex, tumour location and Tumour, Node, Metastasis stage. <i>British Journal of Nutrition</i> , 2015, 114, 959-969.	2.3	23
140	Analysis of the Human Prostate-Specific Proteome Defined by Transcriptomics and Antibody-Based Profiling Identifies TMEM79 and ACOXL as Two Putative, Diagnostic Markers in Prostate Cancer. <i>PLoS ONE</i> , 2015, 10, e0133449.	2.5	23
141	High Estrogen Receptor β Expression Is Prognostic among Adjuvant Chemotherapy-Treated Patients: Results from a Population-Based Breast Cancer Cohort. <i>Clinical Cancer Research</i> , 2017, 23, 766-777.	7.0	23
142	Erythropoietin Receptor Expression and Correlation to Tamoxifen Response and Prognosis in Breast Cancer. <i>Clinical Cancer Research</i> , 2009, 15, 5552-5559.	7.0	22
143	Infiltration of β T cells, IL-17+ T cells and FoxP3+ T cells in human breast cancer. <i>Cancer Biomarkers</i> , 2018, 20, 395-409.	1.7	22
144	Expression and prognostic significance of the polymeric immunoglobulin receptor in epithelial ovarian cancer. <i>Journal of Ovarian Research</i> , 2014, 7, 26.	3.0	21

#	ARTICLE	IF	CITATIONS
145	Pancreatic cancer risk in relation to sex, lifestyle factors, and pre-diagnostic anthropometry in the Malmö Diet and Cancer Study. <i>Biology of Sex Differences</i> , 2016, 7, 66.	4.1	21
146	Low RBM3 Protein Expression Correlates with Clinical Stage, Prognostic Classification and Increased Risk of Treatment Failure in Testicular Non-Seminomatous Germ Cell Cancer. <i>PLoS ONE</i> , 2015, 10, e0121300.	2.5	21
147	Investigation of molecular alterations of <i>AKT</i> in triple-negative breast cancer. <i>Histopathology</i> , 2014, 64, 660-670.	2.9	20
148	Reduced expression of ezrin in urothelial bladder cancer signifies more advanced tumours and an impaired survival: validatory study of two independent patient cohorts. <i>BMC Urology</i> , 2014, 14, 36.	1.4	20
149	Tumor-Associated CD68+, CD163+, and MARCO+ Macrophages as Prognostic Biomarkers in Patients With Treatment-Naïve Gastroesophageal Adenocarcinoma. <i>Frontiers in Oncology</i> , 2020, 10, 534761.	2.8	20
150	Complement inhibitor factor H expressed by breast cancer cells differentiates CD14 ⁺ human monocytes into immunosuppressive macrophages. <i>Oncolmmunology</i> , 2020, 9, 1731135.	4.6	20
151	Multiple Miscarriages Are Associated with the Risk of Ovarian Cancer: Results from the European Prospective Investigation into Cancer and Nutrition. <i>PLoS ONE</i> , 2012, 7, e37141.	2.5	19
152	Intra-tumour IgA1 is common in cancer and is correlated with poor prognosis in bladder cancer.. <i>Heliyon</i> , 2016, 2, e00143.	3.2	19
153	The prognostic impact of COX-2 expression in breast cancer depends on oral contraceptive history, preoperative NSAID use, and tumor size. <i>International Journal of Cancer</i> , 2017, 140, 163-175.	5.1	19
154	Tumor-associated autoantibodies as early detection markers for ovarian cancer? A prospective evaluation. <i>International Journal of Cancer</i> , 2018, 143, 515-526.	5.1	18
155	Relationship between mismatch repair immunophenotype and long-term survival in patients with resected periampullary adenocarcinoma. <i>Journal of Translational Medicine</i> , 2018, 16, 66.	4.4	18
156	BET Inhibition as a Rational Therapeutic Strategy for Invasive Lobular Breast Cancer. <i>Clinical Cancer Research</i> , 2019, 25, 7139-7150.	7.0	18
157	A Functional Genomic Screen Identifies the Deubiquitinase USP11 as a Novel Transcriptional Regulator of ER α in Breast Cancer. <i>Cancer Research</i> , 2020, 80, 5076-5088.	0.9	18
158	Topographical Distribution and Spatial Interactions of Innate and Semi-Innate Immune Cells in Pancreatic and Other Periampullary Adenocarcinoma. <i>Frontiers in Immunology</i> , 2020, 11, 558169.	4.8	18
159	Histological specificity of alterations and expression of <i>KIT</i> and <i>KITLG</i> in non-small cell lung carcinoma. <i>Genes Chromosomes and Cancer</i> , 2013, 52, 1088-1096.	2.8	17
160	Discordant human epidermal growth factor receptor 2 overexpression in primary and metastatic upper gastrointestinal adenocarcinoma signifies poor prognosis. <i>Histopathology</i> , 2016, 68, 230-240.	2.9	17
161	Reaching the limits of prognostication in non-small cell lung cancer: an optimized biomarker panel fails to outperform clinical parameters. <i>Modern Pathology</i> , 2017, 30, 964-977.	5.5	17
162	Genetic variation in the ADIPOQ gene, adiponectin concentrations and risk of colorectal cancer: a Mendelian Randomization analysis using data from three large cohort studies. <i>European Journal of Epidemiology</i> , 2017, 32, 419-430.	5.7	17

#	ARTICLE	IF	CITATIONS
163	Hypoxia Attenuates Trastuzumab Uptake and Trastuzumab-Emtansine (T-DM1) Cytotoxicity through Redistribution of Phosphorylated Caveolin-1. <i>Molecular Cancer Research</i> , 2020, 18, 644-656.	3.4	17
164	Protein Expression Profiling of Formalin-Fixed Paraffin-Embedded Tissue Using Recombinant Antibody Microarrays. <i>Journal of Proteome Research</i> , 2013, 12, 5943-5953.	3.7	16
165	Podocalyxin-like and RNA-binding motif protein 3 are prognostic biomarkers in urothelial bladder cancer: a validity study. <i>Biomarker Research</i> , 2017, 5, 10.	6.8	16
166	Prognostic effect of hENT1, dCK and HuR expression by morphological type in periampullary adenocarcinoma, including pancreatic cancer. <i>Acta Oncologica</i> , 2016, 55, 286-296.	1.8	14
167	CYP27A1 expression is associated with risk of late lethal estrogen receptor-positive breast cancer in postmenopausal patients. <i>Breast Cancer Research</i> , 2020, 22, 123.	5.0	14
168	T cells, B cells, and PD-L1 expression in esophageal and gastric adenocarcinoma before and after neoadjuvant chemotherapy: relationship with histopathological response and survival. <i>Oncology</i> , 2021, 10, 1921443.	4.6	14
169	Human G-MDSCs are neutrophils at distinct maturation stages promoting tumor growth in breast cancer. <i>Life Science Alliance</i> , 2020, 3, e202000893.	2.8	14
170	Low expression of SHP-2 is associated with less favorable prostate cancer outcomes. <i>Tumor Biology</i> , 2013, 34, 637-642.	1.8	13
171	A systematic analysis of commonly used antibodies in cancer diagnostics. <i>Histopathology</i> , 2014, 64, 293-305.	2.9	13
172	Combined and individual tumor-specific expression of insulin-like growth factor-I receptor, insulin receptor and phospho-insulin-like growth factor-I receptor/insulin receptor in primary breast cancer: Implications for prognosis in different treatment groups. <i>Oncotarget</i> , 2017, 8, 9093-9107.	1.8	13
173	Translational study reveals a two-faced role of RBM3 in pancreatic cancer and suggests its potential value as a biomarker for improved patient stratification. <i>Oncotarget</i> , 2018, 9, 6188-6200.	1.8	13
174	FKBPL: a marker of good prognosis in breast cancer. <i>Oncotarget</i> , 2015, 6, 12209-12223.	1.8	13
175	CRIM1 is localized to the podocyte filtration slit diaphragm of the adult human kidney. <i>Nephrology Dialysis Transplantation</i> , 2009, 24, 2038-2044.	0.7	12
176	The Target for Statins, HMG-CoA Reductase, Is Expressed in Ductal Carcinoma-In Situ and May Predict Patient Response to Radiotherapy. <i>Annals of Surgical Oncology</i> , 2014, 21, 2911-2919.	1.5	12
177	Delineating transcriptional networks of prognostic gene signatures refines treatment recommendations for lymph node-negative breast cancer patients. <i>FEBS Journal</i> , 2015, 282, 3455-3473.	4.7	12
178	Early diagnostic value of Bcl-3 localization in colorectal cancer. <i>BMC Cancer</i> , 2015, 15, 341.	2.6	12
179	Significant association and synergistic adverse prognostic effect of podocalyxin-like protein and epidermal growth factor receptor expression in colorectal cancer. <i>Journal of Translational Medicine</i> , 2016, 14, 128.	4.4	12
180	Re-evaluation of HER2 status in 606 breast cancers using gene protein assay on tissue microarrays versus routine pathological assessment. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2020, 477, 317-320.	2.8	12

#	ARTICLE	IF	CITATIONS
181	Cyclin I is expressed in human breast cancer and closely associated with VEGF and KDR expression. <i>Breast Cancer Research and Treatment</i> , 2005, 89, 313-316.	2.5	11
182	Anthropometric factors and ovarian cancer risk in the Malmö Diet and Cancer Study. <i>Cancer Epidemiology</i> , 2011, 35, 432-437.	1.9	11
183	Mucin staining is of limited value in addition to basic immunohistochemical analyses in the diagnostics of non-small cell lung cancer. <i>Scientific Reports</i> , 2019, 9, 1319.	3.3	11
184	Molecular Evolution of Specific Human Antibody against MUC1 Mucin Results in Improved Recognition of the Antigen on Tumor Cells. <i>Tumor Biology</i> , 2009, 30, 221-231.	1.8	10
185	Incident urothelial cancer in the Malmö Diet and Cancer Study: cohort characteristics and further validation of ezrin as a prognostic biomarker. <i>Diagnostic Pathology</i> , 2014, 9, 189.	2.0	10
186	The human complement inhibitor Sushi Domain-Containing Protein 4 (SUSD4) expression in tumor cells and infiltrating T cells is associated with better prognosis of breast cancer patients. <i>BMC Cancer</i> , 2015, 15, 737.	2.6	10
187	Comprehensive analysis of RNA binding motif protein 3 (RBM3) in non-small cell lung cancer. <i>Cancer Medicine</i> , 2020, 9, 5609-5619.	2.8	10
188	Expression and prognostic significance of human epidermal growth factor receptors 1, 2 and 3 in oesophageal and gastric adenocarcinomas preneoadjuvant and postneoadjuvant treatment. <i>Journal of Clinical Pathology</i> , 2018, 71, 451-462.	2.0	9
189	Chemotherapy, host response and molecular dynamics in periampullary cancer: the CHAMP study. <i>BMC Cancer</i> , 2020, 20, 308.	2.6	9
190	Primary Melanoma Tumors from CDKN2A Mutation Carriers Do Not Belong to a Distinct Molecular Subclass. <i>Journal of Investigative Dermatology</i> , 2014, 134, 3000-3003.	0.7	8
191	Multi-parametric profiling of renal cell, colorectal, and ovarian cancer identifies tumour-type-specific stroma phenotypes and a novel vascular biomarker. <i>Journal of Pathology: Clinical Research</i> , 2017, 3, 214-224.	3.0	8
192	Interactions Between ABCB1 Genotype and Preoperative Statin Use Impact Clinical Outcomes Among Breast Cancer Patients. <i>Frontiers in Oncology</i> , 2018, 8, 428.	2.8	8
193	Prognostic impact of tumor-specific insulin-like growth factor binding protein 7 (IGFBP7) levels in breast cancer: a prospective cohort study. <i>Carcinogenesis</i> , 2021, 42, 1314-1325.	2.8	8
194	The β -tubulin meshwork assists in the recruitment of PCNA to chromatin in mammalian cells. <i>Communications Biology</i> , 2021, 4, 767.	4.4	8
195	High Infiltration of CD68 ⁺ /CD163 ⁺ Macrophages Is an Adverse Prognostic Factor after Neoadjuvant Chemotherapy in Esophageal and Gastric Adenocarcinoma. <i>Journal of Innate Immunity</i> , 2022, 14, 615-628.	3.8	8
196	Serum concentrations of human chorionic gonadotropin beta and its association with survival in patients with colorectal cancer. <i>Cancer Biomarkers</i> , 2012, 11, 173-181.	1.7	7
197	Detecting EGFR alterations in clinical specimens – pitfalls and necessities. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2013, 463, 755-764.	2.8	7
198	Podocalyxin-like protein as a predictive biomarker for benefit of neoadjuvant chemotherapy in resectable gastric and esophageal adenocarcinoma. <i>Journal of Translational Medicine</i> , 2018, 16, 290.	4.4	7

#	ARTICLE	IF	CITATIONS
199	Coffee Is Associated With Lower Breast Tumor Insulin-Like Growth Factor Receptor 1 Levels in Normal-Weight Patients and Improved Prognosis Following Tamoxifen or Radiotherapy Treatment. <i>Frontiers in Endocrinology</i> , 2018, 9, 306.	3.5	7
200	Inflammatory macrophage derived TNF α downregulates estrogen receptor α via FOXO3a inactivation in human breast cancer cells. <i>Experimental Cell Research</i> , 2020, 390, 111932.	2.6	7
201	Human Epidermal Growth Factor Receptor-3 Expression Is Regulated at Transcriptional Level in Breast Cancer Settings by Junctional Adhesion Molecule-A via a Pathway Involving Beta-Catenin and FOXA1. <i>Cancers</i> , 2021, 13, 871.	3.7	7
202	Reproductive Factors, Exogenous Hormone Use, and Risk of B-Cell Non-Hodgkin Lymphoma in a Cohort of Women From the European Prospective Investigation Into Cancer and Nutrition. <i>American Journal of Epidemiology</i> , 2019, 188, 274-281.	3.4	6
203	Ovarian cancer subtypes and survival in relation to three comprehensive imaging parameters. <i>Journal of Ovarian Research</i> , 2020, 13, 26.	3.0	6
204	Pre- and Postoperative Circulating IGF-I, IGFBP-3, and IGFBP-7 Levels in Relation to Endocrine Treatment and Breast Cancer Recurrence: A Nested Case-Control Study. <i>Frontiers in Oncology</i> , 2021, 11, 626058.	2.8	6
205	Body Mass Index Influences the Prognostic Impact of Combined Nuclear Insulin Receptor and Estrogen Receptor Expression in Primary Breast Cancer. <i>Frontiers in Endocrinology</i> , 2017, 8, 332.	3.5	5
206	Pre-diagnostic anthropometry, sex, and risk of colorectal cancer according to tumor immune cell composition. <i>Oncolmmunology</i> , 2019, 8, e1664275.	4.6	5
207	Metabolic factors and the risk of small intestine cancers: Pooled study of 800,000 individuals in the metabolic syndrome and cancer project. <i>International Journal of Cancer</i> , 2021, 149, 66-74.	5.1	5
208	PRR11 unveiled as a top candidate biomarker within the RBM3-regulated transcriptome in pancreatic cancer. <i>Journal of Pathology: Clinical Research</i> , 2022, 8, 65-77.	3.0	5
209	MMP7 Modulation by Short- and Long-term Radiotherapy in Patients with Rectal Cancer. <i>In Vivo</i> , 2018, 32, 133-138.	1.3	5
210	OUP accepted manuscript. <i>Clinical Chemistry</i> , 2022, , .	3.2	5
211	Lymphocyte antigen 6 superfamily member D is a marker of urothelial and squamous differentiation: implications for risk stratification of bladder cancer. <i>Biomarker Research</i> , 2020, 8, 51.	6.8	4
212	Additive clinical impact of epidermal growth factor receptor and podocalyxin-like protein expression in pancreatic and periampullary adenocarcinomas. <i>Scientific Reports</i> , 2020, 10, 10373.	3.3	4
213	The Prognostic Impact of Intratumoral Aryl Hydrocarbon Receptor in Primary Breast Cancer Depends on the Type of Endocrine Therapy: A Population-Based Cohort Study. <i>Frontiers in Oncology</i> , 2021, 11, 642768.	2.8	4
214	Delineating the intra-patient heterogeneity of molecular alterations in treatment-naïve colorectal cancer with peritoneal carcinomatosis. <i>Modern Pathology</i> , 2022, , .	5.5	4
215	Clinical significance of RBM3 expression in surgically treated colorectal lung metastases and paired primary tumors. <i>Journal of Surgical Oncology</i> , 2021, 123, 1144-1156.	1.7	3
216	Pre-clinical and clinical studies on the role of RBM3 in muscle-invasive bladder cancer: longitudinal expression, transcriptome-level effects and modulation of chemosensitivity. <i>BMC Cancer</i> , 2022, 22, 131.	2.6	3

#	ARTICLE	IF	CITATIONS
217	Interplay between Caveolin-1 and body and tumor size affects clinical outcomes in breast cancer. <i>Translational Oncology</i> , 2022, 22, 101464.	3.7	3
218	Translational Database Selection and Multiplexed Sequence Capture for Up Front Filtering of Reliable Breast Cancer Biomarker Candidates. <i>PLoS ONE</i> , 2011, 6, e20794.	2.5	2
219	Nuclear T-STAR Protein Expression Correlates with HER2 Status, Hormone Receptor Negativity and Prolonged Recurrence Free Survival in Primary Breast Cancer and Decreased Cancer Cell Growth In Vitro. <i>PLoS ONE</i> , 2013, 8, e70596.	2.5	2
220	Patient Characteristics Influence Activated Signal Transducer and Activator of Transcription 3 (STAT3) Levels in Primary Breast Cancer—Impact on Prognosis. <i>Frontiers in Oncology</i> , 2020, 10, 1278.	2.8	2
221	Clinical significance of stromal ER and PR expression in periampullary adenocarcinoma. <i>Biomarker Research</i> , 2019, 7, 26.	6.8	1
222	Development of a Novel Weighted Ranking Method for Immunohistochemical Quantification of a Heterogeneously Expressed Protein in Gastro-Esophageal Cancers. <i>Cancers</i> , 2021, 13, 1286.	3.7	1