

# Christer Borgfeldt

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8266901/publications.pdf>

Version: 2024-02-01

71  
papers

1,754  
citations

304743

22  
h-index

302126

39  
g-index

72  
all docs

72  
docs citations

72  
times ranked

2357  
citing authors

#	ARTICLE	IF	CITATIONS
1	Cancer risk after hospital discharge diagnosis of benign ovarian cysts and endometriosis. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2004, 83, 395-400.	2.8	115
2	Menopause-related symptoms: What are the background factors? A prospective population-based cohort study of Swedish women (The Women's Health in Lund Area study). <i>American Journal of Obstetrics and Gynecology</i> , 2003, 189, 1646-1653.	1.3	113
3	Robot assisted laparoscopic radical hysterectomy and pelvic lymphadenectomy with short and long term morbidity data. <i>Gynecologic Oncology</i> , 2009, 113, 185-190.	1.4	100
4	A novel diagnostic index combining HE4, CA125 and age may improve triage of women with suspected ovarian cancer – An international multicenter study in women with an ovarian mass. <i>Gynecologic Oncology</i> , 2015, 138, 640-646.	1.4	78
5	The sentinel node concept in early cervical cancer performs well in tumors smaller than 2 cm. <i>Gynecologic Oncology</i> , 2010, 117, 266-269.	1.4	75
6	Radiotherapy Versus Inguinofemoral Lymphadenectomy as Treatment for Vulvar Cancer Patients With Micrometastases in the Sentinel Node: Results of GROINSS-V II. <i>Journal of Clinical Oncology</i> , 2021, 39, 3623-3632.	1.6	69
7	Cleaved Forms of the Urokinase Plasminogen Activator Receptor in Plasma Have Diagnostic Potential and Predict Postoperative Survival in Patients with Ovarian Cancer. <i>Clinical Cancer Research</i> , 2008, 14, 5785-5793.	7.0	67
8	Reproducibility and accuracy of robot-assisted laparoscopic fertility sparing radical trachelectomy. <i>Gynecologic Oncology</i> , 2012, 127, 484-488.	1.4	67
9	Fertility-sparing surgery and outcome in fertile women with ovarian borderline tumors and epithelial invasive ovarian cancer. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2007, 134, 110-114.	1.1	56
10	Lymphovascular space invasion as a predictive factor for lymph node metastases and survival in endometrioid endometrial cancer – a Swedish Gynecologic Cancer Group (SweGCG) study. <i>Acta Oncologica</i> , 2019, 58, 1628-1633.	1.8	55
11	Comparison of use of vaginal HPV self-sampling and offering flexible appointments as strategies to reach long-term non-attending women in organized cervical screening. <i>Journal of Clinical Virology</i> , 2013, 58, 155-160.	3.1	54
12	High preoperative blood levels of HE4 predicts poor prognosis in patients with ovarian cancer. <i>Journal of Ovarian Research</i> , 2012, 5, 20.	3.0	48
13	Dedifferentiation of serous ovarian cancer from cystic to solid tumors is associated with increased expression of mRNA for urokinase plasminogen activator (uPA), its receptor (uPAR) and its inhibitor (PAI-1). <i>International Journal of Cancer</i> , 2001, 92, 497-502.	5.1	45
14	Postmenopausal hormone therapy and risk of stroke: A pooled analysis of data from population-based cohort studies. <i>PLoS Medicine</i> , 2017, 14, e1002445.	8.4	44
15	Use of proton pump inhibitors (PPI) and history of earlier fracture are independent risk factors for fracture in postmenopausal women. <i>The WHILA study. Maturitas</i> , 2014, 78, 310-315.	2.4	43
16	Data quality in the Swedish Quality Register of Gynecologic Cancer – a Swedish Gynecologic Cancer Group (SweGCG) study. <i>Acta Oncologica</i> , 2018, 57, 346-353.	1.8	41
17	Background factors influencing somatic and psychological symptoms in middle-age women with different hormonal status. <i>Maturitas</i> , 2005, 52, 306-318.	2.4	38
18	Self-sampling with HPV mRNA analyses from vagina and urine compared with cervical samples. <i>Journal of Clinical Virology</i> , 2018, 101, 69-73.	3.1	38

#	ARTICLE	IF	CITATIONS
19	Low-dose hormone therapy and carbohydrate metabolism. <i>Fertility and Sterility</i> , 2003, 79, 550-555.	1.0	32
20	Elderly women above screening age diagnosed with cervical cancer have a worse prognosis. <i>Anticancer Research</i> , 2014, 34, 5147-51.	1.1	29
21	A Swedish population-based evaluation of benign hysterectomy, comparing minimally invasive and abdominal surgery. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2018, 222, 113-118.	1.1	25
22	Risk factors for lymph node metastases in women with endometrial cancer: A population-based, nationwide register study—on behalf of the Swedish Gynecological Cancer Group. <i>International Journal of Cancer</i> , 2017, 140, 2693-2700.	5.1	23
23	Population-based study of survival for women with serous cancer of the ovary, fallopian tube, peritoneum or undesignated origin - on behalf of the Swedish gynecological cancer group (SweGCG). <i>Gynecologic Oncology</i> , 2017, 144, 167-173.	1.4	23
24	Vaginal self-sampling without preservative for human papillomavirus testing shows good sensitivity. <i>Journal of Clinical Virology</i> , 2013, 56, 52-56.	3.1	21
25	Robot-assisted surgery in cervical cancer patients reduces the time to normal activities of daily living. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2015, 94, 260-265.	2.8	21
26	Follow up with HPV test and cytology as test of cure, 6 months after conization, is reliable. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2016, 95, 1251-1257.	2.8	21
27	Increased blood cadmium levels were not associated with increased fracture risk but with increased total mortality in women: the Malmö Diet and Cancer Study. <i>Osteoporosis International</i> , 2017, 28, 2401-2408.	3.1	21
28	High tissue content of urokinase plasminogen activator (u-PA) is associated with high stromal expression of u-PA mRNA in poorly differentiated serous ovarian carcinoma. <i>International Journal of Cancer</i> , 1998, 79, 588-595.	5.1	20
29	Changes in lipid and lipoprotein profile in postmenopausal women receiving low-dose combinations of 17 $\beta$ -estradiol and norethisterone acetate. <i>Menopause</i> , 2002, 9, 335-342.	2.0	20
30	Low-dose oral combination of 17 $\beta$ -estradiol and norethisterone acetate in postmenopausal women decreases factor VII, fibrinogen, antithrombin and plasminogen activator inhibitor-1. <i>Climacteric</i> , 2004, 7, 78-85.	2.4	19
31	Vaginal and Urine Self-sampling Compared to Cervical Sampling for HPV-testing with the Cobas 4800 HPV Test. <i>Anticancer Research</i> , 2017, 37, 4183-4187.	1.1	18
32	A population-based registry study evaluating surgery in newly diagnosed uterine cancer. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2016, 95, 901-911.	2.8	17
33	Long-term survival in women with borderline ovarian tumors: a population-based survey of borderline ovarian tumors in Sweden 1960–2007. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2016, 95, 473-479.	2.8	17
34	Lymph node metastases as only qualifier for stage IV serous ovarian cancer confers longer survival than other sites of distant disease—a Swedish Gynecologic Cancer Group (SweGCG) study. <i>Acta Oncologica</i> , 2018, 57, 331-337.	1.8	17
35	High tumor tissue concentration of urokinase plasminogen activator receptor is associated with good prognosis in patients with ovarian cancer. <i>International Journal of Cancer</i> , 2003, 107, 658-665.	5.1	16
36	Preoperative and intraoperative assessment of myometrial invasion in endometrial cancer—A Swedish Gynecologic Cancer Group (SweGCG) study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2021, 100, 1526-1533.	2.8	16

#	ARTICLE	IF	CITATIONS
37	The economic burden of human papillomavirus-related precancers and cancers in Sweden. PLoS ONE, 2017, 12, e0179520.	2.5	16
38	A Biomarker Panel Increases the Diagnostic Performance for Epithelial Ovarian Cancer Type I and II in Young Women. Anticancer Research, 2016, 36, 957-65.	1.1	16
39	Cervical cancer prevention among long-term screening non-attendees by vaginal self-collected samples for hr-HPV mRNA detection. Infectious Agents and Cancer, 2020, 15, 10.	2.6	15
40	A multiplex biomarker assay improves the diagnostic performance of HE4 and CA125 in ovarian tumor patients. PLoS ONE, 2020, 15, e0240418.	2.5	15
41	High preoperative plasma concentration of tissue plasminogen activator (tPA) is an independent marker for shorter overall survival in patients with ovarian cancer. Gynecologic Oncology, 2003, 91, 112-117.	1.4	12
42	HE4 is a novel tissue marker for therapy response and progestin resistance in medium- and low-risk endometrial hyperplasia. British Journal of Cancer, 2016, 115, 725-730.	6.4	12
43	Primary treatment patterns and survival of cervical cancer in Sweden: A population-based Swedish Gynecologic Cancer Group Study. Gynecologic Oncology, 2019, 155, 229-236.	1.4	12
44	Population-based primary HPV mRNA cervical screening compared with cytology screening. Preventive Medicine, 2019, 124, 61-66.	3.4	11
45	Primary treatment and relative survival by stage and age in vulvar squamous cell carcinoma: A population-based SweGCG study. Gynecologic Oncology, 2020, 159, 663-671.	1.4	10
46	Low androstenedione/sex hormone binding globulin ratio increases fracture risk in postmenopausal women. The Women's Health in the Lund Area study. Maturitas, 2013, 75, 270-275.	2.4	9
47	Ovarian Cancer Surgery – A Population-based Registry Study. Anticancer Research, 2017, 37, 1837-1845.	1.1	9
48	Age influences the clinical significance of atypical glandular cells on cytology. Anticancer Research, 2015, 35, 913-9.	1.1	9
49	Prevalence of High-risk HPV in Postmenopausal Women with Benign Cervical Cytology – A Population-based Cohort Study. Anticancer Research, 2018, 38, 4221-4228.	1.1	8
50	Increased HPV detection by the use of a pre-heating step on vaginal self-samples analysed by Aptima HPV assay. Journal of Virological Methods, 2019, 270, 18-20.	2.1	8
51	Detection of HPV mRNA in Self-collected Vaginal Samples Among Women at 69-70 Years of Age. Anticancer Research, 2019, 39, 381-386.	1.1	8
52	Equal prevalence of severe cervical dysplasia by HPV self-sampling and by midwife-collected samples for primary HPV screening: a randomised controlled trial. European Journal of Cancer Prevention, 2021, 30, 334-340.	1.3	8
53	Confounders other than comorbidity explain survival differences in Danish and Swedish ovarian cancer patients – a comparative cohort study. Acta Oncologica, 2018, 57, 1100-1108.	1.8	6
54	Implementation of National Guidelines increased survival in advanced ovarian cancer - A population-based nationwide SweGCG study. Gynecologic Oncology, 2021, 161, 244-250.	1.4	6

#	ARTICLE	IF	CITATIONS
55	Promotion of Cervical Screening among Long-term Non-attendees by Human Papillomavirus Self-sampling. <i>Journal of Cancer Prevention</i> , 2021, 26, 25-31.	2.0	5
56	Cost-effectiveness of HPV testing in cervical screening among 40- to 42-year-old women is unreasonable. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2022, 101, 374-378.	2.8	5
57	14-type HPV mRNA test in triage of HPV DNA-positive postmenopausal women with normal cytology. <i>BMC Cancer</i> , 2020, 20, 1025.	2.6	4
58	HPV73 in cervical cancer and distribution of HPV73 variants in cervical dysplasia. <i>International Journal of Cancer</i> , 2021, 149, 936-943.	5.1	4
59	Circulating Markers of Neutrophil Extracellular Traps (NETs) in Patients With Ovarian Tumors. <i>Anticancer Research</i> , 2022, 42, 965-971.	1.1	4
60	Long-term incidence of endometrial cancer after endometrial resection and ablation: A population based Swedish gynecologic cancer group ( SweGCG ) study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 0, , .	2.8	4
61	Time Trends for Incidence and Net Survival of Cervical Cancer in Sweden 1960-2014: A Nationwide Population-Based Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2022, 31, 1572-1581.	2.5	4
62	Primary screening for increased fracture risk by the FRAX® questionnaire: uptake rates in relation to invitation method. <i>Archives of Osteoporosis</i> , 2019, 14, 51.	2.4	3
63	Areas to Improve Quality of Life After Ovarian Tumor Surgery and Adjuvant Treatment. <i>In Vivo</i> , 2021, 35, 2399-2408.	1.3	2
64	Survival in endometrial cancer in relation to minimally invasive surgery or open surgery - a Swedish Gynecologic Cancer Group (SweGCG) study. <i>BMC Cancer</i> , 2021, 21, 658.	2.6	2
65	Negative Pressure Wound Therapy (NPWT) in Groin Wounds After Lymphadenectomy in Vulvar Cancer Patients. <i>In Vivo</i> , 2020, 34, 3511-3517.	1.3	1
66	Detection of HPV mRNA in Self-collected Vaginal Samples Among Urban Ethiopian Women. <i>Anticancer Research</i> , 2020, 40, 1513-1517.	1.1	1
67	Incidence and survival of epithelial ovarian, fallopian tube, peritoneal, and undesigned abdominal/pelvic cancers in Sweden 1960-2014: A population-based cohort study. <i>BMC Cancer</i> , 2021, 21, 465.	2.6	1
68	The association between common urogenital infections and cervical neoplasia - A nationwide cohort study of over four million women (2002-2018). <i>Lancet Regional Health - Europe</i> , The, 2022, 17, 100378.	5.6	1
69	Robot-assisted removal of a lymphocyst causing severe neuralgic pain and adductor atrophy. <i>Journal of Robotic Surgery</i> , 2011, 5, 299-302.	1.8	0
70	A population-based study of pelvic serous carcinoma in Sweden: Primary site, FIGO stage and survival. <i>Journal of Clinical Oncology</i> , 2015, 33, e16533-e16533.	1.6	0
71	The wait time to primary surgery in endometrial cancer - impact on survival and predictive factors: a population-based SweGCG study. <i>Acta Oncologica</i> , 2022, 61, 30-37.	1.8	0