

# Christer Borgfeldt

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8266901/publications.pdf>

Version: 2024-02-01

71  
papers

1,754  
citations

304743

22  
h-index

302126

39  
g-index

72  
all docs

72  
docs citations

72  
times ranked

2357  
citing authors

#	ARTICLE	IF	CITATIONS
1	The wait time to primary surgery in endometrial cancer – impact on survival and predictive factors: a population-based SweGCG study. <i>Acta Oncologica</i> , 2022, 61, 30-37.	1.8	0
2	Cost-effectiveness of testing in cervical screening among 40-year-old to 42-year-old women is unreasonable. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2022, 101, 374-378.	2.8	5
3	Circulating Markers of Neutrophil Extracellular Traps (NETs) in Patients With Ovarian Tumors. <i>Anticancer Research</i> , 2022, 42, 965-971.	1.1	4
4	The association between common urogenital infections and cervical neoplasia – A nationwide cohort study of over four million women (2002–2018). <i>Lancet Regional Health - Europe</i> , The, 2022, 17, 100378.	5.6	1
5	Time Trends for Incidence and Net Survival of Cervical Cancer in Sweden 1960–2014 – A Nationwide Population-Based Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2022, 31, 1572-1581.	2.5	4
6	Areas to Improve Quality of Life After Ovarian Tumor Surgery and Adjuvant Treatment. <i>In Vivo</i> , 2021, 35, 2399-2408.	1.3	2
7	Promotion of Cervical Screening among Long-term Non-attendees by Human Papillomavirus Self-sampling. <i>Journal of Cancer Prevention</i> , 2021, 26, 25-31.	2.0	5
8	Implementation of National Guidelines increased survival in advanced ovarian cancer - A population-based nationwide SweGCG study. <i>Gynecologic Oncology</i> , 2021, 161, 244-250.	1.4	6
9	Incidence and survival of epithelial ovarian, fallopian tube, peritoneal, and undesignated abdominal/pelvic cancers in Sweden 1960–2014: A population-based cohort study. <i>BMC Cancer</i> , 2021, 21, 465.	2.6	1
10	HPV73 in cervical cancer and distribution of HPV73 variants in cervical dysplasia. <i>International Journal of Cancer</i> , 2021, 149, 936-943.	5.1	4
11	Equal prevalence of severe cervical dysplasia by HPV self-sampling and by midwife-collected samples for primary HPV screening: a randomised controlled trial. <i>European Journal of Cancer Prevention</i> , 2021, 30, 334-340.	1.3	8
12	Preoperative and intraoperative assessment of myometrial invasion in endometrial cancer – A Swedish Gynecologic Cancer Group (SweGCG) study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2021, 100, 1526-1533.	2.8	16
13	Survival in endometrial cancer in relation to minimally invasive surgery or open surgery – a Swedish Gynecologic Cancer Group (SweGCG) study. <i>BMC Cancer</i> , 2021, 21, 658.	2.6	2
14	Radiotherapy Versus Inguinofemoral Lymphadenectomy as Treatment for Vulvar Cancer Patients With Micrometastases in the Sentinel Node: Results of GROINSS-V II. <i>Journal of Clinical Oncology</i> , 2021, 39, 3623-3632.	1.6	69
15	Negative Pressure Wound Therapy (NPWT) in Groin Wounds After Lymphadenectomy in Vulvar Cancer Patients. <i>In Vivo</i> , 2020, 34, 3511-3517.	1.3	1
16	14-type HPV mRNA test in triage of HPV DNA-positive postmenopausal women with normal cytology. <i>BMC Cancer</i> , 2020, 20, 1025.	2.6	4
17	Primary treatment and relative survival by stage and age in vulvar squamous cell carcinoma: A population-based SweGCG study. <i>Gynecologic Oncology</i> , 2020, 159, 663-671.	1.4	10
18	Cervical cancer prevention among long-term screening non-attendees by vaginal self-collected samples for hr-HPV mRNA detection. <i>Infectious Agents and Cancer</i> , 2020, 15, 10.	2.6	15

#	ARTICLE	IF	CITATIONS
19	Detection of HPV mRNA in Self-collected Vaginal Samples Among Urban Ethiopian Women. <i>Anticancer Research</i> , 2020, 40, 1513-1517.	1.1	1
20	A multiplex biomarker assay improves the diagnostic performance of HE4 and CA125 in ovarian tumor patients. <i>PLoS ONE</i> , 2020, 15, e0240418.	2.5	15
21	Lymphovascular space invasion as a predictive factor for lymph node metastases and survival in endometrioid endometrial cancer – a Swedish Gynecologic Cancer Group (SweGCG) study. <i>Acta Oncologica</i> , 2019, 58, 1628-1633.	1.8	55
22	Primary treatment patterns and survival of cervical cancer in Sweden: A population-based Swedish Gynecologic Cancer Group Study. <i>Gynecologic Oncology</i> , 2019, 155, 229-236.	1.4	12
23	Primary screening for increased fracture risk by the FRAX® questionnaire – uptake rates in relation to invitation method. <i>Archives of Osteoporosis</i> , 2019, 14, 51.	2.4	3
24	Population-based primary HPV mRNA cervical screening compared with cytology screening. <i>Preventive Medicine</i> , 2019, 124, 61-66.	3.4	11
25	Increased HPV detection by the use of a pre-heating step on vaginal self-samples analysed by Aptima HPV assay. <i>Journal of Virological Methods</i> , 2019, 270, 18-20.	2.1	8
26	Detection of HPV mRNA in Self-collected Vaginal Samples Among Women at 69-70 Years of Age. <i>Anticancer Research</i> , 2019, 39, 381-386.	1.1	8
27	Confounders other than comorbidity explain survival differences in Danish and Swedish ovarian cancer patients – a comparative cohort study. <i>Acta Oncologica</i> , 2018, 57, 1100-1108.	1.8	6
28	Self-sampling with HPV mRNA analyses from vagina and urine compared with cervical samples. <i>Journal of Clinical Virology</i> , 2018, 101, 69-73.	3.1	38
29	A Swedish population-based evaluation of benign hysterectomy, comparing minimally invasive and abdominal surgery. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2018, 222, 113-118.	1.1	25
30	Lymph node metastases as only qualifier for stage IV serous ovarian cancer confers longer survival than other sites of distant disease – a Swedish Gynecologic Cancer Group (SweGCG) study. <i>Acta Oncologica</i> , 2018, 57, 331-337.	1.8	17
31	Data quality in the Swedish Quality Register of Gynecologic Cancer – a Swedish Gynecologic Cancer Group (SweGCG) study. <i>Acta Oncologica</i> , 2018, 57, 346-353.	1.8	41
32	Prevalence of High-risk HPV in Postmenopausal Women with Benign Cervical Cytology – A Population-based Cohort Study. <i>Anticancer Research</i> , 2018, 38, 4221-4228.	1.1	8
33	Risk factors for lymph node metastases in women with endometrial cancer: A population-based, nationwide register study – On behalf of the Swedish Gynecological Cancer Group. <i>International Journal of Cancer</i> , 2017, 140, 2693-2700.	5.1	23
34	Increased blood cadmium levels were not associated with increased fracture risk but with increased total mortality in women: the Malmö Diet and Cancer Study. <i>Osteoporosis International</i> , 2017, 28, 2401-2408.	3.1	21
35	Population-based study of survival for women with serous cancer of the ovary, fallopian tube, peritoneum or undesignated origin - on behalf of the Swedish gynecological cancer group (SweGCG). <i>Gynecologic Oncology</i> , 2017, 144, 167-173.	1.4	23
36	Postmenopausal hormone therapy and risk of stroke: A pooled analysis of data from population-based cohort studies. <i>PLoS Medicine</i> , 2017, 14, e1002445.	8.4	44

#	ARTICLE	IF	CITATIONS
37	The economic burden of human papillomavirus-related precancers and cancers in Sweden. PLoS ONE, 2017, 12, e0179520.	2.5	16
38	Ovarian Cancer Surgery – A Population-based Registry Study. Anticancer Research, 2017, 37, 1837-1845.	1.1	9
39	Vaginal and Urine Self-sampling Compared to Cervical Sampling for HPV-testing with the Cobas 4800 HPV Test. Anticancer Research, 2017, 37, 4183-4187.	1.1	18
40	A population-based registry study evaluating surgery in newly diagnosed uterine cancer. Acta Obstetrica Et Gynecologica Scandinavica, 2016, 95, 901-911.	2.8	17
41	HE4 is a novel tissue marker for therapy response and progestin resistance in medium- and low-risk endometrial hyperplasia. British Journal of Cancer, 2016, 115, 725-730.	6.4	12
42	Follow up with HPV test and cytology as test of cure, 6 months after conization, is reliable. Acta Obstetrica Et Gynecologica Scandinavica, 2016, 95, 1251-1257.	2.8	21
43	Long-term survival in women with borderline ovarian tumors: a population-based survey of borderline ovarian tumors in Sweden 1960–2007. Acta Obstetrica Et Gynecologica Scandinavica, 2016, 95, 473-479.	2.8	17
44	A Biomarker Panel Increases the Diagnostic Performance for Epithelial Ovarian Cancer Type I and II in Young Women. Anticancer Research, 2016, 36, 957-65.	1.1	16
45	A novel diagnostic index combining HE4, CA125 and age may improve triage of women with suspected ovarian cancer – An international multicenter study in women with an ovarian mass. Gynecologic Oncology, 2015, 138, 640-646.	1.4	78
46	Robot-assisted surgery in cervical cancer patients reduces the time to normal activities of daily living. Acta Obstetrica Et Gynecologica Scandinavica, 2015, 94, 260-265.	2.8	21
47	A population-based study of pelvic serous carcinoma in Sweden: Primary site, FIGO stage and survival.. Journal of Clinical Oncology, 2015, 33, e16533-e16533.	1.6	0
48	Age influences the clinical significance of atypical glandular cells on cytology. Anticancer Research, 2015, 35, 913-9.	1.1	9
49	Use of proton pump inhibitors (PPI) and history of earlier fracture are independent risk factors for fracture in postmenopausal women. The WHILA study. Maturitas, 2014, 78, 310-315.	2.4	43
50	Elderly women above screening age diagnosed with cervical cancer have a worse prognosis. Anticancer Research, 2014, 34, 5147-51.	1.1	29
51	Low androstenedione/sex hormone binding globulin ratio increases fracture risk in postmenopausal women. The Women's Health in the Lund Area study. Maturitas, 2013, 75, 270-275.	2.4	9
52	Comparison of use of vaginal HPV self-sampling and offering flexible appointments as strategies to reach long-term non-attending women in organized cervical screening. Journal of Clinical Virology, 2013, 58, 155-160.	3.1	54
53	Vaginal self-sampling without preservative for human papillomavirus testing shows good sensitivity. Journal of Clinical Virology, 2013, 56, 52-56.	3.1	21
54	Reproducibility and accuracy of robot-assisted laparoscopic fertility sparing radical trachelectomy. Gynecologic Oncology, 2012, 127, 484-488.	1.4	67

#	ARTICLE	IF	CITATIONS
55	High preoperative blood levels of HE4 predicts poor prognosis in patients with ovarian cancer. <i>Journal of Ovarian Research</i> , 2012, 5, 20.	3.0	48
56	Robot-assisted removal of a lymphocyst causing severe neuralgic pain and adductor atrophy. <i>Journal of Robotic Surgery</i> , 2011, 5, 299-302.	1.8	0
57	The sentinel node concept in early cervical cancer performs well in tumors smaller than 2Åcm. <i>Gynecologic Oncology</i> , 2010, 117, 266-269.	1.4	75
58	Robot assisted laparoscopic radical hysterectomy and pelvic lymphadenectomy with short and long term morbidity data. <i>Gynecologic Oncology</i> , 2009, 113, 185-190.	1.4	100
59	Cleaved Forms of the Urokinase Plasminogen Activator Receptor in Plasma Have Diagnostic Potential and Predict Postoperative Survival in Patients with Ovarian Cancer. <i>Clinical Cancer Research</i> , 2008, 14, 5785-5793.	7.0	67
60	Fertility-sparing surgery and outcome in fertile women with ovarian borderline tumors and epithelial invasive ovarian cancer. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2007, 134, 110-114.	1.1	56
61	Background factors influencing somatic and psychological symptoms in middle-age women with different hormonal status. <i>Maturitas</i> , 2005, 52, 306-318.	2.4	38
62	Cancer risk after hospital discharge diagnosis of benign ovarian cysts and endometriosis. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2004, 83, 395-400.	2.8	115
63	Low-dose oral combination of 17 $\beta$ -estradiol and norethisterone acetate in postmenopausal women decreases factor VII, fibrinogen, antithrombin and plasminogen activator inhibitor-1. <i>Climacteric</i> , 2004, 7, 78-85.	2.4	19
64	Menopause-related symptoms: What are the background factors? A prospective population-based cohort study of Swedish women (The Women's Health in Lund Area study). <i>American Journal of Obstetrics and Gynecology</i> , 2003, 189, 1646-1653.	1.3	113
65	High preoperative plasma concentration of tissue plasminogen activator (tPA) is an independent marker for shorter overall survival in patients with ovarian cancer. <i>Gynecologic Oncology</i> , 2003, 91, 112-117.	1.4	12
66	High tumor tissue concentration of urokinase plasminogen activator receptor is associated with good prognosis in patients with ovarian cancer. <i>International Journal of Cancer</i> , 2003, 107, 658-665.	5.1	16
67	Low-dose hormone therapy and carbohydrate metabolism. <i>Fertility and Sterility</i> , 2003, 79, 550-555.	1.0	32
68	Changes in lipid and lipoprotein profile in postmenopausal women receiving low-dose combinations of 17 $\beta$ -estradiol and norethisterone acetate. <i>Menopause</i> , 2002, 9, 335-342.	2.0	20
69	Dedifferentiation of serous ovarian cancer from cystic to solid tumors is associated with increased expression of mRNA for urokinase plasminogen activator (uPA), its receptor (uPAR) and its inhibitor (PAI-1). <i>International Journal of Cancer</i> , 2001, 92, 497-502.	5.1	45
70	High tissue content of urokinase plasminogen activator (u-PA) is associated with high stromal expression of u-PA mRNA in poorly differentiated serous ovarian carcinoma. <i>International Journal of Cancer</i> , 1998, 79, 588-595.	5.1	20
71	Long-term incidence of endometrial cancer after endometrial resection and ablation: A population based Swedish gynecologic cancer group ( <scp>SweGCG</scp> ) study. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 0, , .	2.8	4