Stephen J Sharp

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8318151/publications.pdf

Version: 2024-02-01

38742 11,144 139 50 citations h-index papers

100 g-index 143 143 143 20933 docs citations times ranked citing authors all docs

32842

#	Article	IF	CITATIONS
1	Sedentary behaviour and risk of all-cause, cardiovascular and cancer mortality, and incident type 2 diabetes: a systematic review and dose response meta-analysis. European Journal of Epidemiology, 2018, 33, 811-829.	5.7	777
2	Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge. Nature Genetics, 2010, 42, 142-148.	21.4	591
3	HMG-coenzyme A reductase inhibition, type 2 diabetes, and bodyweight: evidence from genetic analysis and randomised trials. Lancet, The, 2015, 385, 351-361.	13.7	562
4	Association between alcohol and cardiovascular disease: Mendelian randomisation analysis based on individual participant data. BMJ, The, 2014, 349, g4164-g4164.	6.0	528
5	Integrative genomic analysis implicates limited peripheral adipose storage capacity in the pathogenesis of human insulin resistance. Nature Genetics, 2017, 49, 17-26.	21.4	452
6	Differences in the prospective association between individual plasma phospholipid saturated fatty acids and incident type 2 diabetes: the EPIC-InterAct case-cohort study. Lancet Diabetes and Endocrinology,the, 2014, 2, 810-818.	11.4	431
7	Physical distancing interventions and incidence of coronavirus disease 2019: natural experiment in 149 countries. BMJ, The, 2020, 370, m2743.	6.0	427
8	Effect of early intensive multifactorial therapy on 5-year cardiovascular outcomes in individuals with type 2 diabetes detected by screening (ADDITION-Europe): a cluster-randomised trial. Lancet, The, 2011, 378, 156-167.	13.7	406
9	Impact of common genetic determinants of Hemoglobin A1c on type 2 diabetes risk and diagnosis in ancestrally diverse populations: A transethnic genome-wide meta-analysis. PLoS Medicine, 2017, 14, e1002383.	8.4	341
10	Genetic Predisposition to an Impaired Metabolism of the Branched-Chain Amino Acids and Risk of Type 2 Diabetes: A Mendelian Randomisation Analysis. PLoS Medicine, 2016, 13, e1002179.	8.4	324
11	Association Between Low-Density Lipoprotein Cholesterol–Lowering Genetic Variants and Risk of Type 2 Diabetes. JAMA - Journal of the American Medical Association, 2016, 316, 1383.	7.4	310
12	Physical activity and all-cause mortality across levels of overall and abdominal adiposity in European men and women: the European Prospective Investigation into Cancer and Nutrition Study (EPIC). American Journal of Clinical Nutrition, 2015, 101, 613-621.	4.7	284
13	The UKâ€PBC risk scores: Derivation and validation of a scoring system for longâ€term prediction of endâ€stage liver disease in primary biliary cholangitis. Hepatology, 2016, 63, 930-950.	7.3	269
14	Human Gain-of-Function MC4R Variants Show Signaling Bias and Protect against Obesity. Cell, 2019, 177, 597-607.e9.	28.9	192
15	Fatty acids measured in plasma and erythrocyte-membrane phospholipids and derived by food-frequency questionnaire and the risk of new-onset type 2 diabetes: a pilot study in the European Prospective Investigation into Cancer and Nutrition (EPIC)–Norfolk cohort. American Journal of Clinical Nutrition, 2010, 92, 1214-1222.	4.7	190
16	The amount and type of dairy product intake and incident type 2 diabetes: results from the EPIC-InterAct Study. American Journal of Clinical Nutrition, 2012, 96, 382-390.	4.7	183
17	Gene-Lifestyle Interaction and Type 2 Diabetes: The EPIC InterAct Case-Cohort Study. PLoS Medicine, 2014, 11, e1001647.	8.4	180
18	Change in objectively measured physical activity during the transition to adolescence. British Journal of Sports Medicine, 2015, 49, 730-736.	6.7	175

#	Article	IF	CITATIONS
19	Association between circulating 25-hydroxyvitamin D and incident type 2 diabetes: a mendelian randomisation study. Lancet Diabetes and Endocrinology,the, 2015, 3, 35-42.	11.4	164
20	Wearable-device-measured physical activity and future health risk. Nature Medicine, 2020, 26, 1385-1391.	30.7	157
21	The spectrum effect in tests for risk prediction, screening, and diagnosis. BMJ, The, 2016, 353, i3139.	6.0	155
22	Common Genetic Variants Highlight the Role of Insulin Resistance and Body Fat Distribution in Type 2 Diabetes, Independent of Obesity. Diabetes, 2014, 63, 4378-4387.	0.6	153
23	Associations between Potentially Modifiable Risk Factors and Alzheimer Disease: A Mendelian Randomization Study. PLoS Medicine, 2015, 12, e1001841.	8.4	153
24	Association of Genetic Variants Related to Gluteofemoral vs Abdominal Fat Distribution With Type 2 Diabetes, Coronary Disease, and Cardiovascular Risk Factors. JAMA - Journal of the American Medical Association, 2018, 320, 2553.	7.4	152
25	Association of Plasma Phospholipid n-3 and n-6 Polyunsaturated Fatty Acids with Type 2 Diabetes: The EPIC-InterAct Case-Cohort Study. PLoS Medicine, 2016, 13, e1002094.	8.4	150
26	Dietary Protein Intake and Incidence of Type 2 Diabetes in Europe: The EPIC-InterAct Case-Cohort Study. Diabetes Care, 2014, 37, 1854-1862.	8.6	141
27	Prescription of glucose-lowering therapies and risk of COVID-19 mortality in people with type 2 diabetes: a nationwide observational study in England. Lancet Diabetes and Endocrinology,the, 2021, 9, 293-303.	11.4	140
28	Evidence of a Causal Association Between Insulinemia and Endometrial Cancer: A Mendelian Randomization Analysis. Journal of the National Cancer Institute, 2015, 107, .	6.3	129
29	Untargeted Metabolic Profiling Identifies Altered Serum Metabolites of Type 2 Diabetes Mellitus in a Prospective, Nested Case Control Study. Clinical Chemistry, 2015, 61, 487-497.	3.2	113
30	A genomic approach to therapeutic target validation identifies a glucose-lowering <i>GLP1R</i> variant protective for coronary heart disease. Science Translational Medicine, 2016, 8, 341ra76.	12.4	100
31	A Mendelian Randomization Study of Circulating Uric Acid and Type 2 Diabetes. Diabetes, 2015, 64, 3028-3036.	0.6	98
32	Use of the prevented fraction for the population to determine deaths averted by existing prevalence of physical activity: a descriptive study. The Lancet Global Health, 2020, 8, e920-e930.	6.3	86
33	Age-related patterns of vigorous-intensity physical activity in youth: The International Children's Accelerometry Database. Preventive Medicine Reports, 2016, 4, 17-22.	1.8	84
34	Residential neighbourhood greenspace is associated with reduced risk of incident diabetes in older people: a prospective cohort study. BMC Public Health, 2016, 16, 1171.	2.9	80
35	Assessing the impact of the Barbados sugar-sweetened beverage tax on beverage sales: an observational study. International Journal of Behavioral Nutrition and Physical Activity, 2019, 16, 13.	4.6	75
36	Association of plasma biomarkers of fruit and vegetable intake with incident type 2 diabetes: EPIC-InterAct case-cohort study in eight European countries. BMJ, The, 2020, 370, m2194.	6.0	75

#	Article	IF	CITATIONS
37	The absolute and relative risk of type 2 diabetes after gestational diabetes: A systematic review and meta-analysis of 129 studies. Diabetes Research and Clinical Practice, 2021, 171, 108625.	2.8	7 5
38	Alcohol intake in relation to non-fatal and fatal coronary heart disease and stroke: EPIC-CVD case-cohort study. BMJ: British Medical Journal, 2018, 361, k934.	2.3	70
39	Plasma Vitamin C and Type 2 Diabetes: Genome-Wide Association Study and Mendelian Randomization Analysis in European Populations. Diabetes Care, 2021, 44, 98-106.	8.6	68
40	Association of Multiple Biomarkers of Iron Metabolism and Type 2 Diabetes: The EPIC-InterAct Study. Diabetes Care, 2016, 39, 572-581.	8.6	65
41	The combination of cardiorespiratory fitness and muscle strength, and mortality risk. European Journal of Epidemiology, 2018, 33, 953-964.	5.7	64
42	Definitions of Metabolic Health and Risk of Future Type 2 Diabetes in BMI Categories: A Systematic Review and Network Meta-analysis. Diabetes Care, 2015, 38, 2177-2187.	8.6	61
43	A combination of plasma phospholipid fatty acids and its association with incidence of type 2 diabetes: The EPIC-InterAct case-cohort study. PLoS Medicine, 2017, 14, e1002409.	8.4	61
44	Thirdâ€wave cognitive behaviour therapies for weight management: A systematic review and network metaâ€analysis. Obesity Reviews, 2020, 21, e13013.	6.5	61
45	Tea Consumption and Incidence of Type 2 Diabetes in Europe: The EPIC-InterAct Case-Cohort Study. PLoS ONE, 2012, 7, e36910.	2.5	59
46	Impact of follow-up time and analytical approaches to account for reverse causality on the association between physical activity and health outcomes in UK Biobank. International Journal of Epidemiology, 2020, 49, 162-172.	1.9	57
47	Effect of Early Multifactorial Therapy Compared With Routine Care on Microvascular Outcomes at 5 Years in People With Screen-Detected Diabetes: A Randomized Controlled Trial. Diabetes Care, 2014, 37, 2015-2023.	8.6	56
48	Lifestyle Advice Combined with Personalized Estimates of Genetic or Phenotypic Risk of Type 2 Diabetes, and Objectively Measured Physical Activity: A Randomized Controlled Trial. PLoS Medicine, 2016, 13, e1002185.	8.4	55
49	Association of Genetically Enhanced Lipoprotein Lipase–Mediated Lipolysis and Low-Density Lipoprotein Cholesterol–Lowering Alleles With Risk of Coronary Disease and Type 2 Diabetes. JAMA Cardiology, 2018, 3, 957.	6.1	55
50	Association between plasma phospholipid saturated fatty acids and metabolic markers of lipid, hepatic, inflammation and glycaemic pathways in eight European countries: a cross-sectional analysis in the EPIC-InterAct study. BMC Medicine, 2017, 15, 203.	5.5	47
51	Supermarket policies on less-healthy food at checkouts: Natural experimental evaluation using interrupted time series analyses of purchases. PLoS Medicine, 2018, 15, e1002712.	8.4	47
52	Interaction between genes and macronutrient intake on the risk of developing type 2 diabetes: systematic review and findings from European Prospective Investigation into Cancer (EPIC)-InterAct. American Journal of Clinical Nutrition, 2017, 106, 263-275.	4.7	46
53	Meta-analysis investigating the role of interleukin-6 mediated inflammation in type 2 diabetes. EBioMedicine, 2020, 61, 103062.	6.1	46
54	The association between circulating 25-hydroxyvitamin D metabolites and type 2 diabetes in European populations: AÂmeta-analysis and Mendelian randomisation analysis. PLoS Medicine, 2020, 17, e1003394.	8.4	45

#	Article	IF	CITATIONS
55	Population level physical activity before and during the first national COVID-19 lockdown: A nationally representative repeat cross-sectional study of 5 years of Active Lives data in England. Lancet Regional Health - Europe, The, 2022, 12, 100265.	5. 6	44
56	Long-term effects of intensive multifactorial therapy in individuals with screen-detected type 2 diabetes in primary care: 10-year follow-up of the ADDITION-Europe cluster-randomised trial. Lancet Diabetes and Endocrinology,the, 2019, 7, 925-937.	11.4	39
57	A pragmatic and scalable strategy using mobile technology to promote sustained lifestyle changes to prevent type 2 diabetes in India and the UK: a randomised controlled trial. Diabetologia, 2020, 63, 486-496.	6.3	38
58	The impact of adult behavioural weight management interventions on mental health: A systematic review and metaâ€analysis. Obesity Reviews, 2021, 22, e13150.	6.5	38
59	Low Bone Mineral Density Predicts Incident Heart Failure in Men and Women. JACC: Heart Failure, 2014, 2, 380-389.	4.1	37
60	Substituting prolonged sedentary time and cardiovascular risk in children and youth: a meta-analysis within the International Children's Accelerometry database (ICAD). International Journal of Behavioral Nutrition and Physical Activity, 2019, 16, 96.	4.6	35
61	Magnitude and determinants of change in objectively-measured physical activity, sedentary time and sleep duration from ages 15 to 17.5y in UK adolescents: the ROOTS study. International Journal of Behavioral Nutrition and Physical Activity, 2015, 12, 61.	4.6	34
62	Effect of communicating phenotypic and genetic risk of coronary heart disease alongside web-based lifestyle advice: the INFORM Randomised Controlled Trial. Heart, 2019, 105, 982-989.	2.9	34
63	Galanin inhibits GLPâ€1 and GIP secretion via the GAL ₁ receptor in enteroendocrine L and K cells. British Journal of Pharmacology, 2016, 173, 888-898.	5.4	33
64	Using alternatives to the car and risk of all-cause, cardiovascular and cancer mortality. Heart, 2018, 104, 1749-1755.	2.9	32
65	Mortality Risk Reductions from Substituting Screen Time by Discretionary Activities. Medicine and Science in Sports and Exercise, 2017, 49, 1111-1119.	0.4	30
66	Dairy Product Intake and Risk of Type 2 Diabetes in EPIC-InterAct: A Mendelian Randomization Study. Diabetes Care, 2019, 42, 568-575.	8.6	29
67	Effectiveness and cost-effectiveness of the GoActive intervention to increase physical activity among UK adolescents: AÂcluster randomised controlled trial. PLoS Medicine, 2020, 17, e1003210.	8.4	28
68	Metformin in non-diabetic hyperglycaemia: the GLINT feasibility RCT. Health Technology Assessment, 2018, 22, 1-64.	2.8	28
69	Effect of interventions incorporating personalised cancer risk information on intentions and behaviour: a systematic review and meta-analysis of randomised controlled trials. BMJ Open, 2018, 8, e017717.	1.9	26
70	Association of Plasma Vitamin D Metabolites With Incident Type 2 Diabetes: EPIC-InterAct Case-Cohort Study. Journal of Clinical Endocrinology and Metabolism, 2019, 104, 1293-1303.	3.6	25
71	Estimated Substitution of Tea or Coffee for Sugar-Sweetened Beverages Was Associated with Lower Type 2 Diabetes Incidence in Case–Cohort Analysis across 8 European Countries in the EPIC-InterAct Study. Journal of Nutrition, 2019, 149, 1985-1993.	2.9	24
72	The Relationship between Self-Reported Exposure to Sugar-Sweetened Beverage Promotions and Intake: Cross-Sectional Analysis of the 2017 International Food Policy Study. Nutrients, 2019, 11, 3047.	4.1	24

#	Article	IF	Citations
73	A cluster randomised controlled trial to evaluate the effectiveness and cost-effectiveness of the GoActive intervention to increase physical activity among adolescents aged $13\hat{a}\in 14$ years. BMJ Open, 2017, 7, e014419.	1.9	23
74	Impact of Personalised Feedback about Physical Activity on Change in Objectively Measured Physical Activity (the FAB Study): A Randomised Controlled Trial. PLoS ONE, 2013, 8, e75398.	2.5	21
75	The independent prospective associations of activity intensity and dietary energy density with adiposity in young adolescents. British Journal of Nutrition, 2016, 115, 921-929.	2.3	21
76	A multilevel linear mixed model of the association between candidate genes and weight and body mass index using the Framingham longitudinal family data. BMC Proceedings, 2009, 3, S115.	1.6	20
77	Effect of communicating genetic and phenotypic risk for type 2 diabetes in combination with lifestyle advice on objectively measured physical activity: protocol of a randomised controlled trial. BMC Public Health, 2012, 12, 444.	2.9	20
78	The pathway to diagnosis of type 1 diabetes in children: a questionnaire study. BMJ Open, 2015, 5, e006470-e006470.	1.9	20
79	Interplay between genetic predisposition, macronutrient intake and type 2 diabetes incidence: analysis within EPIC-InterAct across eight European countries. Diabetologia, 2018, 61, 1325-1332.	6.3	20
80	Genome-wide association analysis of type 2 diabetes in the EPIC-InterAct study. Scientific Data, 2020, 7, 393.	5.3	19
81	Estimating physical activity from self-reported behaviours in large-scale population studies using network harmonisation: findings from UK Biobank and associations with disease outcomes. International Journal of Behavioral Nutrition and Physical Activity, 2020, 17, 40.	4.6	18
82	Circulating Fetuin-A and Risk of Type 2 Diabetes: A Mendelian Randomization Analysis. Diabetes, 2018, 67, 1200-1205.	0.6	17
83	Changes in plasma phospholipid fatty acid profiles over 13 years and correlates of change: European Prospective Investigation into Cancer and Nutrition-Norfolk Study. American Journal of Clinical Nutrition, 2019, 109, 1527-1534.	4.7	17
84	Associations of genetic susceptibility and healthy lifestyle with incidence of coronary heart disease and stroke in individuals with hypertension. European Journal of Preventive Cardiology, 2022, 29, 2101-2110.	1.8	17
85	Randomised controlled trial of a theory-based behavioural intervention to reduce formula milk intake. Archives of Disease in Childhood, 2018, 103, archdischild-2018-314784.	1.9	16
86	Is occupational physical activity associated with mortality in UK Biobank?. International Journal of Behavioral Nutrition and Physical Activity, 2021, 18, 102.	4.6	16
87	Tinned Fruit Consumption and Mortality in Three Prospective Cohorts. PLoS ONE, 2015, 10, e0117796.	2.5	15
88	Interaction of Dietary and Genetic Factors Influencing Body Iron Status and Risk of Type 2 Diabetes Within the EPIC-InterAct Study. Diabetes Care, 2018, 41, 277-285.	8.6	15
89	Autoimmunity plays a role in the onset of diabetes after 40 years of age. Diabetologia, 2020, 63, 266-277.	6.3	15
90	Plasma vitamin C and risk of hospitalisation with diagnosis of atrial fibrillation in men and women in EPIC-Norfolk prospective study. International Journal of Cardiology, 2014, 177, 830-835.	1.7	14

#	Article	IF	CITATIONS
91	Promoting physical activity in a multi-ethnic population at high risk of diabetes: the 48-month PROPELS randomised controlled trial. BMC Medicine, 2021, 19, 130.	5.5	14
92	Information and Risk Modification Trial (INFORM): design of a randomised controlled trial of communicating different types of information about coronary heart disease risk, alongside lifestyle advice, to achieve change in health-related behaviour. BMC Public Health, 2015, 15, 868.	2.9	13
93	Development and Validation of Lifestyle-Based Models to Predict Incidence of the Most Common Potentially Preventable Cancers. Cancer Epidemiology Biomarkers and Prevention, 2019, 28, 67-75.	2.5	13
94	The impact of participant mental health on attendance and engagement in a trial of behavioural weight management programmes: secondary analysis of the WRAP randomised controlled trial. International Journal of Behavioral Nutrition and Physical Activity, 2021, 18, 146.	4.6	13
95	Effect of interventions including provision of personalised cancer risk information on accuracy of risk perception and psychological responses: A systematic review and meta-analysis. Patient Education and Counseling, 2020, 103, 83-95.	2.2	12
96	A randomised trial of the effect and cost-effectiveness of early intensive multifactorial therapy on 5-year cardiovascular outcomes in individuals with screen-detected type 2 diabetes: the Anglo–Danish–Dutch Study of Intensive Treatment in People with Screen-Detected Diabetes in Primary Care (ADDITION-Europe) study. Health Technology Assessment, 2016, 20, 1-86.	2.8	12
97	Risk models for recurrence and survival after kidney cancer: a systematic review. BJU International, 2022, 130, 562-579.	2.5	12
98	The Association of Low-To-Moderate Alcohol Consumption with Breast Cancer Subtypes Defined by Hormone Receptor Status. PLoS ONE, 2015, 10, e0144680.	2. 5	11
99	Reply to H Pareja-Galeano et al American Journal of Clinical Nutrition, 2015, 101, 1101.	4.7	11
100	Inverse association between bone mineral density and risk of aortic stenosis in men and women in EPIC–Norfolk prospective study. International Journal of Cardiology, 2015, 178, 29-30.	1.7	11
101	A simple risk score using routine data for predicting cardiovascular disease in primary care. British Journal of General Practice, 2010, 60, e327-e334.	1.4	10
102	A method making fewer assumptions gave the most reliable estimates of exposure–outcome associations in stratified case–cohort studies. Journal of Clinical Epidemiology, 2015, 68, 1397-1405.	5.0	10
103	Protocol for Get Moving: a randomised controlled trial to assess the effectiveness of three minimal contact interventions to promote fitness and physical activity in working adults. BMC Public Health, 2015, 15, 296.	2.9	10
104	Anticipatory changes in British household purchases of soft drinks associated with the announcement of the Soft Drinks Industry Levy: A controlled interrupted time series analysis. PLoS Medicine, 2020, 17, e1003269.	8.4	10
105	Repeat Cardiovascular Risk Assessment after Four Years: Is There Improvement in Risk Prediction?. PLoS ONE, 2016, 11, e0147417.	2.5	9
106	Socio-economic and age variations in response to supermarket-led checkout food policies: a repeated measures analysis. International Journal of Behavioral Nutrition and Physical Activity, 2018, 15, 125.	4.6	9
107	Heterogeneity of Associations between Total and Types of Fish Intake and the Incidence of Type 2 Diabetes: Federated Meta-Analysis of 28 Prospective Studies Including 956,122 Participants. Nutrients, 2021, 13, 1223.	4.1	8
108	A randomised controlled trial of the effect of providing online risk information and lifestyle advice for the most common preventable cancers. Preventive Medicine, 2020, 138, 106154.	3.4	7

#	Article	IF	CITATIONS
109	Independent and combined associations between fast-food outlet exposure and genetic risk for obesity: a population-based, cross-sectional study in the UK. BMC Medicine, 2021, 19, 49.	5.5	7
110	A school-based, peer-led programme to increase physical activity among 13- to 14-year-old adolescents: the GoActive cluster RCT. Public Health Research, 2021, 9, 1-134.	1.3	7
111	Behavioural interventions to promote physical activity in a multiethnic population at high risk of diabetes: PROPELS three-arm RCT. Health Technology Assessment, 2021, 25, 1-190.	2.8	7
112	Supporting Weight Management during COVID-19: A Randomized Controlled Trial of a Web-Based, ACT-Based, Guided Self-Help Intervention. Obesity Facts, 2022, 15, 550-559.	3.4	7
113	Generalizability of a Diabetes-Associated Country-Specific Exploratory Dietary Pattern Is Feasible Across European Populations. Journal of Nutrition, 2019, 149, 1047-1055.	2.9	6
114	Association of weight loss and weight loss maintenance following diabetes diagnosis by screening and incidence of cardiovascular disease and allâ€cause mortality: An observational analysis of the ADDITIONâ€Europe trial. Diabetes, Obesity and Metabolism, 2021, 23, 730-741.	4.4	6
115	Interaction Between GAD65 Antibodies and Dietary Fish Intake or Plasma Phospholipid n-3 Polyunsaturated Fatty Acids on Incident Adult-Onset Diabetes: The EPIC-InterAct Study. Diabetes Care, 2021, 44, 416-424.	8.6	6
116	Network Harmonization of Physical Activity Variables Through Indirect Validation. Journal for the Measurement of Physical Behaviour, 2020, 3, 8-18.	0.8	6
117	A randomised controlled trial of the effect of providing online risk information and lifestyle advice for the most common preventable cancers: study protocol. BMC Public Health, 2018, 18, 796.	2.9	5
118	Development and multi-cohort validation of a clinical score for predicting type 2 diabetes mellitus. PLoS ONE, 2019, 14, e0218933.	2.5	5
119	Longitudinal associations between prepubertal childhood total energy and macronutrient intakes and subsequent puberty timing in UK boys and girls. European Journal of Nutrition, 2022, 61, 157-167.	3.9	5
120	Protocol for a clinical trial of text messaging in addition to standard care versus standard care alone in prevention of type 2 diabetes through lifestyle modification in India and the UK. BMC Endocrine Disorders, 2018, 18, 63.	2.2	3
121	Effectiveness of Minimal Contact Interventions: An RCT. American Journal of Preventive Medicine, 2021, 60, e111-e121.	3.0	3
122	Positive maternal attitudes to following healthy infant feeding guidelines attenuate the associations between infant appetitive traits and both infant milk intake and weight. Appetite, 2021, 161, 105124.	3.7	2
123	Association between patient activation, self-management behaviours and clinical outcomes in adults with diabetes or related metabolic disorders: a systematic review and meta-analysis protocol. BMJ Open, 2022, 12, e056293.	1.9	2
124	Acceptability and feasibility of an acceptance and commitment therapy-based guided self-help intervention for weight loss maintenance in adults who have previously completed a behavioural weight loss programme: the SWiM feasibility study protocol. BMJ Open, 2022, 12, e058103.	1.9	2
125	Association between 25-hydroxyvitamin D and type 2 diabetes – Authors' reply. Lancet Diabetes and Endocrinology,the, 2015, 3, 11-12.	11.4	1
126	Impact of achievement and change in achievement of lifestyle recommendations in middle-age on risk of the most common potentially preventable cancers. Preventive Medicine, 2021, 153, 106712.	3.4	1

#	Article	IF	CITATIONS
127	Impacts of new cycle infrastructure on cycling levels in two French cities: an interrupted time series analysis. International Journal of Behavioral Nutrition and Physical Activity, 2022, 19, .	4.6	1
128	Participant Characteristics Associated with Changes in Mental Health in a Trial of Behavioural Weight Management Programmes: Secondary Analysis of the WRAP Trial. Obesity Facts, 2022, 15, 508-518.	3.4	0
129	Title is missing!. , 2020, 17, e1003210.		O
130	Title is missing!. , 2020, 17, e1003210.		0
131	Title is missing!. , 2020, 17, e1003210.		O
132	Title is missing!. , 2020, 17, e1003210.		0
133	Title is missing!. , 2020, 17, e1003210.		O
134	Title is missing!. , 2020, 17, e1003210.		0
135	Title is missing!. , 2020, 17, e1003269.		O
136	Title is missing!. , 2020, 17, e1003269.		0
137	Title is missing!. , 2020, 17, e1003269.		O
138	Title is missing!. , 2020, 17, e1003269.		0
139	Title is missing!. , 2020, 17, e1003269.		O