

Veli-Matti KÃ¤hÃ¤ri

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8340493/publications.pdf>

Version: 2024-02-01

185
papers

15,223
citations

18482

62
h-index

19749

117
g-index

187
all docs

187
docs citations

187
times ranked

14849
citing authors

#	ARTICLE	IF	CITATIONS
1	Regulation of matrix metalloproteinase expression in tumor invasion. <i>FASEB Journal</i> , 1999, 13, 781-792.	0.5	1,390
2	CIP2A Inhibits PP2A in Human Malignancies. <i>Cell</i> , 2007, 130, 51-62.	28.9	662
3	Matrix metalloproteinases in cancer: Prognostic markers and therapeutic targets. <i>International Journal of Cancer</i> , 2002, 99, 157-166.	5.1	547
4	Matrix metalloproteinases in skin. <i>Experimental Dermatology</i> , 1997, 6, 199-213.	2.9	516
5	Trends in Molecular Medicine: Matrix metalloproteinases and their inhibitors in tumour growth and invasion. <i>Annals of Medicine</i> , 1999, 31, 34-45.	3.8	390
6	Matrix metalloproteinases in inflammation. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2014, 1840, 2571-2580.	2.4	344
7	Matrix metalloproteinases in tumor invasion. <i>Cellular and Molecular Life Sciences</i> , 2000, 57, 5-15.	5.4	295
8	MAPK/ERK Overrides the Apoptotic Signaling from Fas, TNF, and TRAIL Receptors. <i>Journal of Biological Chemistry</i> , 2001, 276, 16484-16490.	3.4	287
9	Collagenases in cancer. <i>Biochimie</i> , 2005, 87, 273-286.	2.6	277
10	Integrin $\alpha 2 \beta 1$ Is a Positive Regulator of Collagenase (MMP-1) and Collagen $\alpha 1(I)$ Gene Expression. <i>Journal of Biological Chemistry</i> , 1995, 270, 13548-13552.	3.4	263
11	Collagenase-3 (MMP-13) is expressed by hypertrophic chondrocytes, periosteal cells, and osteoblasts during human fetal bone development. , 1997, 208, 387-397.		262
12	Matrix Metalloproteinases as Therapeutic Targets in Cancer. <i>Current Cancer Drug Targets</i> , 2005, 5, 203-220.	1.6	253
13	Induction of Collagenase-3 (MMP-13) Expression in Human Skin Fibroblasts by Three-dimensional Collagen Is Mediated by p38 Mitogen-activated Protein Kinase. <i>Journal of Biological Chemistry</i> , 1999, 274, 2446-2455.	3.4	248
14	Distinct Populations of Stromal Cells Express Collagenase-3 (MMP-13) and Collagenase-1 (MMP-1) in Chronic Ulcers but Not in Normally Healing Wounds. <i>Journal of Investigative Dermatology</i> , 1997, 109, 96-101.	0.7	233
15	Integrin $\alpha 2 \beta 1$ Mediates Isoform-Specific Activation of p38 and Upregulation of Collagen Gene Transcription by a Mechanism Involving the $\alpha 2$ Cytoplasmic Tail. <i>Journal of Cell Biology</i> , 1999, 147, 401-416.	5.2	206
16	Activation of p38 β MAPK Enhances Collagenase-1 (Matrix Metalloproteinase (MMP)-1) and Stromelysin-1 (MMP-3) Expression by mRNA Stabilization. <i>Journal of Biological Chemistry</i> , 2002, 277, 32360-32368.	3.4	195
17	Transforming Growth Factor- $\beta 2$ Induces Collagenase-3 Expression by Human Gingival Fibroblasts via p38 Mitogen-activated Protein Kinase. <i>Journal of Biological Chemistry</i> , 1999, 274, 37292-37300.	3.4	191
18	Enhancement of Fibroblast Collagenase (Matrix Metalloproteinase-1) Gene Expression by Ceramide Is Mediated by Extracellular Signal-regulated and Stress-activated Protein Kinase Pathways. <i>Journal of Biological Chemistry</i> , 1998, 273, 5137-5145.	3.4	184

#	ARTICLE	IF	CITATIONS
19	p38 Mitogen-Activated Protein Kinase-Dependent Activation of Protein Phosphatases 1 and 2A Inhibits MEK1 and MEK2 Activity and Collagenase 1 (MMP-1) Gene Expression. <i>Molecular and Cellular Biology</i> , 2001, 21, 2373-2383.	2.3	183
20	Transforming growth factor- β 2 signaling in cancer invasion and metastasis. <i>International Journal of Cancer</i> , 2007, 121, 2119-2124.	5.1	179
21	Tumor necrosis factor-alpha and interferon-gamma suppress the activation of human type I collagen gene expression by transforming growth factor-beta 1. Evidence for two distinct mechanisms of inhibition at the transcriptional and posttranscriptional levels. <i>Journal of Clinical Investigation</i> , 1990, 86, 1489-1495.	8.2	170
22	Regulation of Membrane-Type Matrix Metalloproteinase-1 Expression by Growth Factors and Phorbol 12-Myristate 13-Acetate. <i>FEBS Journal</i> , 1996, 239, 239-247.	0.2	167
23	Identification of Fibroblasts Responsible for Increased Collagen Production in Localized Scleroderma by In Situ Hybridization. <i>Journal of Investigative Dermatology</i> , 1988, 90, 664-670.	0.7	164
24	Integrin α 2 β 1 Promotes Activation of Protein Phosphatase 2A and Dephosphorylation of Akt and Glycogen Synthase Kinase 3 β . <i>Molecular and Cellular Biology</i> , 2002, 22, 1352-1359.	2.3	164
25	Tissue inhibitor of metalloproteinases-3 induces apoptosis in melanoma cells by stabilization of death receptors. <i>Oncogene</i> , 2003, 22, 2121-2134.	5.9	162
26	High Serum Levels of Matrix Metalloproteinase-9 and Matrix Metalloproteinase-1 Are Associated with Rapid Progression in Patients with Metastatic Melanoma. <i>Clinical Cancer Research</i> , 2005, 11, 5158-5166.	7.0	161
27	Proteinases in cutaneous wound healing. <i>Cellular and Molecular Life Sciences</i> , 2009, 66, 203-224.	5.4	161
28	Human Collagenase-3 Is Expressed in Malignant Squamous Epithelium of the Skin. <i>Journal of Investigative Dermatology</i> , 1997, 109, 225-231.	0.7	150
29	European Dermatology Forum S1 guideline on the diagnosis and treatment of sclerosing diseases of the skin, Part 1: localized scleroderma, systemic sclerosis and overlap syndromes. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2017, 31, 1401-1424.	2.4	148
30	Evaluation of Transforming Growth Factor β 2 and Type I Procollagen Gene Expression in Fibrotic Skin Disease by In Situ Hybridization. <i>Journal of Investigative Dermatology</i> , 1990, 94, 365-371.	0.7	146
31	Differential regulation of interstitial collagenase (MMP-1) gene expression by ETS transcription factors. <i>Oncogene</i> , 1997, 14, 2651-2660.	5.9	136
32	EGF-R regulates MMP function in fibroblasts through MAPK and AP-1 pathways. <i>Journal of Cellular Physiology</i> , 2007, 212, 489-497.	4.1	133
33	p38 α and p38 β mitogen-activated protein kinase isoforms regulate invasion and growth of head and neck squamous carcinoma cells. <i>Oncogene</i> , 2007, 26, 5267-5279.	5.9	122
34	Interleukin-1 increases collagen production and mRNA levels in cultured skin fibroblasts. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1987, 929, 142-147.	4.1	119
35	Collagenase-3 (MMP-13) is Expressed by Tumor Cells in Invasive Vulvar Squamous Cell Carcinomas. <i>American Journal of Pathology</i> , 1999, 154, 469-480.	3.8	119
36	Senescence Sensitivity of Breast Cancer Cells Is Defined by Positive Feedback Loop between CIP2A and E2F1. <i>Cancer Discovery</i> , 2013, 3, 182-197.	9.4	117

#	ARTICLE	IF	CITATIONS
37	A Role for Decorin in the Structural Organization of Periodontal Ligament. <i>Laboratory Investigation</i> , 2000, 80, 1869-1880.	3.7	112
38	Smad3 and Extracellular Signal-Regulated Kinase 1/2 Coordinately Mediate Transforming Growth Factor- β -Induced Expression of Connective Tissue Growth Factor in Human Fibroblasts. <i>Journal of Investigative Dermatology</i> , 2005, 124, 1162-1169.	0.7	111
39	High expression levels of collagenase-1 and stromelysin-1 correlate with shorter disease-free survival in human metastatic melanoma. <i>International Journal of Cancer</i> , 2002, 97, 432-438.	5.1	108
40	Comparative Effects of Interleukin-1 and Tumor Necrosis Factor- α on Collagen Production and Corresponding Procollagen mRNA Levels in Human Dermal Fibroblasts. <i>Journal of Investigative Dermatology</i> , 1991, 96, 243-249.	0.7	104
41	Expression Profiles and Clinical Correlations of Degradome Components in the Tumor Microenvironment of Head and Neck Squamous Cell Carcinoma. <i>Clinical Cancer Research</i> , 2010, 16, 2022-2035.	7.0	100
42	Smad3 Mediates Transforming Growth Factor- β -induced Collagenase-3 (Matrix Metalloproteinase-13) Expression in Human Gingival Fibroblasts. <i>Journal of Biological Chemistry</i> , 2002, 277, 46338-46346.	3.4	93
43	Endothelial cell-Matrix interactions. <i>Microscopy Research and Technique</i> , 2003, 60, 13-22.	2.2	92
44	Activation of Smad signaling enhances collagenase-3 (MMP-13) expression and invasion of head and neck squamous carcinoma cells. <i>Oncogene</i> , 2006, 25, 2588-2600.	5.9	89
45	Expression of Human Macrophage Metalloelastase (MMP-12) by Tumor Cells in Skin Cancer. <i>Journal of Investigative Dermatology</i> , 2000, 114, 1113-1119.	0.7	88
46	Metalloelastase (MMP-12) expression by tumour cells in squamous cell carcinoma of the vulva correlates with invasiveness, while that by macrophages predicts better outcome. <i>Journal of Pathology</i> , 2002, 198, 258-269.	4.5	88
47	Epidermal growth factor increases collagen production in granulation tissue by stimulation of fibroblast proliferation and not by activation of procollagen genes. <i>Biochemical Journal</i> , 1987, 247, 385-388.	3.7	87
48	MMP-13 Regulates Growth of Wound Granulation Tissue and Modulates Gene Expression Signatures Involved in Inflammation, Proteolysis, and Cell Viability. <i>PLoS ONE</i> , 2012, 7, e42596.	2.5	87
49	A metaphyseal defect model of the femur for studies of murine bone healing. <i>Bone</i> , 2001, 28, 423-429.	2.9	84
50	Isoform-Specific Regulation of the Actin-Organizing Protein Palladin during TGF- β 1-Induced Myofibroblast Differentiation. <i>Journal of Investigative Dermatology</i> , 2006, 126, 2387-2396.	0.7	83
51	European dermatology forum S1 guideline on the diagnosis and treatment of sclerosing diseases of the skin, Part 2: Scleromyxedema, scleredema and nephrogenic systemic fibrosis. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2017, 31, 1581-1594.	2.4	79
52	Enhancement of fibroblast collagenase-1 (MMP-1) gene expression by tumor promoter okadaic acid is mediated by stress-activated protein kinases jun N-terminal kinase and p38. <i>Matrix Biology</i> , 1998, 17, 547-557.	3.6	78
53	Antitumor Activity and Bystander Effect of Adenovirally Delivered Tissue Inhibitor of Metalloproteinases-3. <i>Molecular Therapy</i> , 2002, 5, 705-715.	8.2	75
54	Coordinated regulation of type I and type III collagen production and mRNA levels of pro α 1(I) and pro α 2(I) collagen in cultured morphea fibroblasts. <i>Archives of Dermatological Research</i> , 1987, 279, 154-160.	1.9	73

#	ARTICLE	IF	CITATIONS
55	Expression of matrix metalloproteinase (MMP)-7 and MMP-13 and loss of MMP-19 and p16 are associated with malignant progression in chronic wounds. <i>British Journal of Dermatology</i> , 2005, 152, 720-726.	1.5	73
56	Complement Factor H: A Biomarker for Progression of Cutaneous Squamous Cell Carcinoma. <i>Journal of Investigative Dermatology</i> , 2014, 134, 498-506.	0.7	73
57	p38 Mitogen-activated protein kinase pathway suppresses cell survival by inducing dephosphorylation of mitogen-activated protein/extracellular signal-regulated kinase 1,2. <i>Cancer Research</i> , 2003, 63, 3473-7.	0.9	73
58	Targeted inhibition of human collagenase-3 (MMP-13) expression inhibits squamous cell carcinoma growth in vivo. <i>Oncogene</i> , 2004, 23, 5111-5123.	5.9	70
59	Requirements for Receptor Engagement during Infection by Adenovirus Complexed with Blood Coagulation Factor X. <i>PLoS Pathogens</i> , 2010, 6, e1001142.	4.7	70
60	Serpin Peptidase Inhibitor Clade A Member 1 (SerpinA1) Is a Novel Biomarker for Progression of Cutaneous Squamous Cell Carcinoma. <i>American Journal of Pathology</i> , 2011, 179, 1110-1119.	3.8	69
61	Complement Factor I Promotes Progression of Cutaneous Squamous Cell Carcinoma. <i>Journal of Investigative Dermatology</i> , 2015, 135, 579-588.	0.7	68
62	Increased type I collagen mRNA levels in cultured scleroderma fibroblasts. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1984, 781, 183-186.	2.4	66
63	Matrix metalloproteinase-7 activates heparin-binding epidermal growth factor-like growth factor in cutaneous squamous cell carcinoma. <i>British Journal of Dermatology</i> , 2010, 163, 726-735.	1.5	66
64	Interferon- β and interferon- γ reduce excessive collagen synthesis and procollagen mRNA levels of scleroderma fibroblasts in culture. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1988, 968, 45-50.	4.1	64
65	Adenoviral delivery of p53 gene suppresses expression of collagenase-3 (MMP-13) in squamous carcinoma cells. <i>Oncogene</i> , 2002, 21, 1187-1195.	5.9	64
66	Suppression of TGF β 2 and Angiogenesis by Type VII Collagen in Cutaneous SCC. <i>Journal of the National Cancer Institute</i> , 2016, 108, djv293.	6.3	63
67	Complement Component C3 and Complement Factor B Promote Growth of Cutaneous Squamous Cell Carcinoma. <i>American Journal of Pathology</i> , 2017, 187, 1186-1197.	3.8	63
68	Activation of Tissue Inhibitor of Metalloproteinases-3 (TIMP-3) mRNA Expression in Scleroderma Skin Fibroblasts. <i>Journal of Investigative Dermatology</i> , 1998, 110, 416-421.	0.7	62
69	Accelerated Up-Regulation of L-Sox5, Sox6, and Sox9 by BMP-2 Gene Transfer During Murine Fracture Healing. <i>Journal of Bone and Mineral Research</i> , 2001, 16, 1837-1845.	2.8	62
70	Long Noncoding RNA PICSAR Promotes Growth of Cutaneous Squamous Cell Carcinoma by Regulating ERK1/2 Activity. <i>Journal of Investigative Dermatology</i> , 2016, 136, 1701-1710.	0.7	61
71	New perspectives on role of tumor microenvironment in progression of cutaneous squamous cell carcinoma. <i>Cell and Tissue Research</i> , 2016, 365, 691-702.	2.9	60
72	Expression of human collagenase-3 (MMP-13) by fetal skin fibroblasts is induced by transforming growth factor β 2 via p38 mitogen-activated protein kinase. <i>FASEB Journal</i> , 2001, 15, 1098-1100.	0.5	59

#	ARTICLE	IF	CITATIONS
73	Tumor cell-specific AIM2 regulates growth and invasion of cutaneous squamous cell carcinoma. <i>Oncotarget</i> , 2017, 8, 45825-45836.	1.8	59
74	Expression of collagenase-3 (matrix metalloproteinase-13) in transitional-cell carcinoma of the urinary bladder. <i>International Journal of Cancer</i> , 2000, 88, 417-423.	5.1	58
75	Matrix metalloproteinase-19 is expressed by proliferating epithelium but disappears with neoplastic dedifferentiation. <i>International Journal of Cancer</i> , 2003, 103, 709-716.	5.1	58
76	Scleroderma-like cutaneous syndromes. <i>Current Rheumatology Reports</i> , 2002, 4, 113-122.	4.7	57
77	Transformation-specific matrix metalloproteinases (MMP)-7 and MMP-13 are expressed by tumour cells in epidermolysis bullosa-associated squamous cell carcinomas. <i>British Journal of Dermatology</i> , 2008, 158, 778-785.	1.5	57
78	Elevated $\alpha 2(I)$ collagen mRNA levels in cultured scleroderma fibroblasts result from an increased transcription rate of the corresponding gene. <i>FEBS Letters</i> , 1987, 215, 331-334.	2.8	56
79	Activation of Extracellular Signal-regulated Kinase 1/2 Inhibits Type I Collagen Expression by Human Skin Fibroblasts. <i>Journal of Biological Chemistry</i> , 2000, 275, 34634-34639.	3.4	55
80	Oncolytic Capacity of Attenuated Replicative Semliki Forest Virus in Human Melanoma Xenografts in Severe Combined Immunodeficient Mice. <i>Cancer Research</i> , 2006, 66, 7185-7194.	0.9	55
81	TGF- $\beta 2$ -Elicited Induction of Tissue Inhibitor of Metalloproteinases (TIMP)-3 Expression in Fibroblasts Involves Complex Interplay between Smad3, p38 β , and ERK1/2. <i>PLoS ONE</i> , 2013, 8, e57474.	2.5	55
82	Activation of Dermal Connective Tissue in Scleroderma. <i>Annals of Medicine</i> , 1993, 25, 511-518.	3.8	54
83	Inhibition of collagenase-3 (MMP-13) expression in transformed human keratinocytes by interferon- $\beta 3$ is associated with activation of extracellular signal-regulated kinase-1,2 and STAT1. <i>Oncogene</i> , 2000, 19, 248-257.	5.9	54
84	Matrix Metalloproteinase-13 Promotes Recovery from Experimental Liver Cirrhosis in Rats. <i>Pathobiology</i> , 2011, 78, 239-252.	3.8	54
85	Human Granulation-tissue Fibroblasts Show Enhanced Proteoglycan Gene Expression and Altered Response to TGF- $\beta 2$. <i>Journal of Dental Research</i> , 1996, 75, 1767-1778.	5.2	53
86	Transforming growth factor- β -induced alpha-smooth muscle cell actin expression in renal proximal tubular cells is regulated by p38 β mitogen-activated protein kinase, extracellular signal-regulated protein kinase1,2 and the Smad signalling during epithelial-myofibroblast transdifferentiation. <i>Nephrology Dialysis Transplantation</i> , 2008, 23, 1537-1545.	0.7	52
87	Collagenase-3 (MMP-13) Enhances Remodeling of Three-Dimensional Collagen and Promotes Survival of Human Skin Fibroblasts. <i>Journal of Investigative Dermatology</i> , 2007, 127, 49-59.	0.7	51
88	Matrix Metalloproteinase-19 Expression in Dermal Wounds and by Fibroblasts in Culture. <i>Journal of Investigative Dermatology</i> , 2003, 121, 997-1004.	0.7	50
89	Hypoxia-activated Smad3-specific Dephosphorylation by PP2A. <i>Journal of Biological Chemistry</i> , 2010, 285, 3740-3749.	3.4	49
90	Human TIMP-3 Is Expressed During Fetal Development, Hair Growth Cycle, and Cancer Progression. <i>Journal of Histochemistry and Cytochemistry</i> , 1998, 46, 437-447.	2.5	48

#	ARTICLE	IF	CITATIONS
91	Association between high collagenase-3 expression levels and poor prognosis in patients with head and neck cancer. <i>Head and Neck</i> , 2006, 28, 225-234.	2.0	48
92	EphB2 Promotes Progression of Cutaneous Squamous Cell Carcinoma. <i>Journal of Investigative Dermatology</i> , 2015, 135, 1882-1892.	0.7	48
93	Î±V integrin promotes in vitro and in vivo survival of cells in metastatic melanoma. <i>International Journal of Cancer</i> , 2004, 112, 61-70.	5.1	47
94	Human recombinant interleukin-1 regulates cellular mRNA levels of dermatan sulphate proteoglycan core protein. <i>Biochemical Journal</i> , 1988, 252, 309-312.	3.7	45
95	Expression and activity of matrix metalloproteinase-2 and -9 in experimental granulation tissue. <i>Apms</i> , 2000, 108, 318-328.	2.0	45
96	Expression of collagenase-3 (MMP-13) enhances invasion of human fibrosarcoma HT-1080 cells. <i>International Journal of Cancer</i> , 2002, 97, 283-289.	5.1	44
97	Differential Regulation of Decorin and Biglycan Gene Expression by Dexamethasone and Retinoic Acid in Cultured Human Skin Fibroblasts. <i>Journal of Investigative Dermatology</i> , 1995, 104, 503-508.	0.7	43
98	Regulation of Elastin Gene Expression: Evidence for Functional Promoter Activity in the 5'-Flanking Region of the Human Gene. <i>Journal of Investigative Dermatology</i> , 1990, 94, 191-196.	0.7	42
99	Regulation of Elastin Gene Expression. <i>Annals of the New York Academy of Sciences</i> , 1991, 624, 116-136.	3.8	42
100	Molecular biology and pathology of human elastin. <i>Biochemical Society Transactions</i> , 1991, 19, 824-829.	3.4	40
101	Tumour cell-derived complement components C1r and C1s promote growth of cutaneous squamous cell carcinoma. <i>British Journal of Dermatology</i> , 2020, 182, 658-670.	1.5	40
102	Characterization of One Phenotype of Human Periodontal Granulation-tissue Fibroblasts. <i>Journal of Dental Research</i> , 1989, 68, 20-25.	5.2	39
103	MicroRNA-203 Inversely Correlates with Differentiation Grade, Targets c-MYC, and Functions as a Tumor Suppressor in cSCC. <i>Journal of Investigative Dermatology</i> , 2016, 136, 2485-2494.	0.7	39
104	Differential Regulation of the AP-1 Family Members by UV Irradiation In Vitro and In Vivo. <i>Cellular Signalling</i> , 1998, 10, 191-195.	3.6	38
105	Squamous cell carcinoma of the skin: Emerging need for novel biomarkers. <i>World Journal of Clinical Oncology</i> , 2013, 4, 85.	2.3	37
106	Collagen synthesis in the vaginal connective tissue of patients with and without uterine prolapse. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 1987, 24, 319-325.	1.1	33
107	Efficient infection of tumor endothelial cells by a capsid-modified adenovirus. <i>Gene Therapy</i> , 2006, 13, 52-59.	4.5	33
108	Matrix metalloproteinase (MMP)-1, -9 and -13 as prognostic factors in salivary gland cancer. <i>Acta Oto-Laryngologica</i> , 2008, 128, 482-490.	0.9	33

#	ARTICLE	IF	CITATIONS
109	CCHCR1 Is Up-Regulated in Skin Cancer and Associated with EGFR Expression. PLoS ONE, 2009, 4, e6030.	2.5	33
110	Protodynamic Intracellular Acidification by cis-Urocanic Acid Promotes Apoptosis of Melanoma Cells In Vitro and In Vivo. Journal of Investigative Dermatology, 2010, 130, 2431-2439.	0.7	33
111	p53-Regulated Long Noncoding RNA PRECISIT Promotes Progression of Cutaneous Squamous Cell Carcinoma via STAT3 Signaling. American Journal of Pathology, 2020, 190, 503-517.	3.8	33
112	Collagenase-1, stromelysin-1 and 92 kDa gelatinase are associated with tumor necrosis factor- α induced morphological change of human endothelial cells in Vitro. Matrix Biology, 1998, 17, 293-304.	3.6	32
113	Hypoxic Conversion of SMAD7 Function from an Inhibitor into a Promoter of Cell Invasion. Cancer Research, 2010, 70, 5984-5993.	0.9	32
114	Significant Role of Collagen XVII And Integrin α 24 in Migration and Invasion of The Less Aggressive Squamous Cell Carcinoma Cells. Scientific Reports, 2017, 7, 45057.	3.3	32
115	Serum VEGF-C is associated with metastatic site in patients with malignant melanoma. Acta Oncologica, 2007, 46, 678-684.	1.8	31
116	TIMP-3 promotes apoptosis in nonadherent small cell lung carcinoma cells lacking functional death receptor pathway. International Journal of Cancer, 2011, 128, 991-996.	5.1	31
117	Long non-coding RNA PICSAR decreases adhesion and promotes migration of squamous carcinoma cells by downregulating α 21 and α 51 integrin expression. Biology Open, 2018, 7, .	1.2	31
118	Matrix metalloproteinase (MMP)-7 in salivary gland cancer. Acta Oncologica, 2010, 49, 85-90.	1.8	30
119	Collagens XV and XVIII show different expression and localisation in cutaneous squamous cell carcinoma: type XV appears in tumor stroma, while XVIII becomes upregulated in tumor cells and lost from microvessels. Experimental Dermatology, 2016, 25, 348-354.	2.9	30
120	Epidermal growth factor (EGF) prevents methylprednisolone-induced inhibition of wound healing. Journal of Surgical Research, 1989, 47, 354-359.	1.6	28
121	The Role of p53 in Progression of Cutaneous Squamous Cell Carcinoma. Cancers, 2021, 13, 4507.	3.7	28
122	Fibroblast Activation in Scleroderma. Scandinavian Journal of Rheumatology, 1984, 13, 229-237.	1.1	27
123	Expression of matrix metalloproteinases and tissue inhibitors of metalloproteinases in human chondrosarcomas. Note. Apms, 2001, 109, 305-315.	2.0	27
124	Potential Applications of Tissue Inhibitor of Metalloproteinase (TIMP) Overexpression For Cancer Gene Therapy. Advances in Experimental Medicine and Biology, 2002, 465, 469-483.	1.6	26
125	Collagen Turnover in Wound Repair – A Macrophage Connection. Journal of Investigative Dermatology, 2015, 135, 2350-2352.	0.7	26
126	Complement System in Cutaneous Squamous Cell Carcinoma. International Journal of Molecular Sciences, 2019, 20, 3550.	4.1	26

#	ARTICLE	IF	CITATIONS
127	Induction of periosteal callus formation by bone morphogenetic protein-2 employing adenovirus-mediated gene delivery. <i>Matrix Biology</i> , 2001, 20, 123-127.	3.6	25
128	Dexamethasone Suppresses Elastin Gene Expression in Human Skin Fibroblasts in Culture. <i>Biochemical and Biophysical Research Communications</i> , 1994, 201, 1189-1196.	2.1	24
129	High collagenase-1 expression correlates with a favourable chemoimmunotherapy response in human metastatic melanoma. <i>Melanoma Research</i> , 2001, 11, 157-166.	1.2	24
130	Keratinocyte Growth Factor Induces Gene Expression Signature Associated with Suppression of Malignant Phenotype of Cutaneous Squamous Carcinoma Cells. <i>PLoS ONE</i> , 2012, 7, e33041.	2.5	24
131	Cyclosporin A Enhances Cytokine and Phorbol Ester-Induced Fibroblast Collagenase Expression. <i>Journal of Investigative Dermatology</i> , 1994, 102, 938-944.	0.7	23
132	TNF-R55-Specific Form of Human Tumor Necrosis Factor- β Induces Collagenase Gene Expression By Human Skin Fibroblasts. <i>Journal of Investigative Dermatology</i> , 1995, 105, 197-202.	0.7	23
133	High-efficiency gene transfer to primary T lymphocytes by recombinant adenovirus vectors. <i>Journal of Immunological Methods</i> , 2002, 260, 79-89.	1.4	23
134	H-Ras activation and fibroblast-induced TGF- β signaling promote laminin-332 accumulation and invasion in cutaneous squamous cell carcinoma. <i>Matrix Biology</i> , 2020, 87, 26-47.	3.6	23
135	Matrix metalloproteinases in keratinocyte carcinomas. <i>Experimental Dermatology</i> , 2021, 30, 50-61.	2.9	23
136	Risk Factors and Prognosis for Metastatic Cutaneous Squamous Cell Carcinoma: A Cohort Study. <i>Acta Dermato-Venereologica</i> , 2020, 100, adv00266.	1.3	23
137	Adenovirus mediated intra-articular expression of collagenase-3 (MMP-13) induces inflammatory arthritis in mice. <i>Annals of the Rheumatic Diseases</i> , 2004, 63, 656-664.	0.9	22
138	Transcription of β 2 Integrin Gene in Osteosarcoma Cells Is Enhanced by Tumor Promoters. <i>Experimental Cell Research</i> , 1998, 243, 1-10.	2.6	20
139	Expression of matrix metalloproteinase-1, -7, -9, -13, Ki-67, and HER-2 in epithelial-myoeipithelial salivary gland cancer. <i>Head and Neck</i> , 2010, 32, 1019-1027.	2.0	20
140	Dasatinib promotes apoptosis of cutaneous squamous carcinoma cells by regulating activation of ERK1/2. <i>Experimental Dermatology</i> , 2017, 26, 89-92.	2.9	20
141	C1r Upregulates Production of Matrix Metalloproteinase-13 and Promotes Invasion of Cutaneous Squamous Cell Carcinoma. <i>Journal of Investigative Dermatology</i> , 2022, 142, 1478-1488.e9.	0.7	19
142	Activation of extracellular signal-regulated protein kinase1,2 results in down-regulation of decorin expression in fibroblasts. <i>Biochemical Journal</i> , 2000, 349, 19-25.	3.7	18
143	p38 β mitogen-activated protein kinase regulates the expression of tight junction protein ZO-1 in differentiating human epidermal keratinocytes. <i>Archives of Dermatological Research</i> , 2014, 306, 131-141.	1.9	18
144	Gene expression of fibroblast matrix proteins is altered by indomethacin. <i>FEBS Letters</i> , 1988, 231, 125-129.	2.8	17

#	ARTICLE	IF	CITATIONS
145	Extended release of adenovirus from silica implants in vitro and in vivo. <i>Gene Therapy</i> , 2009, 16, 103-110.	4.5	17
146	Eosinophilic fasciitis. Increased collagen production and type I procollagen messenger RNA levels in fibroblasts cultured from involved skin. <i>Archives of Dermatology</i> , 1990, 126, 613-617.	1.4	17
147	Leukoregulin, A T-cell derived cytokine, upregulates stromelysin-1 gene expression in human dermal fibroblasts: Evidence for the role of AP-1 in transcriptional activation. <i>Journal of Cellular Biochemistry</i> , 1992, 50, 53-61.	2.6	16
148	Transcriptional targeting of adenoviral gene delivery into migrating wound keratinocytes using FiRE, a growth factor-inducible regulatory element. <i>Gene Therapy</i> , 2000, 7, 1640-1647.	4.5	15
149	Human Nidogen Gene: Structural and Functional Characterization of the 5'-Flanking Region. <i>Journal of Investigative Dermatology</i> , 1991, 97, 281-285.	0.7	14
150	New prognostic factors and developing therapy of cutaneous melanoma. <i>Annals of Medicine</i> , 2003, 35, 66-78.	3.8	14
151	Activation of extracellular signal-regulated protein kinase1,2 results in down-regulation of decorin expression in fibroblasts. <i>Biochemical Journal</i> , 2000, 349, 19.	3.7	13
152	Expression of extracellular matrix genes: transforming growth factor (TGF)- β 21 and ras in tibial fracture healing of lathyrictic rats. <i>Bone</i> , 2000, 27, 551-557.	2.9	13
153	Temporospatial expression of matrix metalloproteinases and tissue inhibitors of matrix metalloproteinases in mouse antigen-induced arthritis. <i>Histochemistry and Cell Biology</i> , 2005, 124, 535-545.	1.7	12
154	Expression of claudin-11 by tumor cells in cutaneous squamous cell carcinoma is dependent on the activity of p38 β . <i>Experimental Dermatology</i> , 2017, 26, 771-777.	2.9	12
155	Long non-coding RNAs in cutaneous biology and keratinocyte carcinomas. <i>Cellular and Molecular Life Sciences</i> , 2020, 77, 4601-4614.	5.4	12
156	Discovery of a Novel CIP2A Variant (NOCIVA) with Clinical Relevance in Predicting TKI Resistance in Myeloid Leukemias. <i>Clinical Cancer Research</i> , 2021, 27, 2848-2860.	7.0	11
157	Signaling pathways in human osteoclasts differentiation: ERK1/2 as a key player. <i>Molecular Biology Reports</i> , 2021, 48, 1243-1254.	2.3	11
158	Identification of metastatic primary cutaneous squamous cell carcinoma utilizing artificial intelligence analysis of whole slide images. <i>Scientific Reports</i> , 2022, 12, .	3.3	11
159	Inhibition of c-Abl Kinase Activity Renders Cancer Cells Highly Sensitive to Mitoxantrone. <i>PLoS ONE</i> , 2014, 9, e105526.	2.5	10
160	High Level Expression of Tissue Inhibitors of Metalloproteinases-1,-2 and -3 in Melanoma Cells Achieved by Adenovirus Mediated Gene Transfer. <i>Advances in Experimental Medicine and Biology</i> , 1998, 451, 69-72.	1.6	9
161	ADAMTS5. <i>American Journal of Pathology</i> , 2012, 181, 743-745.	3.8	8
162	Complement factor I upregulates expression of matrix metalloproteinase-13 and -2 and promotes invasion of cutaneous squamous carcinoma cells. <i>Experimental Dermatology</i> , 2021, 30, 1631-1641.	2.9	8

#	ARTICLE	IF	CITATIONS
163	Expression of Collagenase-3 (MMP-13) by Tumor Cells in Squamous Cell Carcinomas of the Head and Neck. <i>Advances in Experimental Medicine and Biology</i> , 1998, 451, 63-68.	1.6	8
164	Complement Factor D Is a Novel Biomarker and Putative Therapeutic Target in Cutaneous Squamous Cell Carcinoma. <i>Cancers</i> , 2022, 14, 305.	3.7	8
165	Collagen in the Extracellular Matrix of Cultured Scleroderma Skin Fibroblasts: Changes Related to Ascorbic Acid-Treatment. <i>Matrix Biology</i> , 1989, 9, 34-39.	1.7	7
166	Natural killer cells in wound healing. , 2010, , 519-525.		7
167	Loss of the laminin subunit alpha β 3 induces cell invasion and macrophage infiltration in cutaneous squamous cell carcinoma*. <i>British Journal of Dermatology</i> , 2021, 184, 923-934.	1.5	7
168	The Viability and Growth of HaCaT Cells After Exposure to Bioactive Glass S53P4-Containing Cell Culture Media. <i>Otology and Neurotology</i> , 2021, 42, e559-e567.	1.3	7
169	Expression of human collagenase β 3 (MMP β 13) by fetal skin fibroblasts is induced by transforming growth factor β 2 via p38 mitogen β activated protein kinase. <i>FASEB Journal</i> , 2001, 15, 1098-1100.	0.5	6
170	The protein phosphatase inhibitor okadaic acid suppresses type I collagen gene expression in cultured fibroblasts at the transcriptional level. <i>Biochemical Journal</i> , 1995, 308, 995-999.	3.7	5
171	Suppression of Elastin Gene Expression in Dermal Fibroblasts by Protein Phosphatase Inhibitor Okadaic Acid. <i>Biochemical and Biophysical Research Communications</i> , 1995, 209, 175-181.	2.1	4
172	Clinical and Pathological Aspects of Melanoma among Children in Finland. <i>Acta Dermato-Venereologica</i> , 2016, 96, 718-720.	1.3	4
173	Increased incidence of melanoma in children and adolescents in Finland in 1990 β 2014: nationwide re-evaluation of histopathological characteristics. <i>Annals of Medicine</i> , 2022, 54, 244-252.	3.8	4
174	Diagnostic and prognostic role of matrix metalloproteases in cancer. <i>Expert Opinion on Medical Diagnostics</i> , 2008, 2, 1025-1039.	1.6	3
175	Different expression of BRAFV600E, ALK and PD-L1 in melanoma in children and adolescents: a nationwide retrospective study in Finland in 1990 β 2014. <i>Acta Oncolβgica</i> , 2021, 60, 165-172.	1.8	3
176	Flow Cytometry of Fibroblasts Cultured from Skin of Patients with Localized Scleroderma. <i>Dermatology</i> , 1988, 177, 348-353.	2.1	1
177	Introduction: Cell invasion: cooperation between gene families at distinct levels. <i>Cellular and Molecular Life Sciences</i> , 2000, 57, 3-4.	5.4	1
178	Stromal Collagenase in Melanoma: A Vascular Connection. <i>Journal of Investigative Dermatology</i> , 2009, 129, 2545-2547.	0.7	1
179	Tumor cell-specific Serpin A1 expression in vulvar squamous cell carcinoma. <i>Archives of Gynecology and Obstetrics</i> , 2019, 299, 1345-1351.	1.7	1
180	Abstract 1098: MiR-203 suppresses cutaneous squamous cell carcinoma growth and targets the myc oncogene. , 2016, , .		1

#	ARTICLE	IF	CITATIONS
181	Forty Years of the European Society for Dermatological Research as European Dermatology Goes from Strength to Strength. <i>Journal of Investigative Dermatology</i> , 2010, 130, 1957-1959.	0.7	0
182	Ulpu Saarialho-Kere (1960–2009). <i>Journal of Investigative Dermatology</i> , 2010, 130, 640.	0.7	0
183	Abstract 1074: Keratinocyte growth factor suppresses malignant phenotype of cutaneous squamous carcinoma cells. , 2012, , .		0
184	Abstract 3201: Complement component C3 and complement factor B regulate growth of cutaneous squamous cell carcinoma. , 2015, , .		0
185	Targeting Degradome Genes via Engineered Viral Vectors. , 0, , 877-894.		0