

Nicolle H R Litjens

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8380039/publications.pdf>

Version: 2024-02-01

68
papers

2,416
citations

186265
28
h-index

214800
47
g-index

70
all docs

70
docs citations

70
times ranked

3041
citing authors

#	ARTICLE	IF	CITATIONS
1	Activated CD4+ T Cells and Highly Differentiated Alloreactive CD4+ T Cells Distinguish Operationally Tolerant Liver Transplantation Recipients. <i>Liver Transplantation</i> , 2022, 28, 98-112.	2.4	8
2	ARHGDI1 and AT1R autoantibodies are differentially related to the development and presence of chronic antibody-mediated rejection and fibrosis in kidney allografts. <i>Human Immunology</i> , 2021, 82, 89-96.	2.4	10
3	Expression of Senescence Marker TIGIT Identifies Polyfunctional Donor-Reactive CD4+ T Cells Preferentially Lost After Kidney Transplantation. <i>Frontiers in Immunology</i> , 2021, 12, 656846.	4.8	15
4	The FCGR3A 158V/V-genotype is associated with decreased survival of renal allografts with chronic active antibody-mediated rejection. <i>Scientific Reports</i> , 2021, 11, 7903.	3.3	12
5	Effects of Morbid Obesity and Metabolic Syndrome on the Composition of Circulating Immune Subsets. <i>Frontiers in Immunology</i> , 2021, 12, 675018.	4.8	5
6	Alemtuzumab as Second-Line Treatment for Late Antibody-Mediated Rejection of Transplanted Kidneys. <i>Transplantation Proceedings</i> , 2021, 53, 2206-2211.	0.6	8
7	Current Tolerance-Associated Peripheral Blood Gene Expression Profiles After Liver Transplantation Are Influenced by Immunosuppressive Drugs and Prior Cytomegalovirus Infection. <i>Frontiers in Immunology</i> , 2021, 12, 738837.	4.8	1
8	Validation of a Combined Transcriptome and T Cell Receptor Alpha/Beta (TRA/TRB) Repertoire Assay at the Single Cell Level for Paucicellular Samples. <i>Frontiers in Immunology</i> , 2020, 11, 1999.	4.8	3
9	High numbers of differentiated CD28null CD8+ T cells are associated with a lowered risk for late rejection and graft loss after kidney transplantation. <i>PLoS ONE</i> , 2020, 15, e0228096.	2.5	12
10	A very low thymus function identifies patients with substantial increased risk for long-term mortality after kidney transplantation. <i>Immunity and Ageing</i> , 2020, 17, 4.	4.2	15
11	Title is missing!. , 2020, 15, e0228096.		0
12	Title is missing!. , 2020, 15, e0228096.		0
13	Title is missing!. , 2020, 15, e0228096.		0
14	Title is missing!. , 2020, 15, e0228096.		0
15	Title is missing!. , 2020, 15, e0228096.		0
16	Title is missing!. , 2020, 15, e0228096.		0
17	Increased CD16 expression on NK cells is indicative of antibody-dependent cell-mediated cytotoxicity in chronic-active antibody-mediated rejection. <i>Transplant Immunology</i> , 2019, 54, 52-58.	1.2	22
18	Immunosuppressive drug withdrawal late after liver transplantation improves the lipid profile and reduces infections. <i>European Journal of Gastroenterology and Hepatology</i> , 2019, 31, 1444-1451.	1.6	5

#	ARTICLE	IF	CITATIONS
19	Co-inhibitory profile and cytotoxicity of CD57+PD-1 ^{hi} T cells in end-stage renal disease patients. <i>Clinical and Experimental Immunology</i> , 2018, 191, 363-372.	2.6	6
20	CD4+CD28null T cells are not alloreactive unless stimulated by interleukin-15. <i>American Journal of Transplantation</i> , 2018, 18, 341-350.	4.7	10
21	A comprehensive characterization of aggravated aging-related changes in T lymphocytes and monocytes in end-stage renal disease: the iESRD study. <i>Immunity and Ageing</i> , 2018, 15, 27.	4.2	43
22	Potential Beneficial Effects of Cytomegalovirus Infection after Transplantation. <i>Frontiers in Immunology</i> , 2018, 9, 389.	4.8	49
23	Lymph node and circulating T cell characteristics are strongly correlated in end-stage renal disease patients, but highly differentiated T cells reside within the circulation. <i>Clinical and Experimental Immunology</i> , 2017, 188, 299-310.	2.6	15
24	Pretransplant Numbers of CD16 + Monocytes as a Novel Biomarker to Predict Acute Rejection After Kidney Transplantation: A Pilot Study. <i>American Journal of Transplantation</i> , 2017, 17, 2659-2667.	4.7	29
25	Natural regulatory T cells from patients with end-stage renal disease can be used for large-scale generation of highly suppressive alloantigen-specific Tregs. <i>Kidney International</i> , 2017, 91, 1203-1213.	5.2	10
26	pERK-dependent defective TCR-mediated activation of CD4+ T cells in end-stage renal disease patients. <i>Immunity and Ageing</i> , 2017, 14, 14.	4.2	12
27	Belatacept Does Not Inhibit Follicular T Cell-Dependent B-Cell Differentiation in Kidney Transplantation. <i>Frontiers in Immunology</i> , 2017, 8, 641.	4.8	25
28	Protective Cytomegalovirus (CMV)-Specific T-Cell Immunity Is Frequent in Kidney Transplant Patients without Serum Anti-CMV Antibodies. <i>Frontiers in Immunology</i> , 2017, 8, 1137.	4.8	22
29	T-Cell Composition of the Lymph Node Is Associated with the Risk for Early Rejection after Renal Transplantation. <i>Frontiers in Immunology</i> , 2017, 8, 1416.	4.8	9
30	End-Stage Renal Disease Causes Skewing in the TCR V β 2-Repertoire Primarily within CD8+ T Cell Subsets. <i>Frontiers in Immunology</i> , 2017, 8, 1826.	4.8	19
31	Latency for cytomegalovirus impacts T cell ageing significantly in elderly end-stage renal disease patients. <i>Clinical and Experimental Immunology</i> , 2016, 186, 239-248.	2.6	13
32	Uremia-Associated Premature Aging of T Cells Does Not Predict Infectious Complications After Renal Transplantation. <i>American Journal of Transplantation</i> , 2016, 16, 2324-2333.	4.7	17
33	Loss of CD28 on Peripheral T Cells Decreases the Risk for Early Acute Rejection after Kidney Transplantation. <i>PLoS ONE</i> , 2016, 11, e0150826.	2.5	46
34	End stage renal disease patients have a skewed T cell receptor V β 2 repertoire. <i>Immunity and Ageing</i> , 2015, 12, 28.	4.2	20
35	Primary Cytomegalovirus Infection Significantly Impacts Circulating T Cells in Kidney Transplant Recipients. <i>American Journal of Transplantation</i> , 2015, 15, 3143-3156.	4.7	28
36	Encapsulating peritoneal sclerosis is associated with T-cell activation. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, 1568-1576.	0.7	13

#	ARTICLE	IF	CITATIONS
37	Chronic Kidney Disease and Premature Ageing of the Adaptive Immune Response. <i>Current Urology Reports</i> , 2015, 16, 471.	2.2	48
38	Allogeneic Mature Human Dendritic Cells Generate Superior Alloreactive Regulatory T Cells in the Presence of IL-15. <i>Journal of Immunology</i> , 2015, 194, 5282-5293.	0.8	12
39	Follicular T helper cells and humoral reactivity in kidney transplant patients. <i>Clinical and Experimental Immunology</i> , 2015, 180, 329-340.	2.6	107
40	Rotterdam: Main port for organ transplantation research in the Netherlands. <i>Transplant Immunology</i> , 2014, 31, 200-206.	1.2	1
41	Uremia-associated immunological aging is stably imprinted in the T-cell system and not reversed by kidney transplantation. <i>Transplant International</i> , 2014, 27, 1272-1284.	1.6	55
42	Substantial Proliferation of Human Renal Tubular Epithelial Cell-Reacting CD4+CD28null Memory T Cells, Which Is Resistant to Tacrolimus and Everolimus. <i>Transplantation</i> , 2014, 97, 47-55.	1.0	17
43	T-cell ageing in end-stage renal disease patients: Assessment and clinical relevance. <i>World Journal of Nephrology</i> , 2014, 3, 268.	2.0	30
44	Mesenchymal stem cells control alloreactive CD8+CD28 ⁻ T cells. <i>Clinical and Experimental Immunology</i> , 2013, 174, 449-458.	2.6	41
45	Cytomegalovirus contributes partly to uraemia-associated premature immunological ageing of the T cell compartment. <i>Clinical and Experimental Immunology</i> , 2013, 174, 424-432.	2.6	36
46	Activation-induced CD137 is a fast assay for identification and multi-parameter flow cytometric analysis of alloreactive T cells. <i>Clinical and Experimental Immunology</i> , 2013, 174, 179-191.	2.6	48
47	Human adipose-tissue derived mesenchymal stem cells induce functional <i>de-novo</i> regulatory T cells with methylated FOXP3 gene DNA. <i>Clinical and Experimental Immunology</i> , 2013, 173, 343-354.	2.6	79
48	Loss of Renal Function Causes Premature Aging of the Immune System. <i>Blood Purification</i> , 2013, 36, 173-178.	1.8	61
49	Kinetics of Homeostatic Proliferation and Thymopoiesis after rATG Induction Therapy in Kidney Transplant Patients. <i>Transplantation</i> , 2013, 96, 904-913.	1.0	36
50	Circulating CD4+CD28null T Cells May Increase the Risk of an Atherosclerotic Vascular Event Shortly after Kidney Transplantation. <i>Journal of Transplantation</i> , 2013, 2013, 1-8.	0.5	15
51	Identification of Circulating Human Antigen-Reactive CD4+FOXP3+ Natural Regulatory T Cells. <i>Journal of Immunology</i> , 2012, 188, 1083-1090.	0.8	32
52	Terminally Differentiated CD8+ Temra Cells Are Associated With the Risk for Acute Kidney Allograft Rejection. <i>Transplantation</i> , 2012, 94, 63-69.	1.0	75
53	Systemic varicella zoster virus reactive effector memory T cells impaired in the elderly and in kidney transplant recipients. <i>Journal of Medical Virology</i> , 2012, 84, 2018-2025.	5.0	26
54	A killer on the road: circulating CD4+CD28null T cells as cardiovascular risk factor in ESRD patients. <i>Journal of Nephrology</i> , 2012, 25, 183-191.	2.0	40

#	ARTICLE	IF	CITATIONS
55	Premature aging of circulating T cells in patients with end-stage renal disease. <i>Kidney International</i> , 2011, 80, 208-217.	5.2	181
56	Circulating pro-inflammatory CD4 ^{pos} CD28 ^{null} T cells are independently associated with cardiovascular disease in ESRD patients. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 3640-3646.	0.7	55
57	CMV Seropositivity Determines Epoetin Dose and Hemoglobin Levels in Patients with CKD. <i>Journal of the American Society of Nephrology: JASN</i> , 2009, 20, 2661-2666.	6.1	26
58	The human alloreactive CD4 ⁺ T-cell repertoire is biased to a Th17 response and the frequency is inversely related to the number of HLA class II mismatches. <i>Blood</i> , 2009, 114, 3947-3955.	1.4	33
59	Hepatitis B vaccine-specific CD4 ⁺ T cells can be detected and characterised at the single cell level: Limited usefulness of dendritic cells as signal enhancers. <i>Journal of Immunological Methods</i> , 2008, 330, 1-11.	1.4	31
60	IL-2 Producing Memory CD4 ⁺ T Lymphocytes Are Closely Associated with the Generation of IgG-Secreting Plasma Cells. <i>Journal of Immunology</i> , 2008, 181, 3665-3673.	0.8	50
61	Expansion of cytolytic CD4 ⁺ CD28 [~] T cells in end-stage renal disease. <i>Kidney International</i> , 2008, 74, 760-767.	5.2	95
62	Seropositivity for cytomegalovirus in patients with end-stage renal disease is strongly associated with atherosclerotic disease. <i>Nephrology Dialysis Transplantation</i> , 2007, 22, 3298-3303.	0.7	65
63	Progressive loss of renal function is associated with activation and depletion of naive T lymphocytes. <i>Clinical Immunology</i> , 2006, 118, 83-91.	3.2	126
64	Pharmacokinetics of oral fumarates in healthy subjects. <i>British Journal of Clinical Pharmacology</i> , 2004, 58, 429-432.	2.4	124
65	Psoriasis Is Not Associated with IL-12p70/IL-12p40 Production and IL12B Promoter Polymorphism. <i>Journal of Investigative Dermatology</i> , 2004, 122, 923-926.	0.7	22
66	In vitro pharmacokinetics of anti-psoriatic fumaric acid esters. <i>BMC Pharmacology</i> , 2004, 4, 22.	0.4	62
67	Monomethylfumarate affects polarization of monocyte-derived dendritic cells resulting in down-regulated Th1 lymphocyte responses. <i>European Journal of Immunology</i> , 2004, 34, 565-575.	2.9	99
68	Time Course of Atrial Fibrillation-induced Cellular Structural Remodeling in Atria of the Goat. <i>Journal of Molecular and Cellular Cardiology</i> , 2001, 33, 2083-2094.	1.9	186