

Dani BergÃ©

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8399161/publications.pdf>

Version: 2024-02-01

48
papers

1,038
citations

471509

17
h-index

434195

31
g-index

60
all docs

60
docs citations

60
times ranked

1929
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Improvement in prefrontal thalamic connectivity during the early course of the illness in recent-onset psychosis: a 12-month longitudinal follow-up resting-state fMRI study. <i>Psychological Medicine</i> , 2022, 52, 2713-2721. | 4.5 | 10 |
| 2 | Multidimensional predictors of negative symptoms in antipsychotic-naive first-episode psychosis. <i>Journal of Psychiatry and Neuroscience</i> , 2022, 47, E21-E31. | 2.4 | 5 |
| 3 | Brain activity and connectivity differences in reward value discrimination during effort computation in schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 647-659. | 3.2 | 3 |
| 4 | Dificultades en el parto y sintomatologÃa depresiva en la esquizofrenia. <i>Revista De PsiquiatrÃa Y Salud Mental</i> , 2021, 14, 66-68. | 1.8 | 5 |
| 5 | The prevention of relapses in first episodes of schizophrenia: The 2EPs Project, background, rationale and study design. <i>Revista De PsiquiatrÃa Y Salud Mental (English Edition)</i> , 2021, 14, 164-176. | 0.3 | 3 |
| 6 | The prevention of relapses in first episodes of schizophrenia: The 2EPs Project, background, rationale and study design. <i>Revista De PsiquiatrÃa Y Salud Mental</i> , 2021, 14, 164-176. | 1.8 | 13 |
| 7 | The effect of family environment and psychiatric family history on psychosocial functioning in first-episode psychosis at baseline and after 2 years. <i>European Neuropsychopharmacology</i> , 2021, 49, 54-68. | 0.7 | 12 |
| 8 | Cognitive clusters in first-episode psychosis. <i>Schizophrenia Research</i> , 2021, 237, 31-39. | 2.0 | 10 |
| 9 | Exploring Risk and Resilient Profiles for Functional Impairment and Baseline Predictors in a 2-Year Follow-Up First-Episode Psychosis Cohort Using Latent Class Growth Analysis. <i>Journal of Clinical Medicine</i> , 2021, 10, 73. | 2.4 | 14 |
| 10 | Influence of BDNF and MTHFR polymorphisms on hippocampal volume in first-episode psychosis. <i>Schizophrenia Research</i> , 2020, 223, 345-352. | 2.0 | 8 |
| 11 | Family Features of Social Withdrawal Syndrome (Hikikomori). <i>Frontiers in Psychiatry</i> , 2020, 11, 138. | 2.6 | 25 |
| 12 | Birth weight and antipsychotic induced weight gain: A prenatal programming approach in the PEPs study. <i>Schizophrenia Research</i> , 2020, 218, 292-294. | 2.0 | 6 |
| 13 | The impact of sex and cannabis on clinical features in first-admitted patients with psychosis. <i>European Neuropsychopharmacology</i> , 2020, 36, 235-243. | 0.7 | 15 |
| 14 | Influence of secondary sources in the Brief Negative Symptom Scale. <i>Schizophrenia Research</i> , 2019, 204, 452-454. | 2.0 | 5 |
| 15 | Smoking does not impact social and non-social cognition in patients with first episode psychosis. <i>Schizophrenia Research</i> , 2018, 199, 64-74. | 2.0 | 13 |
| 16 | A 12-month study of the hikikomori syndrome of social withdrawal: Clinical characterization and different subtypes proposal. <i>Psychiatry Research</i> , 2018, 270, 1039-1046. | 3.3 | 42 |
| 17 | Reduced willingness to invest effort in schizophrenia with high negative symptoms regardless of reward stimulus presentation and reward value. <i>Comprehensive Psychiatry</i> , 2018, 87, 153-160. | 3.1 | 8 |
| 18 | Obstetric Phenotypes in the Heterogeneity of Schizophrenia. <i>Journal of Nervous and Mental Disease</i> , 2018, 206, 882-886. | 1.0 | 18 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Affective versus non-affective first episode psychoses: A longitudinal study. <i>Journal of Affective Disorders</i> , 2018, 238, 297-304. | 4.1 | 26 |
| 20 | Cannabis use, COMT, BDNF and age at first-episode psychosis. <i>Psychiatry Research</i> , 2017, 250, 38-43. | 3.3 | 26 |
| 21 | The course of negative symptoms in first-episode schizophrenia and its predictors: A prospective two-year follow-up study. <i>Schizophrenia Research</i> , 2017, 189, 84-90. | 2.0 | 49 |
| 22 | Increased nucleus accumbens volume in first-episode psychosis. <i>Psychiatry Research - Neuroimaging</i> , 2017, 263, 57-60. | 1.8 | 11 |
| 23 | Effort-based reward task, a behavioral measure to study negative symptoms in schizophrenia. <i>European Psychiatry</i> , 2017, 41, S343-S344. | 0.2 | 0 |
| 24 | Default Mode Network Aberrant Connectivity Associated with Neurological Soft Signs in Schizophrenia Patients and Unaffected Relatives. <i>Frontiers in Psychiatry</i> , 2017, 8, 298. | 2.6 | 29 |
| 25 | Social withdrawal and suicide risk: A descriptive study. <i>European Psychiatry</i> , 2016, 33, S175-S175. | 0.2 | 1 |
| 26 | Age at First Episode Modulates Diagnosis-Related Structural Brain Abnormalities in Psychosis. <i>Schizophrenia Bulletin</i> , 2016, 42, 344-357. | 4.3 | 58 |
| 27 | Predictors of Relapse and Functioning in First-Episode Psychosis: A Two-Year Follow-Up Study. <i>Psychiatric Services</i> , 2016, 67, 227-233. | 2.0 | 69 |
| 28 | Association between neurological soft signs, temperament and character in patients with schizophrenia and non-psychotic relatives. <i>PeerJ</i> , 2016, 4, e1651. | 2.0 | 10 |
| 29 | The Neuroanatomical Basis of Panic Disorder and Social Phobia in Schizophrenia: A Voxel Based Morphometric Study. <i>PLoS ONE</i> , 2015, 10, e0119847. | 2.5 | 6 |
| 30 | Schizophrenia, antipsychotic drugs and cardiovascular risk: Descriptive study in primary care. <i>European Psychiatry</i> , 2015, 30, 535-541. | 0.2 | 11 |
| 31 | Premorbid adjustment and clinical correlates of cognitive impairment in first-episode psychosis. The PEPsCog Study. <i>Schizophrenia Research</i> , 2015, 164, 65-73. | 2.0 | 92 |
| 32 | Patient Management and Psychopharmacological Treatment Associated to Smoking Ban in an Acute Psychiatric Unit. <i>Community Mental Health Journal</i> , 2015, 51, 746-752. | 2.0 | 4 |
| 33 | The negative syndrome of schizophrenia: Three -underlying components are better than two. <i>Schizophrenia Research</i> , 2015, 166, 115-118. | 2.0 | 18 |
| 34 | Relationship between cannabis and psychosis: Reasons for use and associated clinical variables. <i>Psychiatry Research</i> , 2015, 229, 70-74. | 3.3 | 43 |
| 35 | P.3.b.002 Cortex morphology and subcortical brain grey matter deficits in schizophrenia and unaffected relatives. <i>European Neuropsychopharmacology</i> , 2015, 25, S463-S464. | 0.7 | 0 |
| 36 | Limbic activity in antipsychotic naïve first-episode psychotic subjects during facial emotion discrimination. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2014, 264, 271-283. | 3.2 | 18 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Spanish adaptation and validation of the Brief Negative Symptoms Scale. <i>Comprehensive Psychiatry</i> , 2014, 55, 1726-1729. | 3.1 | 54 |
| 38 | P.3.b.035 Temperament, character and neurological soft signs in patients with schizophrenia and unaffected siblings. <i>European Neuropsychopharmacology</i> , 2013, 23, S450. | 0.7 | 0 |
| 39 | P.3.f.015 Abnormal functioning of the default mode network in schizophrenia and unaffected relatives: a study of functional magnetic resonance. <i>European Neuropsychopharmacology</i> , 2013, 23, S500-S501. | 0.7 | 0 |
| 40 | Poster #189 PREDICTION OF RELAPSE AFTER A FIRST EPISODE OF PSYCHOSIS: A FOLLOW-UP CLINICAL STUDY. <i>Schizophrenia Research</i> , 2012, 136, S159. | 2.0 | 0 |
| 41 | Gray matter volume deficits and correlation with insight and negative symptoms in first-psychotic-episode subjects. <i>Acta Psychiatrica Scandinavica</i> , 2011, 123, 431-439. | 4.5 | 81 |
| 42 | Neural correlates of impaired emotional discrimination in borderline personality disorder: An fMRI study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 1537-1545. | 4.8 | 61 |
| 43 | Diagnostic Stability and Clinical Characteristics in First-episode Psychosis. <i>European Psychiatry</i> , 2009, 24, . | 0.2 | 0 |
| 44 | Onset of Mania After CPAP in a Man With Obstructive Sleep Apnea. <i>Psychosomatics</i> , 2008, 49, 447-449. | 2.5 | 11 |
| 45 | Abnormal P300 in people with high risk of developing psychosis. <i>NeuroImage</i> , 2008, 41, 553-560. | 4.2 | 87 |
| 46 | Dual Diagnosis in an Inpatient Drug-Abuse Detoxification Unit. <i>European Addiction Research</i> , 2007, 13, 192-200. | 2.4 | 18 |
| 47 | WC5G NEUROPHYSIOLOGIC MARKERS OF RISK OF DEVELOPING PSYCHOSIS. <i>Schizophrenia Research</i> , 2006, 86, S14. | 2.0 | 0 |
| 48 | The relationship between frequency and intensity of fears and a collagen condition. <i>Depression and Anxiety</i> , 2006, 23, 412-417. | 4.1 | 29 |