

Hyoun-Ah Kim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8430389/publications.pdf>

Version: 2024-02-01

114
papers

2,274
citations

236925

25
h-index

276875

41
g-index

116
all docs

116
docs citations

116
times ranked

3072
citing authors

#	ARTICLE	IF	CITATIONS
1	Cytokine IL-6 and IL-10 as Biomarkers in Systemic Lupus Erythematosus. <i>Journal of Clinical Immunology</i> , 2007, 27, 461-466.	3.8	321
2	Dickkopf-1 level is lower in patients with ankylosing spondylitis than in healthy people and is not influenced by anti-tumor necrosis factor therapy. <i>Rheumatology International</i> , 2012, 32, 2523-2527.	3.0	98
3	Therapeutic responses and prognosis in adult-onset Still's disease. <i>Rheumatology International</i> , 2012, 32, 1291-1298.	3.0	64
4	Serum S100A8/A9, But Not Follistatin-like Protein 1 and Interleukin 18, May Be a Useful Biomarker of Disease Activity in Adult-onset Still's Disease. <i>Journal of Rheumatology</i> , 2012, 39, 1399-1406.	2.0	60
5	C-reactive Protein Is a More Sensitive and Specific Marker for Diagnosing Bacterial Infections in Systemic Lupus Erythematosus Compared to S100A8/A9 and Procalcitonin. <i>Journal of Rheumatology</i> , 2012, 39, 728-734.	2.0	55
6	Korean Observational Study Network for Arthritis (KORONA): Establishment of a Prospective Multicenter Cohort for Rheumatoid Arthritis in South Korea. <i>Seminars in Arthritis and Rheumatism</i> , 2012, 41, 745-751.	3.4	54
7	Vitamin D may not be a good marker of disease activity in Korean patients with systemic lupus erythematosus. <i>Rheumatology International</i> , 2011, 31, 1189-1194.	3.0	51
8	Determinants of quality of life in patients with fibromyalgia: A structural equation modeling approach. <i>PLoS ONE</i> , 2017, 12, e0171186.	2.5	51
9	Elevated circulating levels of the interferon- γ -induced chemokines are associated with disease activity and cutaneous manifestations in adult-onset Still's disease. <i>Scientific Reports</i> , 2017, 7, 46652.	3.3	49
10	Serum S100A12 May Be a Useful Biomarker of Disease Activity in Adult-onset Still's Disease. <i>Journal of Rheumatology</i> , 2014, 41, 2403-2408.	2.0	47
11	Prevalence of work-related symptoms and serum-specific antibodies to wheat flour in exposed workers in the bakery industry. <i>Respiratory Medicine</i> , 2008, 102, 548-555.	2.9	41
12	Association of CXCL10 and CXCL13 levels with disease activity and cutaneous manifestation in active adult-onset Still's disease. <i>Arthritis Research and Therapy</i> , 2015, 17, 260.	3.5	39
13	Cytokines and their receptors as biomarkers of systemic lupus erythematosus. <i>Expert Review of Molecular Diagnostics</i> , 2008, 8, 189-198.	3.1	38
14	Prevalence of Human Papilloma Virus Infections and Cervical Cytological Abnormalities among Korean Women with Systemic Lupus Erythematosus. <i>Journal of Korean Medical Science</i> , 2010, 25, 1431.	2.5	37
15	TLR4 Endogenous Ligand S100A8/A9 Levels in Adult-Onset Still's Disease and Their Association with Disease Activity and Clinical Manifestations. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1342.	4.1	37
16	Systemic immune-inflammation index combined with ferritin can serve as a reliable assessment score for adult-onset Still's disease. <i>Clinical Rheumatology</i> , 2021, 40, 661-668.	2.2	37
17	The Pathologic Findings of Skin, Lymph Node, Liver, and Bone Marrow in Patients With Adult-Onset Still Disease. <i>Medicine (United States)</i> , 2015, 94, e787.	1.0	36
18	Usefulness of neutrophil-to-lymphocyte ratio as a biomarker for diagnosing infections in patients with systemic lupus erythematosus. <i>Clinical Rheumatology</i> , 2017, 36, 2479-2485.	2.2	36

#	ARTICLE	IF	CITATIONS
19	Beta2-Microglobulin Can Be a Disease Activity Marker in Systemic Lupus Erythematosus. American Journal of the Medical Sciences, 2010, 339, 337-340.	1.1	32
20	Women with Fibromyalgia Have Lower Levels of Calcium, Magnesium, Iron and Manganese in Hair Mineral Analysis. Journal of Korean Medical Science, 2011, 26, 1253.	2.5	32
21	Prevalence and Fracture Risk of Osteoporosis in Patients with Rheumatoid Arthritis: A Multicenter Comparative Study of the FRAX and WHO Criteria. Journal of Clinical Medicine, 2018, 7, 507.	2.4	32
22	Neutrophil Extracellular Traps May Contribute to the Pathogenesis in Adult-onset Still Disease. Journal of Rheumatology, 2019, 46, 1560-1569.	2.0	31
23	Associated clinical factors for serious infections in patients with systemic lupus erythematosus. Scientific Reports, 2019, 9, 9704.	3.3	30
24	Polymorphisms of the <i>TRPV2</i> and <i>TRPV3</i> genes associated with fibromyalgia in a Korean population. Rheumatology, 2016, 55, 1518-1527.	1.9	27
25	The neutrophil-to-lymphocyte ratio could be a good diagnostic marker and predictor of relapse in patients with adult-onset Still's disease. Medicine (United States), 2017, 96, e7546.	1.0	27
26	Treatment of Atopic Dermatitis with a Combination of Allergen-Specific Immunotherapy and a Histamine-Immunoglobulin Complex. International Archives of Allergy and Immunology, 2008, 146, 235-240.	2.1	26
27	Elevated Salivary Alpha-Amylase Level, Association Between Depression and Disease Activity, and Stress as a Predictor of Disease Flare in Systemic Lupus Erythematosus. Medicine (United States), 2015, 94, e1184.	1.0	26
28	Clinical characteristics of lupus myocarditis in Korea. Rheumatology International, 2008, 28, 275-280.	3.0	25
29	C-Reactive Protein Gene Polymorphisms in Disease Susceptibility and Clinical Manifestations of Korean Systemic Lupus Erythematosus. Journal of Rheumatology, 2009, 36, 2238-2243.	2.0	25
30	Lupus Low Disease Activity State Achievement Is Important for Reducing Adverse Outcomes in Pregnant Patients With Systemic Lupus Erythematosus. Journal of Rheumatology, 2021, 48, 707-716.	2.0	23
31	An Antinuclear Antibody-Negative Patient With Lupus Nephritis. Korean Journal of Internal Medicine, 2009, 24, 76.	1.7	23
32	Interleukin 6 Gene Polymorphisms Are Associated with Systemic Lupus Erythematosus in Koreans. Journal of Rheumatology, 2010, 37, 2251-2258.	2.0	22
33	Serum Growth Arrest-Specific Protein 6 Levels are a Reliable Biomarker of Disease Activity in Systemic Lupus Erythematosus. Journal of Clinical Immunology, 2013, 33, 143-150.	3.8	22
34	Neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio are associated with disease activity in polymyalgia rheumatica. Journal of Clinical Laboratory Analysis, 2019, 33, e23000.	2.1	21
35	Elevated high-mobility group B1 levels in active adult-onset Still's disease associated with systemic score and skin rash. Clinical Rheumatology, 2016, 35, 1937-1942.	2.2	20
36	Switching profiles in a population-based cohort of rheumatoid arthritis receiving biologic therapy: results from the KOBIO registry. Clinical Rheumatology, 2017, 36, 1013-1022.	2.2	20

#	ARTICLE	IF	CITATIONS
37	Roles of Interactions Between Toll-Like Receptors and Their Endogenous Ligands in the Pathogenesis of Systemic Juvenile Idiopathic Arthritis and Adult-Onset Still's Disease. <i>Frontiers in Immunology</i> , 2020, 11, 583513.	4.8	20
38	The role of damage-associated molecular pattern for pathogenesis and biomarkers in adult-onset Still's disease. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 459-468.	3.1	19
39	Tocilizumab-induced anaphylaxis in patients with adult-onset Still's disease and systemic juvenile idiopathic arthritis: a case-based review. <i>Rheumatology International</i> , 2020, 40, 791-798.	3.0	19
40	KOBIO, the First Web-based Korean Biologics Registry Operated With a Unified Platform Among Distinct Disease Entities. <i>Journal of Rheumatic Diseases</i> , 2021, 28, 176-182.	1.1	19
41	Smoking habits influence pain and functional and psychiatric features in fibromyalgia. <i>Joint Bone Spine</i> , 2011, 78, 259-265.	1.6	18
42	Multicenter Retrospective Analysis of the Effectiveness and Safety of Rituximab in Korean Patients with Refractory Systemic Lupus Erythematosus. <i>Autoimmune Diseases</i> , 2012, 2012, 1-6.	0.6	18
43	Effect of fenofibrate on uric acid level in patients with gout. <i>Scientific Reports</i> , 2018, 8, 16767.	3.3	17
44	Off-label use of tocilizumab to treat non-juvenile idiopathic arthritis in pediatric rheumatic patients: a literature review. <i>Pediatric Rheumatology</i> , 2018, 16, 79.	2.1	17
45	An Update on the Pathogenic Role of Neutrophils in Systemic Juvenile Idiopathic Arthritis and Adult-Onset Still's Disease. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13038.	4.1	17
46	Obesity increases the incidence of new-onset lupus nephritis and organ damage during follow-up in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020, 29, 578-586.	1.6	16
47	The antichromatin antibodies can be useful as a diagnostic tool and disease activity marker of systemic lupus erythematosus in Koreans. <i>Clinical Immunology</i> , 2008, 128, 277-283.	3.2	15
48	Liver X receptors alpha gene (NR1H3) promoter polymorphisms are associated with systemic lupus erythematosus in Koreans. <i>Arthritis Research and Therapy</i> , 2014, 16, R112.	3.5	15
49	Serum Levels of Interleukin 33 and Soluble ST2 Are Associated with the Extent of Disease Activity and Cutaneous Manifestations in Patients with Active Adult-onset Still's Disease. <i>Journal of Rheumatology</i> , 2017, 44, 740-747.	2.0	15
50	Histopathology and expression of the chemokines CXCL10, CXCL13, and CXCR3 and the endogenous TLR-4 ligand S100A8/A9 in lymph nodes of patients with adult-onset Still's disease. <i>Scientific Reports</i> , 2019, 9, 7517.	3.3	15
51	Identification of Dioscorea Batatas (Sanyak) Allergen as an Inhalant and Oral Allergen. <i>Journal of Korean Medical Science</i> , 2008, 23, 72.	2.5	14
52	Seasonality of Gout in Korea: A Multicenter Study. <i>Journal of Korean Medical Science</i> , 2015, 30, 240.	2.5	14
53	Outcome and predictors of renal survival in patients with lupus nephritis: Comparison between cyclophosphamide and mycophenolate mofetil. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 1031-1039.	1.9	14
54	The Practice Guideline for Vaccinating Korean Patients with Autoimmune Inflammatory Rheumatic Disease. <i>Infection and Chemotherapy</i> , 2020, 52, 252.	2.3	14

#	ARTICLE	IF	CITATIONS
55	Salivary Biomarkers in Patients with Sjögren's Syndrome—A Systematic Review. <i>International Journal of Molecular Sciences</i> , 2021, 22, 12903.	4.1	14
56	Characteristic patterns of HLA presentation and T cell differentiation in adult-onset Still's disease. <i>International Journal of Immunopathology and Pharmacology</i> , 2018, 32, 205873841879128.	2.1	13
57	Trabecular Bone Score Is a Useful Parameter for the Prediction of Vertebral Fractures in Patients With Polymyalgia Rheumatica. <i>Journal of Clinical Densitometry</i> , 2020, 23, 373-380.	1.2	13
58	Retention Rate and Safety of Biosimilar CT-P13 in Rheumatoid Arthritis: Data from the Korean College of Rheumatology Biologics Registry. <i>BioDrugs</i> , 2020, 34, 89-98.	4.6	13
59	Induction of Remission is Difficult due to Frequent Relapse during Tapering Steroids in Korean Patients with Polymyalgia Rheumatica. <i>Journal of Korean Medical Science</i> , 2012, 27, 22.	2.5	12
60	Flare of adult-onset Still's disease following mRNA COVID-19 vaccination: a case report and review of literature. <i>Clinical Rheumatology</i> , 2022, 41, 1583-1589.	2.2	12
61	Association between brain-derived neurotrophic factor gene polymorphisms and fibromyalgia in a Korean population: a multicenter study. <i>Arthritis Research and Therapy</i> , 2018, 20, 220.	3.5	11
62	Mindfulness-based cognitive therapy in Korean patients with systemic lupus erythematosus: A pilot study. <i>Complementary Therapies in Clinical Practice</i> , 2019, 35, 18-21.	1.7	11
63	Body mass index and glucocorticoid dose contribute to subclinical atherosclerosis in Korean patients with systemic lupus erythematosus: A prospective 4-year follow-up study. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 1410-1418.	1.9	11
64	Association of β_2 -Adrenergic Receptor Polymorphism with the Phenotype of Aspirin-Intolerant Acute Urticaria. <i>Yonsei Medical Journal</i> , 2007, 48, 1079.	2.2	10
65	Autoantibodies to C-Reactive Protein in Incomplete Lupus and Systemic Lupus Erythematosus. <i>Journal of Investigative Medicine</i> , 2014, 62, 890-893.	1.6	10
66	Characteristics and outcomes of rheumatoid arthritis patients who started biosimilar infliximab. <i>Rheumatology International</i> , 2017, 37, 1007-1014.	3.0	10
67	Highly Expression of CD11b and CD32 on Peripheral Blood Mononuclear Cells from Patients with Adult-Onset Still's Disease. <i>International Journal of Molecular Sciences</i> , 2017, 18, 202.	4.1	10
68	Seasonal Variations and Associated Factors of Gout Attacks: a Prospective Multicenter Study in Korea. <i>Journal of Korean Medical Science</i> , 2020, 35, e133.	2.5	10
69	S100A8 in Serum, Urine, and Saliva as a Potential Biomarker for Systemic Lupus Erythematosus. <i>Frontiers in Immunology</i> , 2022, 13, 886209.	4.8	10
70	Effects of risk factors for and components of metabolic syndrome on the quality of life of patients with systemic lupus erythematosus: a structural equation modeling approach. <i>Quality of Life Research</i> , 2018, 27, 105-113.	3.1	9
71	The Assessment of Muscle Mass and Function in Patients with Long-Standing Rheumatoid Arthritis. <i>Journal of Clinical Medicine</i> , 2021, 10, 3458.	2.4	9
72	An unusual clinical manifestation of Takayasu's arteritis: Spinal cord compression. <i>Joint Bone Spine</i> , 2009, 76, 209-212.	1.6	8

#	ARTICLE	IF	CITATIONS
73	Comparative effectiveness of treatment options after conventional DMARDs failure in rheumatoid arthritis. <i>Rheumatology International</i> , 2017, 37, 975-982.	3.0	8
74	The Correlation of CD206, CD209, and Disease Severity in Behçet's Disease with Arthritis. <i>Mediators of Inflammation</i> , 2017, 2017, 1-13.	3.0	8
75	Clinical characteristics and role of whole-body bone scan in multifocal osteonecrosis. <i>BMC Musculoskeletal Disorders</i> , 2019, 20, 23.	1.9	8
76	Retention Rate and Efficacy of the Biosimilar CT-P13 Versus Reference Infliximab in Patients with Ankylosing Spondylitis: A Propensity Score-Matched Analysis from the Korean College of Rheumatology Biologics Registry. <i>BioDrugs</i> , 2020, 34, 529-539.	4.6	7
77	Association of depression with socioeconomic status, anticardiolipin antibodies, and organ damage in patients with systemic lupus erythematosus: results from the KORNET registry. <i>Clinical and Experimental Rheumatology</i> , 2018, 36, 627-635.	0.8	7
78	Prevalence and Factors of Osteoporosis and High Risk of Osteoporotic Fracture in Patients with Ankylosing Spondylitis: A Multicenter Comparative Study of Bone Mineral Density and the Fracture Risk Assessment Tool. <i>Journal of Clinical Medicine</i> , 2022, 11, 2830.	2.4	7
79	Prediction for TNF Inhibitor Users in RA Patients According to Reimbursement Criteria Based on DAS28. <i>Journal of Rheumatic Diseases</i> , 2014, 21, 64.	1.1	6
80	FCI3RIIB Gene Polymorphisms Are Associated with Disease Risk and Clinical Manifestations of Systemic Lupus Erythematosus in Koreans. <i>Tohoku Journal of Experimental Medicine</i> , 2015, 236, 185-191.	1.2	6
81	Application of the international league against rheumatism classification criteria for systemic juvenile idiopathic arthritis as a prognostic factor in patients with adults-onset Still's disease. <i>Pediatric Rheumatology</i> , 2018, 16, 9.	2.1	6
82	Comparison of Retention Rates Between Tumor Necrosis Factor- α Inhibitors in Patients With Ankylosing Spondylitis: Data From the Korean College of Rheumatology Biologics Registry. <i>Frontiers in Medicine</i> , 2021, 8, 689609.	2.6	6
83	Retention rate and long-term safety of biosimilar CT-P13 in patients with ankylosing spondylitis: data from the Korean College of Rheumatology Biologics registry. <i>Clinical and Experimental Rheumatology</i> , 2020, 38, 267-274.	0.8	6
84	Role of bone scan in the assessment of polymyositis/dermatomyositis. <i>Clinical Rheumatology</i> , 2015, 34, 699-706.	2.2	5
85	Impact of Childbearing Decisions on Family Size of Korean Women with Systemic Lupus Erythematosus. <i>Journal of Korean Medical Science</i> , 2016, 31, 729.	2.5	5
86	Clinical Impact of the Fracture Risk Assessment Tool on the Treatment Decision for Osteoporosis in Patients with Knee Osteoarthritis: A Multicenter Comparative Study of the Fracture Risk Assessment Tool and World Health Organization Criteria. <i>Journal of Clinical Medicine</i> , 2019, 8, 918.	2.4	5
87	Rituximab biosimilar CT-P10 for the treatment of rheumatoid arthritis. <i>Expert Opinion on Biological Therapy</i> , 2019, 19, 979-986.	3.1	5
88	Prevalence of osteoporosis in patients with systemic lupus erythematosus: A multicenter comparative study of the World Health Organization and fracture risk assessment tool criteria. <i>Osteoporosis and Sarcopenia</i> , 2020, 6, 173-178.	1.9	5
89	Anti-Inflammatory Effects of Low-Dose Glucocorticoids Compensate for Their Detrimental Effects on Bone Mineral Density in Patients with Rheumatoid Arthritis. <i>Journal of Clinical Medicine</i> , 2021, 10, 2944.	2.4	5
90	Standardized, musculoskeletal ultrasonographic reference values for healthy Korean adults. <i>Korean Journal of Internal Medicine</i> , 2019, 34, 1372-1380.	1.7	5

#	ARTICLE	IF	CITATIONS
91	Three Clinical Clusters Identified through Hierarchical Cluster Analysis Using Initial Laboratory Findings in Korean Patients with Systemic Lupus Erythematosus. <i>Journal of Clinical Medicine</i> , 2022, 11, 2406.	2.4	5
92	Nonsteroidal Antiinflammatory Drugs (NSAID) Versus NSAID with Hydroxychloroquine in Treatment of Chemotherapy-related Arthropathy: Open-label Multicenter Pilot Study. <i>Journal of Rheumatology</i> , 2012, 39, 1902-1903.	2.0	4
93	Patient with Systemic Lupus Erythematosus Combined with Erosive Arthritis was Treated Successfully with Tocilizumab: A Case Report. <i>Journal of Rheumatic Diseases</i> , 2018, 25, 144.	1.1	4
94	The Liver X Receptor Is Upregulated in Monocyte-Derived Macrophages and Modulates Inflammatory Cytokines Based on LXR Polymorphism. <i>Mediators of Inflammation</i> , 2019, 2019, 1-12.	3.0	4
95	Severity of fibromyalgia symptoms is associated with socioeconomic status and not obesity in Korean patients. <i>Clinical and Experimental Rheumatology</i> , 2016, 34, S83-8.	0.8	4
96	The levels of CXCL12 and its receptor, CXCR4, as a biomarker of disease activity and cutaneous manifestation in adult-onset Still's disease. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 121, 67-73.	0.8	4
97	Chronic periaortitis with antiphospholipid syndrome. <i>International Journal of Rheumatic Diseases</i> , 2010, 13, 91-93.	1.9	3
98	Serum growth arrest-specific protein 6 levels are elevated in adult-onset Still's disease. <i>Clinical Rheumatology</i> , 2014, 33, 865-868.	2.2	3
99	Biomarkers for adult-onset still's disease: up-to-date. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 655-657.	3.1	3
100	Differences in Expression of Human Leukocyte Antigen Class II Subtypes and T Cell Subsets in Behçet's Disease with Arthritis. <i>International Journal of Molecular Sciences</i> , 2019, 20, 5044.	4.1	3
101	The Role of Natural Killer Cells in Inflammation in Active Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2019, 46, 1253-1255.	2.0	3
102	Salivary Immunoglobulin Gamma-3 Chain C Is a Promising Noninvasive Biomarker for Systemic Lupus Erythematosus. <i>International Journal of Molecular Sciences</i> , 2021, 22, 1374.	4.1	3
103	Biomarkers for systemic lupus erythematosus: an update. <i>International Journal of Clinical Rheumatology</i> , 2015, 10, 195-204.	0.3	2
104	The Practice Guideline for Vaccinating Korean Patients With Autoimmune Inflammatory Rheumatic Disease. <i>Journal of Rheumatic Diseases</i> , 2020, 27, 182-202.	1.1	2
105	Intestinal pseudo-obstruction caused by neuromyopathy in a patient with systemic sclerosis. <i>Modern Rheumatology</i> , 2012, 22, 912-918.	1.8	2
106	Implications of Persistent Pain in Patients With Rheumatoid Arthritis Despite Remission Status: Data From the KOBIO Registry. <i>Journal of Rheumatic Diseases</i> , 2022, , .	1.1	2
107	SLE developed in a preadolescent child with Klinefelter syndrome who had no detectable sex hormone. <i>Rheumatology International</i> , 2009, 29, 975-977.	3.0	1
108	An update: current and prospective treatments for adult-onset Still's disease. <i>Expert Opinion on Orphan Drugs</i> , 2017, 5, 421-429.	0.8	1

#	ARTICLE	IF	CITATIONS
109	Clinical Features and Drug Retention of TNF Inhibitors in Older Patients with Ankylosing Spondylitis: Results from the KOBIO Registry. BioDrugs, 2022, , 1.	4.6	1
110	Comparison of remission criteria in patients with rheumatoid arthritis treated with biologic or targeted synthetic disease-modifying anti-rheumatic drugs: results from a nationwide registry. Therapeutic Advances in Musculoskeletal Disease, 2022, 14, 1759720X2210963.	2.7	1
111	Elevated expression of TLR2 and its correlation with disease activity and clinical manifestations in adult-onset Still's disease. Scientific Reports, 2022, 12, .	3.3	1
112	Correlations between salivary gland scintigraphy and histopathologic data of salivary glands in patients with primary Sjogren's syndrome. Clinical Rheumatology, 2022, 41, 3083-3093.	2.2	1
113	Neurogenic Bladder in a Patient With Systemic Lupus Erythematosus and Cerebral Involvement. Journal of Clinical Rheumatology, 2009, 15, 40-41.	0.9	0
114	Les habitudes tabagiques influencent la douleur et les Œ©ments fonctionnels et psychiatriques dans la fibromyalgie. Revue Du Rhumatisme (Edition Francaise), 2011, 78, 173-179.	0.0	0