

Miguel Xavier

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8472883/publications.pdf>

Version: 2024-02-01

90
papers

6,558
citations

136950

32
h-index

69250

77
g-index

95
all docs

95
docs citations

95
times ranked

10018
citing authors

#	ARTICLE	IF	CITATIONS
1	The Prevalence and Correlates of Binge Eating Disorder in the World Health Organization World Mental Health Surveys. <i>Biological Psychiatry</i> , 2013, 73, 904-914.	1.3	1,025
2	Barriers to mental health treatment: results from the WHO World Mental Health surveys. <i>Psychological Medicine</i> , 2014, 44, 1303-1317.	4.5	825
3	Posttraumatic stress disorder in the World Mental Health Surveys. <i>Psychological Medicine</i> , 2017, 47, 2260-2274.	4.5	675
4	The descriptive epidemiology of DSM-IV Adult ADHD in the World Health Organization World Mental Health Surveys. <i>ADHD Attention Deficit and Hyperactivity Disorders</i> , 2017, 9, 47-65.	1.7	507
5	Anxious and non-anxious major depressive disorder in the World Health Organization World Mental Health Surveys. <i>Epidemiology and Psychiatric Sciences</i> , 2015, 24, 210-226.	3.9	355
6	The cross-national epidemiology of social anxiety disorder: Data from the World Mental Health Survey Initiative. <i>BMC Medicine</i> , 2017, 15, 143.	5.5	258
7	Affective disorders and risk of developing dementia: systematic review. <i>British Journal of Psychiatry</i> , 2013, 202, 177-186.	2.8	249
8	Prevalence of common mental disorders in general practice attendees across Europe. <i>British Journal of Psychiatry</i> , 2008, 192, 362-367.	2.8	227
9	The cross-national epidemiology of specific phobia in the World Mental Health Surveys. <i>Psychological Medicine</i> , 2017, 47, 1744-1760.	4.5	148
10	Development and Validation of an International Risk Prediction Algorithm for Episodes of Major Depression in General Practice Attendees. <i>Archives of General Psychiatry</i> , 2008, 65, 1368.	12.3	134
11	The association between childhood adversities and subsequent first onset of psychotic experiences: a cross-national analysis of 23 998 respondents from 17 countries. <i>Psychological Medicine</i> , 2017, 47, 1230-1245.	4.5	108
12	Associations between unemployment and major depressive disorder: Evidence from an international, prospective study (the predict cohort). <i>Social Science and Medicine</i> , 2011, 73, 1627-1634.	3.8	105
13	Associations between mental disorders and subsequent onset of hypertension. <i>General Hospital Psychiatry</i> , 2014, 36, 142-149.	2.4	102
14	Suicidal thoughts and behaviors among college students and same-aged peers: results from the World Health Organization World Mental Health Surveys. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2018, 53, 279-288.	3.1	79
15	Association of Huntington's disease and schizophrenia-like psychosis in a Huntington's disease pedigree. <i>Clinical Practice and Epidemiology in Mental Health</i> , 2006, 2, 1.	1.2	78
16	Role of common mental and physical disorders in partial disability around the world. <i>British Journal of Psychiatry</i> , 2012, 200, 454-461.	2.8	77
17	Association Between Psychotic Experiences and Subsequent Suicidal Thoughts and Behaviors. <i>JAMA Psychiatry</i> , 2017, 74, 1136.	11.0	75
18	Prediction of depression in European general practice attendees: the PREDICT study. <i>BMC Public Health</i> , 2006, 6, 6.	2.9	71

#	ARTICLE	IF	CITATIONS
19	Social and clinical factors influencing the choice of coping strategies in relatives of patients with schizophrenia: results of the BIOMED I study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 1998, 33, 413-419.	3.1	62
20	The association between psychotic experiences and disability: results from the <sc>WHO</sc> World Mental Health Surveys. <i>Acta Psychiatrica Scandinavica</i> , 2017, 136, 74-84.	4.5	58
21	MAJOR DEPRESSIVE DISORDER SUBTYPES TO PREDICT LONG-TERM COURSE. <i>Depression and Anxiety</i> , 2014, 31, 765-777.	4.1	56
22	Obsessiveâ€compulsive symptoms in primary focal dystonia: A controlled study. <i>Movement Disorders</i> , 2011, 26, 2274-2278.	3.9	51
23	Spiritual and religious beliefs as risk factors for the onset of major depression: an international cohort study. <i>Psychological Medicine</i> , 2013, 43, 2109-2120.	4.5	50
24	Depression, anxiety and physical function: exploring the strength of causality. <i>Journal of Epidemiology and Community Health</i> , 2012, 66, e25-e25.	3.7	47
25	Unmet needs in the management of schizophrenia. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 97.	2.2	47
26	A comparative analysis of role attainment and impairment in binge-eating disorder and bulimia nervosa: results from the WHO World Mental Health Surveys. <i>Epidemiology and Psychiatric Sciences</i> , 2014, 23, 27-41.	3.9	46
27	Setâ€shifting and behavioral dysfunction in primary focal dystonia. <i>Movement Disorders</i> , 2008, 23, 200-206.	3.9	45
28	Predicting the onset of major depression in primary care: international validation of a risk prediction algorithm from Spain. <i>Psychological Medicine</i> , 2011, 41, 2075-2088.	4.5	45
29	Psychosocial wellbeing and psychiatric care in the European Communities: analysis of macro indicators. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2004, 39, 883-892.	3.1	42
30	Impact of psychosis on Portuguese caregivers: a cross-cultural exploration of burden, distress, positive aspects and clinical-functional correlates. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013, 48, 325-335.	3.1	42
31	The influence of clerkship on studentsâ€™ stigma towards mental illness: a meta-analysis. <i>Medical Education</i> , 2018, 52, 694-704.	2.1	39
32	Implementing the World Mental Health Survey Initiative in Portugal â€“ rationale, design and fieldwork procedures. <i>International Journal of Mental Health Systems</i> , 2013, 7, 19.	2.7	38
33	Arachnoid cyst in a patient with psychosis: Case report. <i>Annals of General Psychiatry</i> , 2007, 6, 16.	2.7	37
34	A Crossâ€National Examination of Differences in Classification of Lifetime Alcohol Use Disorder Between <sc>DSM</sc>â€IV</sc> and <sc>DSM</sc>â€5: Findings from the World Mental Health Survey. <i>Alcoholism: Clinical and Experimental Research</i> , 2016, 40, 1728-1736.	2.4	35
35	Comparison of risk factors for the onset and maintenance of depression. <i>British Journal of Psychiatry</i> , 2010, 196, 13-17.	2.8	31
36	The Associations Between Preexisting Mental Disorders andâ€Subsequent Onset of Chronic Headaches: A Worldwide Epidemiologic Perspective. <i>Journal of Pain</i> , 2015, 16, 42-52.	1.4	29

#	ARTICLE	IF	CITATIONS
37	Impact of clerkship in the attitudes toward psychiatry among Portuguese medical students. BMC Medical Education, 2010, 10, 56.	2.4	27
38	Proportion of patients without mental disorders being treated in mental health services worldwide. British Journal of Psychiatry, 2015, 206, 101-109.	2.8	27
39	Reducing the Number of People with Dementia Through Primary Prevention in Mozambique, Brazil, and Portugal: An Analysis of Population-Based Data. Journal of Alzheimer's Disease, 2019, 70, S283-S291.	2.6	27
40	Disability Mediates the Impact of Common Conditions on Perceived Health. PLoS ONE, 2013, 8, e65858.	2.5	27
41	An international risk prediction algorithm for the onset of generalized anxiety and panic syndromes in general practice attendees: predictA. Psychological Medicine, 2011, 41, 1625-1639.	4.5	26
42	Differential Impact of Risk Factors for Women and Men on the Risk of Major Depressive Disorder. Annals of Epidemiology, 2012, 22, 388-396.	1.9	26
43	The cross-national epidemiology of DSM-IV intermittent explosive disorder. Psychological Medicine, 2016, 46, 3161-3172.	4.5	26
44	The effects of co-morbidity in defining major depression subtypes associated with long-term course and severity. Psychological Medicine, 2014, 44, 3289-3302.	4.5	25
45	Days out of role due to common physical and mental conditions in Portugal: Results from the WHO World Mental Health Survey. BJPsych Open, 2017, 3, 15-21.	0.7	24
46	The prevalence of dementia in a Portuguese community sample: a 10/66 Dementia Research Group study. BMC Geriatrics, 2017, 17, 261.	2.7	24
47	Schizophrenia and quality of life: a one-year follow-up in four EU countries. BMC Psychiatry, 2006, 6, 39.	2.6	22
48	Heavy Episodic Drinking in Europe: A Cross Section Study in Primary Care in Six European Countries. Alcohol and Alcoholism, 2011, 46, 600-606.	1.6	22
49	Patterns of care and dropout rates from outpatient mental healthcare in low-, middle- and high-income countries from the World Health Organization's World Mental Health Survey Initiative. Psychological Medicine, 2021, 51, 2104-2116.	4.5	21
50	Recent life events pose greatest risk for onset of major depressive disorder during mid-life. Journal of Affective Disorders, 2012, 136, 505-513.	4.1	19
51	The role of religious advisors in mental health care in the World Mental Health surveys. Social Psychiatry and Psychiatric Epidemiology, 2017, 52, 353-367.	3.1	19
52	Predicting onset of major depression in general practice attendees in Europe: extending the application of the predictD risk algorithm from 12 to 24 months. Psychological Medicine, 2013, 43, 1929-1939.	4.5	17
53	Associations between DSM-IV mental disorders and subsequent non-fatal, self-reported stroke. Journal of Psychosomatic Research, 2015, 79, 130-136.	2.6	17
54	Barriers and facilitators to implementing a continuing medical education intervention in a primary health care setting. BMC Health Services Research, 2022, 22, 638.	2.2	16

#	ARTICLE	IF	CITATIONS
55	Risk factors for onset of multiple or long major depressive episodes versus single and short episodes. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013, 48, 1067-1075.	3.1	13
56	A Gaussian random field model for similarity-based smoothing in Bayesian disease mapping. <i>Statistical Methods in Medical Research</i> , 2016, 25, 1166-1184.	1.5	13
57	Substance dependence among those without symptoms of substance abuse in the World Mental Health Survey. <i>International Journal of Methods in Psychiatric Research</i> , 2017, 26, .	2.1	13
58	Sudden psychotic episode probably due to meningoencephalitis and <i>Chlamydia pneumoniae</i> acute infection. <i>Clinical Practice and Epidemiology in Mental Health</i> , 2005, 1, 15.	1.2	12
59	Needs for care among patients with schizophrenia in six European countries: a one-year follow-up study. <i>Clinical Practice and Epidemiology in Mental Health</i> , 2006, 2, 22.	1.2	12
60	The prevalence of late-life depression in a Portuguese community sample: A 10/66 Dementia Research Group study. <i>Journal of Affective Disorders</i> , 2019, 246, 674-681.	4.1	11
61	CANFOR Portuguese version: validation study. <i>BMC Psychiatry</i> , 2013, 13, 157.	2.6	10
62	Perceived helpfulness of treatment for generalized anxiety disorder: a World Mental Health Surveys report. <i>BMC Psychiatry</i> , 2021, 21, 392.	2.6	10
63	Suicidality in primary care patients who present with sadness and anhedonia: a prospective European study. <i>BMC Psychiatry</i> , 2016, 16, 94.	2.6	9
64	Obsessive-compulsive disorder and executive deficits in two patients with primary dystonia. <i>Parkinsonism and Related Disorders</i> , 2006, 12, 388-391.	2.2	8
65	Caregiving in severe mental illness: the psychometric properties of the Involvement Evaluation Questionnaire in Portugal. <i>Annals of General Psychiatry</i> , 2012, 11, 8.	2.7	8
66	The Marist's stigma scale: a standardized international measure of the stigma of schizophrenia and other psychoses. <i>BMC Psychiatry</i> , 2014, 14, 182.	2.6	8
67	How Many People Live with Dementia in Portugal? A Discussion Paper of National Estimates. <i>Portuguese Journal of Public Health</i> , 2021, 39, 58-68.	0.5	8
68	Trauma exposure and ptsd in portugal: Findings from the world mental health survey initiative. <i>Psychiatry Research</i> , 2020, 284, 112644.	3.3	7
69	How did the use of psychotropic drugs change during the Great Recession in Portugal? A follow-up to the National Mental Health Survey. <i>BMC Psychiatry</i> , 2020, 20, 215.	2.6	7
70	Perceived helpfulness of treatment for social anxiety disorder: findings from the WHO World Mental Health Surveys. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2022, 57, 2079-2095.	3.1	7
71	Sequential Combined Treatment of Heroin Addicted Patients in Portugal with Naltrexone and Family Therapy. <i>European Addiction Research</i> , 1997, 3, 138-145.	2.4	6
72	Munchausen syndrome mimicking psychiatric disease with concomitant genuine physical illness. <i>BMJ Case Reports</i> , 2010, 2010, bcr0720103142-bcr0720103142.	0.5	6

#	ARTICLE	IF	CITATIONS
73	Development and Validation of a Risk Model for Prediction of Hazardous Alcohol Consumption in General Practice Attendees: The PredictAL Study. PLoS ONE, 2011, 6, e22175.	2.5	6
74	Barriers to mental health services utilisation in Portugal – results from the National Mental Health Survey. Journal of Mental Health, 2022, 31, 453-461.	1.9	5
75	Implementação em Portugal de um estudo de prevalência da demência e da depressão geriátrica: a metodologia do 10/66 Dementia Research Group. Revista Portuguesa De Saude Publica, 2016, 34, 134-143.	0.3	4
76	Suicide time-series structural change analysis in Portugal (1913-2018): Impact of register bias on suicide trends. Journal of Affective Disorders, 2021, 291, 65-75.	4.1	4
77	Acute psychotic episode in a patient with thyrotoxicosis factitia. BMJ Case Reports, 2009, 2009, bcr0820080676-bcr0820080676.	0.5	4
78	Dealing with the COVID-19 pandemic in Europe: five lessons from the European Society for Traumatic Stress Studies. European Journal of Psychotraumatology, 2022, 13, 2046330.	2.5	4
79	Standardization of the MaristÅn Scale of Informal Care in people with schizophrenia and other psychoses. Acta Psychiatrica Scandinavica, 2013, 128, 468-474.	4.5	3
80	Multistage modeling for the classification of numerical and categorical datasets. , 2017, , .		3
81	Needs Assessment Framework for Mental Health Policy in Treating Persons With Serious Mental Illness in Azerbaijan. Psychiatric Services, 2019, 70, 432-435.	2.0	3
82	Standardised measures of needs, stigma and informal care in schizophrenia using a bottom-up, cross-cultural approach. Mental Health in Family Medicine, 2012, 9, 125-34.	0.2	3
83	Perceived helpfulness of service sectors used for mental and substance use disorders: Findings from the WHO World Mental Health Surveys. International Journal of Mental Health Systems, 2022, 16, 6.	2.7	3
84	Bipolar disorder, homocysteine and white matter hyperintensities. Bipolar Disorders, 2008, 10, 748-749.	1.9	2
85	Standardization of the MARISTÅN scale to measure needs in people with schizophrenia and related psychoses. International Journal of Social Psychiatry, 2014, 60, 219-226.	3.1	2
86	An Evaluation of the Therapeutic Programme Conducted by the Southern Regional Alcohol-Abuse Treatment Centre: Study on the Programme’s Results One Year after Discharge from Inpatient Care. European Addiction Research, 2001, 7, 61-68.	2.4	1
87	Transdiagnostic development of internalizing psychopathology throughout the life course up to age 45: a World Mental Health Surveys report. Psychological Medicine, 2020, , 1-10.	4.5	1
88	Childhood adversities and the comorbidity between mood and general medical disorders in adults: Results from the WHO World Mental Health Survey Portugal. Brain, Behavior, & Immunity - Health, 2021, 17, 100329.	2.5	1
89	Resistant schizophrenia in a patient with epilepsy. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2010, 34, 559-560.	4.8	0
90	Social Anxiety Disorder and Childhood adversities in Portugal: Findings from the World Mental Health Survey Initiative. Psychiatry Research, 2022, , 114734.	3.3	0