## Lars A Akslen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8479039/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Molecular portraits of human breast tumours. Nature, 2000, 406, 747-752.	27.8	13,397
2	Immunohistochemical and Clinical Characterization of the Basal-Like Subtype of Invasive Breast Carcinoma. Clinical Cancer Research, 2004, 10, 5367-5374.	7.0	2,393
3	Germline BRCA1 Mutations and a Basal Epithelial Phenotype in Breast Cancer. Journal of the National Cancer Institute, 2003, 95, 1482-1485.	6.3	807
4	Specific P53 mutations are associated with de novo resistance to doxorubicin in breast cancer patients. Nature Medicine, 1996, 2, 811-814.	30.7	797
5	Subtyping of Breast Cancer by Immunohistochemistry to Investigate a Relationship between Subtype and Short and Long Term Survival: A Collaborative Analysis of Data for 10,159 Cases from 12 Studies. PLoS Medicine, 2010, 7, e1000279.	8.4	764
6	EZH2 Expression Is Associated With High Proliferation Rate and Aggressive Tumor Subgroups in Cutaneous Melanoma and Cancers of the Endometrium, Prostate, and Breast. Journal of Clinical Oncology, 2006, 24, 268-273.	1.6	745
7	Landscape of genomic alterations in cervical carcinomas. Nature, 2014, 506, 371-375.	27.8	708
8	Axl is an essential epithelial-to-mesenchymal transition-induced regulator of breast cancer metastasis and patient survival. Proceedings of the National Academy of Sciences of the United States of America, 2010, 107, 1124-1129.	7.1	503
9	A Switch from E-Cadherin to N-Cadherin Expression Indicates Epithelial to Mesenchymal Transition and Is of Strong and Independent Importance for the Progress of Prostate Cancer. Clinical Cancer Research, 2007, 13, 7003-7011.	7.0	472
10	Role of Angiogenesis in Human Tumor Dormancy: Animal Models of the Angiogenic Switch. Cell Cycle, 2006, 5, 1779-1787.	2.6	450
11	Triple-Negative Breast Cancer: Distinguishing between Basal and Nonbasal Subtypes. Clinical Cancer Research, 2009, 15, 2302-2310.	7.0	422
12	A Model of Human Tumor Dormancy: An Angiogenic Switch From the Nonangiogenic Phenotype. Journal of the National Cancer Institute, 2006, 98, 316-325.	6.3	388
13	The Prognostic Implication of the Basal-Like (Cyclin) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 267 Td (Ehigh -Related Breast Cancer. Cancer Research, 2004, 64, 830-835.	/p27low/p 0.9	53+/Glome 345
14	Drug-sensitive <i>FGFR2</i> mutations in endometrial carcinoma. Proceedings of the National Academy of Sciences of the United States of America, 2008, 105, 8713-8717.	7.1	329
15	Use of immunohistochemical markers can refine prognosis in triple negative breast cancer. BMC Cancer, 2007, 7, 134.	2.6	316
16	Combined Vascular Endothelial Growth Factor Receptor and Epidermal Growth Factor Receptor (EGFR) Blockade Inhibits Tumor Growth in Xenograft Models of EGFR Inhibitor Resistance. Clinical Cancer Research, 2009, 15, 3484-3494.	7.0	297
17	PTEN methylation is associated with advanced stage and microsatellite instability in endometrial carcinoma. International Journal of Cancer, 2001, 91, 22-26.	5.1	270
18	Integrated Genome-Wide DNA Copy Number and Expression Analysis Identifies Distinct Mechanisms of Primary Chemoresistance in Ovarian Carcinomas. Clinical Cancer Research, 2009, 15, 1417-1427.	7.0	266

#	Article	IF	CITATIONS
19	Expression of enhancer of zeste homologue 2 is significantly associated with increased tumor cell proliferation and is a marker of aggressive breast cancer Clinical Cancer Research, 2006, 12, 1168-1174.	7.0	249
20	Genome-wide association study identifies three new melanoma susceptibility loci. Nature Genetics, 2011, 43, 1108-1113.	21.4	230
21	Genome-wide meta-analysis identifies five new susceptibility loci for cutaneous malignant melanoma. Nature Genetics, 2015, 47, 987-995.	21.4	218
22	Prognostic significance of histologic grading compared with subclassification of papillary thyroid carcinoma. Cancer, 2000, 88, 1902-1908.	4.1	208
23	A Basal Epithelial Phenotype Is More Frequent in Interval Breast Cancers Compared with Screen Detected Tumors. Cancer Epidemiology Biomarkers and Prevention, 2005, 14, 1108-1112.	2.5	185
24	Importance of P-Cadherin, β-Catenin, and Wnt5a/Frizzled for Progression of Melanocytic Tumors and Prognosis in Cutaneous Melanoma. Clinical Cancer Research, 2005, 11, 8606-8614.	7.0	179
25	Placental Cadherin and the Basal Epithelial Phenotype of BRCA1-Related Breast Cancer. Clinical Cancer Research, 2005, 11, 4003-4011.	7.0	157
26	Tumorâ€vascular interactions and tumor dormancy. Apmis, 2008, 116, 569-585.	2.0	152
27	Prognostic Significance of Angiogenesis and <i>Ki-67</i> , <i>p53</i> , and <i>p21</i> Expression: A Population-Based Endometrial Carcinoma Study. Journal of Clinical Oncology, 1999, 17, 1382-1382.	1.6	149
28	Significance of tumour-associated macrophages, vascular endothelial growth factor and thrombospondin-1 expression for tumour angiogenesis and prognosis in endometrial carcinomas. International Journal of Cancer, 1999, 84, 538-543.	5.1	146
29	Prognostic importance of histologic grading in papillary thyroid carcinoma. Cancer, 1993, 72, 2680-2685.	4.1	139
30	Prognostic Impact of Alterations in P-Cadherin Expression and Related Cell Adhesion Markers in Endometrial Cancer. Journal of Clinical Oncology, 2004, 22, 1242-1252.	1.6	138
31	Genome-wide association meta-analyses combining multiple risk phenotypes provide insights into the genetic architecture of cutaneous melanoma susceptibility. Nature Genetics, 2020, 52, 494-504.	21.4	138
32	Genome-Wide Profiling of Histone H3 Lysine 4 and Lysine 27 Trimethylation Reveals an Epigenetic Signature in Prostate Carcinogenesis. PLoS ONE, 2009, 4, e4687.	2.5	130
33	Combined loss of PTEN and p27 expression is associated with tumor cell proliferation by Ki-67 and increased risk of recurrent disease in localized prostate cancer. Clinical Cancer Research, 2003, 9, 1474-9.	7.0	124
34	Hormone receptor loss in endometrial carcinoma curettage predicts lymph node metastasis and poor outcome in prospective multicentre trial. European Journal of Cancer, 2013, 49, 3431-3441.	2.8	123
35	Lack of Estrogen Receptor-α Is Associated with Epithelial–Mesenchymal Transition and PI3K Alterations in Endometrial Carcinoma. Clinical Cancer Research, 2013, 19, 1094-1105.	7.0	120
36	Vascular Proliferation Is Important for Clinical Progress of Endometrial Cancer. Cancer Research, 2006, 66, 3303-3309.	0.9	116

#	Article	IF	CITATIONS
37	Gene expression profiles do not consistently predict the clinical treatment response in locally advanced breast cancer. Molecular Cancer Therapeutics, 2006, 5, 2914-2918.	4.1	114
38	Suppression of heat shock protein 27 induces long-term dormancy in human breast cancer. Proceedings of the National Academy of Sciences of the United States of America, 2012, 109, 8699-8704.	7.1	114
39	Stathmin Overexpression Identifies High-Risk Patients and Lymph Node Metastasis in Endometrial Cancer. Clinical Cancer Research, 2011, 17, 3368-3377.	7.0	113
40	Bone Marrow–Derived Gr1+ Cells Can Generate a Metastasis-Resistant Microenvironment Via Induced Secretion of Thrombospondin-1. Cancer Discovery, 2013, 3, 578-589.	9.4	113
41	Gene Expression in Poorly Differentiated Papillary Thyroid Carcinomas. Thyroid, 2006, 16, 161-175.	4.5	112
42	Melanoma prone families with <i>CDK4</i> germline mutation: phenotypic profile and associations with <i>MC1R</i> variants. Journal of Medical Genetics, 2013, 50, 264-270.	3.2	112
43	Tumor necrosis is an important hallmark of aggressive endometrial cancer and associates with hypoxia, angiogenesis and inflammation responses. Oncotarget, 2015, 6, 39676-39691.	1.8	112
44	A variant in FTO shows association with melanoma risk not due to BMI. Nature Genetics, 2013, 45, 428-432.	21.4	111
45	BRAF and NRAS Mutations Are Frequent in Nodular Melanoma but Are not Associated with Tumor Cell Proliferation or Patient Survival. Journal of Investigative Dermatology, 2005, 125, 312-317.	0.7	109
46	Early detection of breast cancer based on gene-expression patterns in peripheral blood cells. Breast Cancer Research, 2005, 7, R634-44.	5.0	109
47	Prolonged dormancy of human liposarcoma is associated with impaired tumor angiogenesis. FASEB Journal, 2006, 20, 947-949.	0.5	109
48	The Effect on Melanoma Risk of Genes Previously Associated With Telomere Length. Journal of the National Cancer Institute, 2014, 106, .	6.3	109
49	Ki-67 expression is superior to mitotic count and novel proliferation markers PHH3, MCM4 and mitosin as a prognostic factor in thick cutaneous melanoma. BMC Cancer, 2010, 10, 140.	2.6	104
50	Alterations and prognostic significance of p16 and p53 protein expression in subgroups of cutaneous melanoma. , 1997, 74, 535-539.		99
51	A large Norwegian family with inherited malignant melanoma, multiple atypical nevi, andCDK4 mutation. Genes Chromosomes and Cancer, 2005, 44, 10-18.	2.8	94
52	A single nucleotide polymorphism in the 3?untranslated region of theCDKN2A gene is common in sporadic primary melanomas but mutations in theCDKN2B,CDKN2C,CDK4 andp53 genes are rare. International Journal of Cancer, 2001, 95, 388-393.	5.1	90
53	Proliferation of Immature Tumor Vessels Is a Novel Marker of Clinical Progression in Prostate Cancer. Cancer Research, 2009, 69, 4708-4715.	0.9	90
54	Prognostic importance of various clinicopathological features in papillary thyroid carcinoma. European Journal of Cancer, 1993, 29, 44-51.	2.8	85

#	Article	IF	CITATIONS
55	Prosaposin inhibits tumor metastasis via paracrine and endocrine stimulation of stromal p53 and Tsp-1. Proceedings of the National Academy of Sciences of the United States of America, 2009, 106, 12115-12120.	7.1	85
56	Prognostic importance of glomeruloid microvascular proliferation indicates an aggressive angiogenic phenotype in human cancers. Cancer Research, 2002, 62, 6808-11.	0.9	81
57	Loss of PTEN expression is associated with metastatic disease in patients with endometrial carcinoma. Cancer, 2002, 94, 2185-2191.	4.1	80
58	Oncoproteins and tumor progression in papillary thyroid carcinoma. Presence of epidermal growth factor receptor, c-erbB-2 protein, estrogen receptor related protein, p21-ras protein, and proliferation indicators in relation to tumor recurrences and patient survival. Cancer, 1995, 76, 1643-1654.	4.1	79
59	Biologic markers in endometrial cancer treatment. Apmis, 2009, 117, 693-707.	2.0	76
60	Thyroid Carcinoma with Mixed Tall-Cell and Columnar-Cell Features. American Journal of Clinical Pathology, 1990, 94, 442-445.	0.7	75
61	Mutation analysis of the EGFR–NRAS–BRAF pathway in melanomas from black Africans and other subgroups of cutaneous melanoma. Melanoma Research, 2008, 18, 29-35.	1.2	75
62	Two-view digital breast tomosynthesis versus digital mammography in a population-based breast cancer screening programme (To-Be): a randomised, controlled trial. Lancet Oncology, The, 2019, 20, 795-805.	10.7	75
63	GATA3 expression in estrogen receptor α-negative endometrial carcinomas identifies aggressive tumors with high proliferation and poor patient survival. American Journal of Obstetrics and Gynecology, 2008, 199, 543.e1-543.e7.	1.3	73
64	Molecular pathogenesis and prognostic factors in endometrial carcinoma. Apmis, 2002, 110, 673-689.	2.0	71
65	Integrated Genomic Analysis of the 8q24 Amplification in Endometrial Cancers Identifies ATAD2 as Essential to MYC-Dependent Cancers. PLoS ONE, 2013, 8, e54873.	2.5	70
66	High-throughput interrogation of PIK3CA, PTEN, KRAS, FBXW7 and TP53 mutations in primary endometrial carcinoma. Gynecologic Oncology, 2013, 128, 327-334.	1.4	68
67	<i>TMPRSS2:ERG</i> fusion transcripts in urine from prostate cancer patients correlate with a less favorable prognosis. Apmis, 2009, 117, 575-582.	2.0	66
68	AXL Targeting Abrogates Autophagic Flux and Induces Immunogenic Cell Death in Drug-Resistant Cancer Cells. Journal of Thoracic Oncology, 2020, 15, 973-999.	1.1	66
69	Risk of primary childhood brain tumors related to birth characteristics: A Norwegian prospective study. International Journal of Cancer, 1998, 77, 498-503.	5.1	65
70	Molecular determinants of outcome with mammalian target of rapamycin inhibition in endometrial cancer. Cancer, 2014, 120, 603-610.	4.1	64
71	Expression of p53 protein in cutaneous melanoma. International Journal of Cancer, 1992, 52, 13-16.	5.1	63
72	ARID1A loss is prevalent in endometrial hyperplasia with atypia and low-grade endometrioid carcinomas. Modern Pathology, 2013, 26, 428-434.	5.5	61

#	Article	IF	CITATIONS
73	Independent prognostic importance of vascular invasion in nodular melanomas. , 1996, 78, 1211-1219.		60
74	Pathologic expression of p53 or p16 in preoperative curettage specimens identifies high-risk endometrial carcinomas. American Journal of Obstetrics and Gynecology, 2006, 195, 979-986.	1.3	60
75	Incidence of thyroid cancer in Norway 1970–1985. Apmis, 1990, 98, 549-558.	2.0	58
76	Increased angiogenesis in papillarythyroid carcinoma but lack of prognostic importance. Human Pathology, 2000, 31, 439-442.	2.0	58
77	Prognostic Importance of the Mitotic Marker Phosphohistone H3 in Cutaneous Nodular Melanoma. Journal of Investigative Dermatology, 2012, 132, 1247-1252.	0.7	56
78	BRAF-V600E expression in primary nodular melanoma is associated with aggressive tumour features and reduced survival. British Journal of Cancer, 2016, 114, 801-808.	6.4	56
79	Increased Expression of VEGF-Receptors (FLT-1, KDR, NRP-1) and Thrombospondin-1 is Associated with Glomeruloid Microvascular Proliferation, an Aggressive Angiogenic Phenotype, in Malignant Melanoma. Angiogenesis, 2003, 6, 295-301.	7.2	55
80	Loss of BMI-1 expression is associated with clinical progress of malignant melanoma. Modern Pathology, 2008, 21, 583-590.	5.5	55
81	Incidence pattern of thyroid cancer in norway: Influence of birth cohort and time period. International Journal of Cancer, 1993, 53, 183-187.	5.1	54
82	Evaluation of Ki67 Expression across Distinct Categories of Breast Cancer Specimens: A Population-Based Study of Matched Surgical Specimens, Core Needle Biopsies and Tissue Microarrays. PLoS ONE, 2014, 9, e112121.	2.5	54
83	Development of a prosaposin-derived therapeutic cyclic peptide that targets ovarian cancer via the tumor microenvironment. Science Translational Medicine, 2016, 8, 329ra34.	12.4	54
84	Tumor-associated macrophages are strongly related to vascular invasion, non-luminal subtypes, and interval breast cancer. Human Pathology, 2017, 69, 72-80.	2.0	53
85	Epithelial to Mesenchymal Transition of a Primary Prostate Cell Line with Switches of Cell Adhesion Modules but without Malignant Transformation. PLoS ONE, 2008, 3, e3368.	2.5	52
86	Prognostic impact and concordance of TERT promoter mutation and protein expression in matched primary and metastatic cutaneous melanoma. British Journal of Cancer, 2018, 118, 98-105.	6.4	52
87	Impact of pre-diagnostic triglycerides and HDL-cholesterol on breast cancer recurrence and survival by breast cancer subtypes. BMC Cancer, 2018, 18, 654.	2.6	52
88	Significance of CD 105 expression for tumour angiogenesis and prognosis in endometrial carcinomas. Apmis, 2003, 111, 1011-1018.	2.0	51
89	Poorly Differentiated Thyroid Carcinoma—It Is Important. American Journal of Surgical Pathology, 2000, 24, 310.	3.7	50
90	NATH, a novel gene overexpressed in papillary thyroid carcinomas. Oncogene, 2002, 21, 5056-5068.	5.9	49

#	Article	IF	CITATIONS
91	Secondary breast cancer: a 5â€year populationâ€based study with review of the literature. Apmis, 2009, 117, 762-767.	2.0	49
92	Efficient in vivo vascularization of tissue-engineering scaffolds. Journal of Tissue Engineering and Regenerative Medicine, 2011, 5, e52-e62.	2.7	49
93	Vascular proliferation is increased in basal-like breast cancer. Breast Cancer Research and Treatment, 2011, 130, 1063-1071.	2.5	48
94	MSI-low, a real phenomenon which varies in frequency among cancer types. Journal of Pathology, 2003, 201, 389-394.	4.5	47
95	Prognostic significance of p16 and CDK4 proteins in localized prostate carcinoma. , 2000, 88, 416-424.		46
96	Low frequency ofBRAFandCDKN2Amutations in endometrial cancer. International Journal of Cancer, 2005, 115, 930-934.	5.1	46
97	Clinical Efficacy and Safety of Bevacizumab Monotherapy in Patients with Metastatic Melanoma: Predictive Importance of Induced Early Hypertension. PLoS ONE, 2012, 7, e38364.	2.5	46
98	Populationâ€based prevalence of <i>CDKN2A</i> and <i>CDK4</i> mutations in patients with multiple primary melanomas. Genes Chromosomes and Cancer, 2008, 47, 175-184.	2.8	44
99	Tumor necrosis is associated with increased alphavbeta3integrin expression and poor prognosis in nodular cutaneous melanomas. BMC Cancer, 2008, 8, 362.	2.6	44
100	Lipocalin 2 expression is associated with aggressive features of endometrial cancer. BMC Cancer, 2012, 12, 169.	2.6	44
101	Gene Expression Patterns Related to Vascular Invasion and Aggressive Features in Endometrial Cancer. American Journal of Pathology, 2011, 178, 861-871.	3.8	42
102	Importance of tumour cell invasion in blood and lymphatic vasculature among patients with endometrial carcinoma. Histopathology, 2009, 54, 174-183.	2.9	41
103	Expression and alternative splicing of c-ret RNA in papillary thyroid carcinomas. Oncogene, 2001, 20, 885-892.	5.9	39
104	Expression of bFGF/FGFR-1 and vascular proliferation related to clinicopathologic features and tumor progress in localized prostate cancer. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2006, 448, 68-74.	2.8	39
105	GRK3 is essential for metastatic cells and promotes prostate tumor progression. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, 1521-1526.	7.1	39
106	Strong Expression of Hypoxia-Inducible Factor-1α (HIF-1α) Is Associated with Axl Expression and Features of Aggressive Tumors in African Breast Cancer. PLoS ONE, 2016, 11, e0146823.	2.5	39
107	Loss of hMSH2 and hMSH6 expression is frequent in sporadic endometrial carcinomas with microsatellite instability: a population-based study. Clinical Cancer Research, 2002, 8, 138-43.	7.0	39
108	Frequent mutations of the p53 gene in cutaneous melanoma of the nodular type. , 1998, 79, 91-95.		37

#	Article	IF	CITATIONS
109	Frequency of the basalâ€like phenotype in African breast cancer. Apmis, 2007, 115, 1391-1399.	2.0	36
110	Endothelial microvascular networks affect gene-expression profiles and osteogenic potential of tissue-engineered constructs. Stem Cell Research and Therapy, 2013, 4, 52.	5.5	36
111	Dynamic contrast-enhanced MRI in endometrial carcinoma identifies patients at increased risk of recurrence. European Radiology, 2013, 23, 2916-2925.	4.5	36
112	Loss of p63 and cytokeratin 5/6 expression is associated with more aggressive tumors in endometrial carcinoma patients. International Journal of Cancer, 2006, 118, 1227-1233.	5.1	35
113	High Phospho-Stathmin(Serine38) Expression Identifies Aggressive Endometrial Cancer and Suggests an Association with PI3K Inhibition. Clinical Cancer Research, 2013, 19, 2331-2341.	7.0	35
114	Automated Volumetric Analysis of Mammographic Density in a Screening Setting: Worse Outcomes for Women with Dense Breasts. Radiology, 2018, 288, 343-352.	7.3	35
115	Outcome after surgery for primary hyperaldosteronism may depend on KCNJ5 tumor mutation status: a population-based study from Western Norway. Langenbeck's Archives of Surgery, 2013, 398, 869-874.	1.9	34
116	Lowâ€molecular contrast agent dynamic contrastâ€enhanced (DCE)â€MRI and diffusionâ€weighted (DW)â€MRI i early assessment of bevacizumab treatment in breast cancer xenografts. Journal of Magnetic Resonance Imaging, 2013, 38, 1043-1053.	in 3.4	34
117	Stathmin Protein Level, a Potential Predictive Marker for Taxane Treatment Response in Endometrial Cancer. PLoS ONE, 2014, 9, e90141.	2.5	34
118	Vascular proliferation is a prognostic factor in breast cancer. Breast Cancer Research and Treatment, 2012, 133, 501-510.	2.5	33
119	1.5-T multiparametric MRI using PI-RADS: a region by region analysis to localize the index-tumor of prostate cancer in patients undergoing prostatectomy. Acta Radiologica, 2015, 56, 500-511.	1.1	33
120	Expression of EGFR and câ€kit is associated with the basalâ€like phenotype in breast carcinomas of African women. Apmis, 2008, 116, 515-525.	2.0	32
121	High mortality due to cutaneous melanoma in Norway: a study of prognostic factors in a nationwide cancer registry. Clinical Epidemiology, 2018, Volume 10, 537-548.	3.0	32
122	Poorer survival of nulliparous women with endometrial carcinoma. , 1998, 82, 1328-1333.		30
123	Fine mapping of genetic susceptibility loci for melanoma reveals a mixture of single variant and multiple variant regions. International Journal of Cancer, 2015, 136, 1351-1360.	5.1	30
124	Expression of heregulins and associations with the ErbB family of tyrosine kinase receptors in papillary thyroid carcinomas. International Journal of Cancer, 2000, 87, 763-770.	5.1	29
125	A registry-based study of thyroid paraganglioma: histological and genetic characteristics. Endocrine-Related Cancer, 2015, 22, 191-204.	3.1	29
126	In vivo MRI and histopathological assessment of tumor microenvironment in luminalâ€like and basalâ€like breast cancer xenografts. Journal of Magnetic Resonance Imaging, 2012, 35, 1098-1107.	3.4	27

#	Article	IF	CITATIONS
127	Deoxyribonucleic acid ploidy in endometrial carcinoma: a reproducible and valid prognostic marker in a routine diagnostic setting. American Journal of Obstetrics and Gynecology, 2009, 201, 603.e1-603.e7.	1.3	25
128	QSOX1 expression is associated with aggressive tumor features and reduced survival in breast carcinomas. Modern Pathology, 2016, 29, 1485-1491.	5.5	25
129	FOXC2 expression and epithelial–mesenchymal phenotypes are associated with castration resistance, metastasis and survival in prostate cancer. Journal of Pathology: Clinical Research, 2019, 5, 272-286.	3.0	25
130	Exploring the effects of lifestyle on breast cancer risk, age at diagnosis, and survival: the EBBA-Life study. Breast Cancer Research and Treatment, 2020, 182, 215-227.	2.5	25
131	Post-PET ultrasound improves specificity of 18F-FDG-PET for recurrent differentiated thyroid cancer while maintaining sensitivity. Acta Radiologica, 2015, 56, 1350-1360.	1.1	24
132	Estrogen receptor β– an independent prognostic marker in estrogen receptor α and progesterone receptorâ€positive breast cancer?. Apmis, 2009, 117, 644-650.	2.0	23
133	Concomitant inactivation of the p53―and pRB―functional pathways predicts resistance to DNA damaging drugs in breast cancer inÂvivo. Molecular Oncology, 2015, 9, 1553-1564.	4.6	23
134	Microvascular proliferation in luminal A and basal-like breast cancer subtypes. Journal of Clinical Pathology, 2015, 68, 891-897.	2.0	23
135	Prognostic impact of morphometric nuclear grade of endometrial carcinoma. Cancer, 1998, 83, 956-964.	4.1	22
136	Microenvironment-Induced Non-sporadic Expression of the AXL and cKIT Receptors Are Related to Epithelial Plasticity and Drug Resistance. Frontiers in Cell and Developmental Biology, 2018, 6, 41.	3.7	22
137	Switch in FOXA1 Status Associates with Endometrial Cancer Progression. PLoS ONE, 2014, 9, e98069.	2.5	22
138	<i>MC1R, ASIP, TYR,</i> and <i>TYRP1</i> gene variants in a populationâ€based series of multiple primary melanomas. Genes Chromosomes and Cancer, 2012, 51, 654-661.	2.8	21
139	Expression of thyroid transcription factor-1 is associated with a basal-like phenotype in breast carcinomas. Diagnostic Pathology, 2013, 8, 80.	2.0	21
140	Breast cancer stromal elastosis is associated with mammography screening detection, low Ki67 expression and favourable prognosis in a population-based study. Diagnostic Pathology, 2014, 9, 230.	2.0	21
141	Interval and Subsequent Round Breast Cancer in a Randomized Controlled Trial Comparing Digital Breast Tomosynthesis and Digital Mammography Screening. Radiology, 2021, 300, 66-76.	7.3	21
142	Cytoplasmic localization of EGF receptor in papillary thyroid carcinomas: Association with the 150-kDa receptor form. , 1996, 65, 161-167.		20
143	Prognostic value of uPAR expression and angiogenesis in primary and metastatic melanoma. PLoS ONE, 2019, 14, e0210399.	2.5	20
144	AXL Is a Driver of Stemness in Normal Mammary Gland and Breast Cancer. IScience, 2020, 23, 101649.	4.1	20

#	Article	IF	CITATIONS
145	The epithelial–mesenchymal transition regulators Twist, Slug, and Snail are associated with aggressive tumour features and poor outcome in prostate cancer patients. Journal of Pathology: Clinical Research, 2021, 7, 253-270.	3.0	20
146	Effect of antiangiogenic therapy on tumor growth, vasculature and kinase activity in basal―and luminalâ€like breast cancer xenografts. Molecular Oncology, 2012, 6, 418-427.	4.6	19
147	Spontaneous Reversion of the Angiogenic Phenotype to a Nonangiogenic and Dormant State in Human Tumors. Molecular Cancer Research, 2014, 12, 754-764.	3.4	19
148	Expression of Nestin associates with BRCA1 mutations, a basal-like phenotype and aggressive breast cancer. Scientific Reports, 2017, 7, 1089.	3.3	19
149	Angiogenesis does not predict clinical response to doxorubicin monotherapy in patients with locally advanced breast cancer. International Journal of Cancer, 1997, 74, 138-140.	5.1	18
150	ls preoperative serum prostate-specific antigen level significantly related to clinical recurrence after radical retropubic prostatectomy for localized prostate cancer?. BJU International, 2006, 97, 51-55.	2.5	18
151	Inverse correlation between PDGFC expression and lymphocyte infiltration in human papillary thyroid carcinomas. BMC Cancer, 2009, 9, 425.	2.6	18
152	Tumour cell invasion into blood vessels is significantly related to breast cancer subtypes and decreased survival. Journal of Clinical Pathology, 2017, 70, 313-319.	2.0	18
153	Optimising preoperative risk stratification tools for prostate cancer using mpMRI. European Radiology, 2018, 28, 1016-1026.	4.5	18
154	Asparaginase-like protein 1 is an independent prognostic marker in primary endometrial cancer, and is frequently lost in metastatic lesions. Gynecologic Oncology, 2018, 148, 197-203.	1.4	18
155	Integrin α11β1 is expressed in breast cancer stroma and associates with aggressive tumor phenotypes. Journal of Pathology: Clinical Research, 2020, 6, 69-82.	3.0	18
156	Stathmin expression associates with vascular and immune responses in aggressive breast cancer subgroups. Scientific Reports, 2020, 10, 2914.	3.3	18
157	Vascular proliferation is associated with survival in pancreatic ductal adenocarcinoma. Apmis, 2013, 121, 1037-1046.	2.0	17
158	Vascular invasion is an adverse prognostic factor in resected non–smallâ€cell lung cancer. Apmis, 2017, 125, 197-206.	2.0	17
159	Assessing Extraprostatic Extension with Multiparametric MRI of the Prostate: Mehralivand Extraprostatic Extension Grade or Extraprostatic Extension Likert Scale?. Radiology Imaging Cancer, 2020, 2, e190071.	1.6	17
160	Human Organotypic Airway and Lung Organoid Cells of Bronchiolar and Alveolar Differentiation Are Permissive to Infection by Influenza and SARS-CoV-2 Respiratory Virus. Frontiers in Cellular and Infection Microbiology, 2022, 12, 841447.	3.9	17
161	Balloon Cell Melanoma Mimicking Clear Cell Carcinoma. Pathology Research and Practice, 1989, 184, 548-550.	2.3	16
162	Molecular assessment of depth of myometrial invasion in stage I endometrial cancer: a model based on K-ras mutation analysis. Gynecologic Oncology, 2003, 91, 218-225.	1.4	16

#	Article	IF	CITATIONS
163	WDR19 Expression is Increased in Prostate Cancer Compared with Normal Cells, but Low-Intensity Expression in Cancers is Associated with Shorter Time to Biochemical Failures and Local Recurrence. Clinical Cancer Research, 2008, 14, 1397-1406.	7.0	16
164	Identification of a CDK4 R24H mutation-positive melanoma family by analysis of early-onset melanoma patients in Latvia. Melanoma Research, 2009, 19, 119-122.	1.2	16
165	Stathmin is superior to AKT and phosphoâ€AKT staining for the detection of phosphoinositide 3â€kinase activation and aggressive endometrial cancer. Histopathology, 2010, 57, 641-646.	2.9	16
166	Reprogramming of cell junction modules during stepwise epithelial to mesenchymal transition and accumulation of malignant features in vitro in a prostate cell model. Experimental Cell Research, 2011, 317, 234-247.	2.6	16
167	Polymorphisms in hormone metabolism and growth factor genes and mammographic density in Norwegian postmenopausal hormone therapy users and non-users. Breast Cancer Research, 2012, 14, R135.	5.0	16
168	An 18-Gene Signature for Vascular Invasion Is Associated with Aggressive Features and Reduced Survival in Breast Cancer. PLoS ONE, 2014, 9, e98787.	2.5	16
169	A positive real-time elastography is an independent marker for detection of high-risk prostate cancers in the primary biopsy setting. BJU International, 2014, 113, E90-E97.	2.5	16
170	Extra-nodal extension is a significant prognostic factor in lymph node positive breast cancer. PLoS ONE, 2017, 12, e0171853.	2.5	16
171	Tumour cell expression of interleukin 6 receptor α is associated with response rates in patients treated with sunitinib for metastatic clear cell renal cell carcinoma. Journal of Pathology: Clinical Research, 2018, 4, 114-123.	3.0	15
172	Influence of pre-operative oral carbohydrate loading vs. standard fasting on tumor proliferation and clinical outcome in breast cancer patients ─ a randomized trial. BMC Cancer, 2019, 19, 1076.	2.6	15
173	Expression of Heat Shock Protein 27 in Melanoma Metastases Is Associated with Overall Response to Bevacizumab Monotherapy: Analyses of Predictive Markers in a Clinical Phase II Study. PLoS ONE, 2016, 11, e0155242.	2.5	15
174	Increased angiogenesis is associated with a 32-gene expression signature and 6p21 amplification in aggressive endometrial cancer. Oncotarget, 2015, 6, 10634-10645.	1.8	15
175	Hyperthermia improves the antitumour effect of metronomic cyclophosphamide in a rat transplantable brain tumour. Radiotherapy and Oncology, 2008, 86, 435-442.	0.6	14
176	Receptor for Activated Protein C Kinase 1 (RACK1) Is Overexpressed in Papillary Thyroid Carcinoma. Thyroid, 2011, 21, 1217-1225.	4.5	14
177	Gene expression in tumor cells and stroma in dsRed 4T1 tumors in eGFP-expressing mice with and without enhanced oxygenation. BMC Cancer, 2012, 12, 21.	2.6	14
178	Neuropilin 1 expression correlates with differentiation status of epidermal cells and cutaneous squamous cell carcinomas. Laboratory Investigation, 2014, 94, 752-765.	3.7	14
179	Nestin expression is associated with aggressive cutaneous melanoma of the nodular type. Modern Pathology, 2014, 27, 396-401.	5.5	14
180	Expression of glucocorticoid receptor is associated with aggressive primary endometrial cancer and increases from primary to metastatic lesions. Gynecologic Oncology, 2017, 147, 672-677.	1.4	14

#	Article	IF	CITATIONS
181	CD24-targeted fluorescence imaging in patient-derived xenograft models of high-grade serous ovarian carcinoma. EBioMedicine, 2020, 56, 102782.	6.1	14
182	Baseline microvessel density predicts response to neoadjuvant bevacizumab treatment of locally advanced breast cancer. Scientific Reports, 2021, 11, 3388.	3.3	14
183	Altered expression of cell cycle regulators Cyclin D1, p14, p16, CDK4 and Rb in nodular melanomas. International Journal of Oncology, 2004, 25, 1559-65.	3.3	14
184	Analysis of Latvian familial melanoma patients shows novel variants in the noncoding regions of CDKN2A and that the CDK4 mutation R24H is a founder mutation. Melanoma Research, 2013, 23, 221-226.	1.2	13
185	Microvascular proliferation is associated with aggressive tumour features and reduced survival in lung adenocarcinoma. Journal of Pathology: Clinical Research, 2017, 3, 249-257.	3.0	13
186	HER-2/neu expression is associated with high tumor cell proliferation and aggressive phenotype in a population based patient series of endometrial carcinomas. International Journal of Oncology, 2008, 32, 307-16.	3.3	13
187	Thyroid cancer in children in Norway 1953–1987. European Journal of Cancer, 1993, 29, 365-366.	2.8	12
188	Variation in Inflammatory Cytokine/Growth-Factor Genes and Mammographic Density in Premenopausal Women Aged 50–55. PLoS ONE, 2013, 8, e65313.	2.5	12
189	A positive Real-Time Elastography (RTE) combined with a Prostate Cancer Gene 3 (PCA3) score above 35 convey a high probability of intermediate- or high-risk prostate cancer in patient admitted for primary prostate biopsy. BMC Urology, 2016, 16, 39.	1.4	12
190	Deblender: a semiâ^'/unsupervised multi-operational computational method for complete deconvolution of expression data from heterogeneous samples. BMC Bioinformatics, 2018, 19, 408.	2.6	12
191	Expression of p 16 protein in prostatic adenocarcinomas, intraepithelial neoplasia, and benign/hyperplastic glands. Urologic Oncology: Seminars and Original Investigations, 1997, 3, 59-66.	1.6	11
192	Can We Rely on Pathologic Parameters to Define Conservative Treatment of Papillary Thyroid Carcinoma?. International Journal of Surgical Pathology, 2002, 10, 267-272.	0.8	11
193	Increased tumor cell expression of Axl is a marker of aggressive features in breast cancer among African women. Apmis, 2015, 123, 688-696.	2.0	11
194	Tumor dormancy-from basic mechanisms to clinical practice. Apmis, 2008, 116, 545-547.	2.0	10
195	Plasma calprotectin concentrations in women with endometrial carcinoma. Gynecologic Oncology, 2009, 114, 491-495.	1.4	10
196	Combination of real-time elastography and urine prostate cancer gene 3 (PCA3) detects more than 97% of significant prostate cancers. Scandinavian Journal of Urology, 2013, 47, 211-216.	1.0	10
197	Predictive value of C-reactive protein in patients treated with sunitinib for metastatic clear cell renal cell carcinoma. BMC Urology, 2017, 17, 74.	1.4	10
198	Low expression of SerpinB2 is associated with reduced survival in lung adenocarcinomas. Oncotarget, 2017, 8, 90706-90718.	1.8	10

#	Article	IF	CITATIONS
199	Malignt melanom – diagnostikk, behandling og oppfÃ,lging i Norge. Tidsskrift for Den Norske Laegeforening, 2013, 133, 2154-2158.	0.2	10
200	Quality of life among women with symptomatic, screen-detected, and interval breast cancer, and for women without breast cancer: a retrospective cross-sectional study from Norway. Quality of Life Research, 2022, 31, 1057-1068.	3.1	10
201	Quantifying tumour vascularity in non-luminal breast cancers. Journal of Clinical Pathology, 2017, 70, 766-774.	2.0	9
202	Characterization of FGD5 Expression in Primary Breast Cancers and Lymph Node Metastases. Journal of Histochemistry and Cytochemistry, 2018, 66, 787-799.	2.5	9
203	Metabolic consequences of perioperative oral carbohydrates in breast cancer patients $\hat{a} \in$ " an explorative study. BMC Cancer, 2019, 19, 1183.	2.6	9
204	Evaluation of Tumor Cell Proliferation by Ki-67 Expression and Mitotic Count in Lymph Node Metastases from Breast Cancer. PLoS ONE, 2016, 11, e0150979.	2.5	9
205	Mammographic density and histopathologic characteristics of screen-detected tumors in the Norwegian Breast Cancer Screening Program. Acta Radiologica Open, 2015, 4, 205846011560434.	0.6	8
206	Prostate cancer antigen-3 (PCA3) and PCA3-based nomograms in the diagnosis of prostate cancer: an external validation of Hansen's nomogram on a Norwegian cohort. Scandinavian Journal of Urology, 2015, 49, 8-15.	1.0	8
207	High PDGFRb Expression Predicts Resistance to Radiotherapy in DCIS within the SweDCIS Randomized Trial. Clinical Cancer Research, 2021, 27, 3469-3477.	7.0	8
208	P53 immunostaining in melanocytic lesions. European Journal of Cancer, 1993, 29, 652-653.	2.8	7
209	Predictive value of angiogenic proteins in patients with metastatic melanoma treated with bevacizumab monotherapy. Journal of Pathology: Clinical Research, 2019, 5, 53-62.	3.0	7
210	Preoperative imaging markers and PDZ-binding kinase tissue expression predict low-risk disease in endometrial hyperplasias and low grade cancers. Oncotarget, 2017, 8, 68530-68541.	1.8	7
211	Intrinsic Differences in Spatiotemporal Organization and Stromal Cell Interactions Between Isogenic Lung Cancer Cells of Epithelial and Mesenchymal Phenotypes Revealed by High-Dimensional Single-Cell Analysis of Heterotypic 3D Spheroid Models. Frontiers in Oncology, 2022, 12, 818437.	2.8	7
212	Fetal Heart Block, Anti–Ssa and Anti–Ssb Antibodies Association with intra-uterine growth retardation, fetal death and lupus anticoagulant. Acta Obstetricia Et Gynecologica Scandinavica, 1987, 66, 737-739.	2.8	6
213	Tumorâ€associated lymphocytes and macrophages are related to stromal elastosis and vascular invasion in breast cancer. Journal of Pathology: Clinical Research, 2021, 7, 517-527.	3.0	6
214	Frequency of somaticBRAFmutations in melanocytic lesions from patients in aCDK4melanoma family. Pigment Cell and Melanoma Research, 2014, 27, 149-151.	3.3	5
215	Terminal digit preference: a source of measurement error in breast cancer diameter reporting. Acta Oncológica, 2020, 59, 260-267.	1.8	5
216	Aneuploidy related transcriptional changes in endometrial cancer link low expression of chromosome 15q genes to poor survival. Oncotarget, 2017, 8, 9696-9707.	1.8	4

#	Article	IF	CITATIONS
217	Squamous Cell Carcinoma of the Urinary Bladder. Scandinavian Journal of Urology and Nephrology, 1991, 25, 275-276.	1.4	3
218	Enhanced expression of EGF receptor and low frequency of ras mutations in X-ray-induced rat thyroid tumours. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 1999, 435, 434-441.	2.8	3
219	Akt1 Activity Regulates Vessel Maturation in a Tissue Engineering Model of Angiogenesis. Tissue Engineering - Part A, 2014, 20, 2590-2603.	3.1	3
220	HSP27 Expression as a Novel Predictive Biomarker for Bevacizumab: is it Cost Effective?. PharmacoEconomics - Open, 2020, 4, 529-539.	1.8	3
221	Molecular Phenotypes of Endothelial Cells in Malignant Tumors. , 2022, , 31-52.		3
222	Significance of tumourâ€associated macrophages, vascular endothelial growth factor and thrombospondinâ€1 expression for tumour angiogenesis and prognosis in endometrial carcinomas. International Journal of Cancer, 1999, 84, 538-543.	5.1	2
223	Papillary Thyroid Carcinoma with Lipomatous Stroma. American Journal of Surgical Pathology, 1997, 21, 1256-1257.	3.7	2
224	Abstract 135: Associations between VEGF polymorphisms and clinical outcome in breast cancer. , 2015, , $\cdot$		2
225	Enhanced expression of EGF receptor and low frequency of. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 1999, 435, 434.	2.8	2
226	Hva er ansvarlig kreftforskning?. Tidsskrift for Den Norske Laegeforening, 2017, 137, 292-294.	0.2	2
227	Prognostic Importance of Tumor Diameter in Nodular Melanomas. Annals of Plastic Surgery, 1997, 38, 550-551.	0.9	1
228	Fibulin-2 expression associates with vascular invasion and patient survival in breast cancer. PLoS ONE, 2021, 16, e0249767.	2.5	1
229	Abstract 1290: The ability of the transcription factor p63 to induce selected gene expression modules associated with mesenchymal to epithelial transition of prostate cells. , 2012, , .		1
230	Abstract 2651: Nestin expression in tumor cells is associated with the basal-like phenotype and aggressive features in breast cancer , 2013, , .		1
231	Abstract 2465: Gene expression, phenotypic and karyotypic analyses of stepwise prostate carcinogenesisin vitro. , 2010, , .		1
232	Abstract 3158: Stathmin expression predicts response to taxanes in metastatic endometrial cancer. , 2011, , .		1
233	Abstract 1384: Axl and VE-cadherin expression as markers of angiogenesis in human breast cancers. , 2012, , .		1
234	Abstract 101: Analysisof GNAQmutations in subtypes of malignant melanoma. , 2012, , .		1

#	Article	IF	CITATIONS
235	Early screening outcomes among non-immigrants and immigrants targeted by BreastScreen Norway, 2010–2019. Scandinavian Journal of Public Health, 2022, , 140349482210787.	2.3	1
236	Letter Thyroid carcinomas with tall-cell and columnar-cell features. Current Diagnostic Pathology, 1996, 3, 67.	0.4	0
237	Angiogenesis – recent developments in molecular pathogenesis, morphological markers, and therapeutic applications. Apmis, 2004, 112, 399-401.	2.0	0
238	Dr. Folkman's legacy—a life of angiogenesis research. Apmis, 2008, 116, 89-92.	2.0	0
239	Gene Expression Signatures of the Tumor Microenvironment: Relation to Tumor Progress in Breast Cancer. , 2017, , 379-407.		0
240	Tissue-Based Biomarkers of Tumor-Vascular Interactions. , 2017, , 55-75.		0
241	Detection and significance of small and low proliferation breast cancer. Journal of Medical Screening, 2021, , 096914132110239.	2.3	0
242	Abstract 330: Gene expression signature characterized by PI3Kinase activation and Stathmin overexpression validates to identify aggressive disease and a potential for PI3Kinase-inhibitors in endometrial carcinomas. , 2011, , .		0
243	Abstract 4191: Prognostic impact of phosphohistone H3 in cutaneous nodular melanoma. , 2011, , .		0
244	Abstract 528: A tumor-stroma interaction study in red mammary tumors in green mice with and without enhanced oxygenation. , 2011, , .		0
245	Abstract 869: Characterization of malignant melanoma families withCDK4germ-line mutation. , 2011, , .		0
246	Abstract 4169: Loss of GPR30 expression identifies estrogen receptor positive endometrial carcinoma patients with poor outcome. , 2011, , .		0
247	Abstract 3089: The role of p63 regulation during epithelial to mesenchymal transition (EMT) and subsequent accumulation of malignant features of primary immortalized prostate cells. , 2011, , .		0
248	Abstract LB-299: Effect of antiangiogenic therapy on tumor growth, vasculature and kinase activity in luminal and basal-like breast cancer xenografts. , 2011, , .		0
249	Abstract 3828: An 18-gene signature for vascular invasion is associated with aggressive features and reduced survival in breast cancer. , 2014, , .		0
250	Abstract 2828: High expression of heat shock protein 27 in metastases is correlated with response to bevacizumab monotherapy in patients with metastatic melanoma. , 2014, , .		0
251	Abstract 2429: Nestin expression in tumor cells is associated with BRCA1 positive genotype and a basal-like phenotype in breast cancer. , 2015, , .		0
252	Abstract 4329: Expression of serpin B2, neuroserpin and L1CAM in association with metastasis and survival in non-small cell lung cancer. , 2015, , .		0

#	Article	IF	CITATIONS
253	Abstract A18: Development of a novel peptide therapeutic that targets ovarian cancer via the tumor microenvironment. , 2016, , .		0
254	Abstract 3937: Nestin expression is associated with a basal-like phenotype and poor prognosis in breast cancer. , 2016, , .		0
255	Abstract 1138: Prognostic impact of BRAF-V600E expression and correlation with mutation status in cutaneous melanoma. , 2016, , .		Ο
256	Abstract P6-10-24: High stromal platelet derived growth factor receptor beta predicts absence of benefit of radiotherapy after breast conserving surgery in breast ductal carcinoma in situ within the SweDCIS randomized trial. , 2020, , .		0