

Lu Zhu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8481693/publications.pdf>

Version: 2024-02-01

34
papers

1,308
citations

304743

22
h-index

377865

34
g-index

35
all docs

35
docs citations

35
times ranked

2119
citing authors

#	ARTICLE	IF	CITATIONS
1	Î²-Arrestins as Important Regulators of Glucose and Energy Homeostasis. Annual Review of Physiology, 2022, 84, 17-40.	13.1	14
2	Exercise during pregnancy may have more benefits than we thought. EBioMedicine, 2022, 77, 103889.	6.1	3
3	Î²-Arrestins as regulators of key metabolic processes. , 2022, , 69-85.		0
4	Mitochondrial GCN5L1 regulates glutaminase acetylation and hepatocellular carcinoma. Clinical and Translational Medicine, 2022, 12, e852.	4.0	14
5	Mitochondrial GCN5L1 regulates cytosolic redox state and hepatic gluconeogenesis via glycerol phosphate shuttle GPD2. Biochemical and Biophysical Research Communications, 2022, 621, 1-7.	2.1	2
6	Use of DREADD Technology to Identify Novel Targets for Antidiabetic Drugs. Annual Review of Pharmacology and Toxicology, 2021, 61, 421-440.	9.4	26
7	Mitochondrial General Control of Amino Acid Synthesis 5 Like 1 Regulates Glutaminolysis, Mammalian Target of Rapamycin Complex 1 Activity, and Murine Liver Regeneration. Hepatology, 2020, 71, 643-657.	7.3	13
8	Adipocyte Gi signaling is essential for maintaining whole-body glucose homeostasis and insulin sensitivity. Nature Communications, 2020, 11, 2995.	12.8	27
9	Î²-arrestin-1 suppresses myogenic reprogramming of brown fat to maintain euglycemia. Science Advances, 2020, 6, eaba1733.	10.3	15
10	Beta-cell M3 muscarinic acetylcholine receptors as potential targets for novel antidiabetic drugs. International Immunopharmacology, 2020, 81, 106267.	3.8	6
11	Adipocyte Î²-arrestin-2 is essential for maintaining whole body glucose and energy homeostasis. Nature Communications, 2019, 10, 2936.	12.8	43
12	Selective activation of Gs signaling in adipocytes causes striking metabolic improvements in mice. Molecular Metabolism, 2019, 27, 83-91.	6.5	25
13	Allosteric modulation of Î²-cell M ₃ muscarinic acetylcholine receptors greatly improves glucose homeostasis in lean and obese mice. Proceedings of the National Academy of Sciences of the United States of America, 2019, 116, 18684-18690.	7.1	22
14	Intra-islet glucagon signaling is critical for maintaining glucose homeostasis. JCI Insight, 2019, 4, .	5.0	102
15	Î² Cellâ€™s intrinsic Î²-arrestin 1 signaling enhances sulfonylurea-induced insulin secretion. Journal of Clinical Investigation, 2019, 129, 3732-3737.	8.2	32
16	Hepatic Gi signaling regulates whole-body glucose homeostasis. Journal of Clinical Investigation, 2018, 128, 746-759.	8.2	34
17	Î²-arrestin-2 is an essential regulator of pancreatic Î²-cell function under physiological and pathophysiological conditions. Nature Communications, 2017, 8, 14295.	12.8	63
18	GCN5L1 modulates cross-talk between mitochondria and cell signaling to regulate FoxO1 stability and gluconeogenesis. Nature Communications, 2017, 8, 523.	12.8	41

#	ARTICLE	IF	CITATIONS
19	Central IGF1 improves glucose tolerance and insulin sensitivity in mice. <i>Nutrition and Diabetes</i> , 2017, 7, 2.	3.2	36
20	Hepatic β -arrestin 2 is essential for maintaining euglycemia. <i>Journal of Clinical Investigation</i> , 2017, 127, 2941-2945.	8.2	40
21	A G Protein-biased Designer G Protein-coupled Receptor Useful for Studying the Physiological Relevance of Gq/11-dependent Signaling Pathways. <i>Journal of Biological Chemistry</i> , 2016, 291, 7809-7820.	3.4	29
22	CK2 acts as a potent negative regulator of receptor-mediated insulin release in vitro and in vivo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, E6818-24.	7.1	27
23	Virus-Mediated Expression of DREADDs for In Vivo Metabolic Studies. <i>Methods in Molecular Biology</i> , 2015, 1335, 205-221.	0.9	2
24	Regulation of Glucose Homeostasis and Lipid Metabolism by PPP1R3G-mediated Hepatic Glycogenesis. <i>Molecular Endocrinology</i> , 2014, 28, 116-126.	3.7	38
25	PAQR3 Modulates Insulin Signaling by Shunting Phosphoinositide 3-Kinase p110 β to the Golgi Apparatus. <i>Diabetes</i> , 2013, 62, 444-456.	0.6	52
26	PAQR3 Has Modulatory Roles in Obesity, Energy Metabolism, and Leptin Signaling. <i>Endocrinology</i> , 2013, 154, 4525-4535.	2.8	38
27	PAQR3 Plays a Suppressive Role in the Tumorigenesis of Colorectal Cancers. <i>Carcinogenesis</i> , 2012, 33, 2228-2235.	2.8	51
28	Tollip, an Intracellular Trafficking Protein, Is a Novel Modulator of the Transforming Growth Factor- β Signaling Pathway. <i>Journal of Biological Chemistry</i> , 2012, 287, 39653-39663.	3.4	69
29	Unraveling the biological functions of Smad7 with mouse models. <i>Cell and Bioscience</i> , 2011, 1, 44.	4.8	20
30	Fasting-Induced Protein Phosphatase 1 Regulatory Subunit Contributes to Postprandial Blood Glucose Homeostasis via Regulation of Hepatic Glycogenesis. <i>Diabetes</i> , 2011, 60, 1435-1445.	0.6	51
31	Hepatic Deletion of Smad7 in Mouse Leads to Spontaneous Liver Dysfunction and Aggravates Alcoholic Liver Injury. <i>PLoS ONE</i> , 2011, 6, e17415.	2.5	27
32	Hydroxytyrosol protects against oxidative damage by simultaneous activation of mitochondrial biogenesis and phase II detoxifying enzyme systems in retinal pigment epithelial cells. <i>Journal of Nutritional Biochemistry</i> , 2010, 21, 1089-1098.	4.2	140
33	Amelioration of high fat diet induced liver lipogenesis and hepatic steatosis by interleukin-22. <i>Journal of Hepatology</i> , 2010, 53, 339-347.	3.7	129
34	Hydroxytyrosol protects retinal pigment epithelial cells from acrolein-induced oxidative stress and mitochondrial dysfunction. <i>Journal of Neurochemistry</i> , 2007, 103, 2690-2700.	3.9	76