

# Jianqiang Wu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/848876/publications.pdf>

Version: 2024-02-01

86  
papers

4,706  
citations

117625

34  
h-index

106344

65  
g-index

90  
all docs

90  
docs citations

90  
times ranked

5163  
citing authors

#	ARTICLE	IF	CITATIONS
1	P2RY14 cAMP signaling regulates Schwann cell precursor self-renewal, proliferation, and nerve tumor initiation in a mouse model of neurofibromatosis. <i>ELife</i> , 2022, 11, .	6.0	5
2	ZmMPK6 and ethylene signalling negatively regulate the accumulation of anti-insect metabolites DIMBOA and DIMBOA- $\beta$ -Glc in maize inbred line A188. <i>New Phytologist</i> , 2021, 229, 2273-2287.	7.3	19
3	<i>Mythimna separata</i> herbivory primes maize resistance in systemic leaves. <i>Journal of Experimental Botany</i> , 2021, 72, 3792-3805.	4.8	12
4	Proteomic analysis of urine reveals biomarkers for the diagnosis and phenotyping of abdominal-type Henoch-Schonlein purpura. <i>Translational Pediatrics</i> , 2021, 10, 510-524.	1.2	5
5	Proteomics applications in biomarker discovery and pathogenesis for abdominal aortic aneurysm. <i>Expert Review of Proteomics</i> , 2021, 18, 305-314.	3.0	16
6	Exploration of Crucial Mediators for Carotid Atherosclerosis Pathogenesis Through Integration of Microbiome, Metabolome, and Transcriptome. <i>Frontiers in Physiology</i> , 2021, 12, 645212.	2.8	11
7	Distinguishing Benign and Malignant Thyroid Nodules and Identifying Lymph Node Metastasis in Papillary Thyroid Cancer by Plasma N-Glycomics. <i>Frontiers in Endocrinology</i> , 2021, 12, 692910.	3.5	10
8	Herbivory-induced systemic signals are likely to be evolutionarily conserved in euphyllophytes. <i>Journal of Experimental Botany</i> , 2021, 72, 7274-7284.	4.8	6
9	Single-Center Experience in the Endovascular Management of the Combination of Isolated Common and Internal Iliac Artery Aneurysms. <i>Frontiers in Surgery</i> , 2021, 8, 693233.	1.4	1
10	Proteomic Analysis of Copper Toxicity in Human Fungal Pathogen <i>Cryptococcus neoformans</i> . <i>Frontiers in Cellular and Infection Microbiology</i> , 2021, 11, 662404.	3.9	5
11	Diagnostic Potential of Plasma IgG N-glycans in Discriminating Thyroid Cancer from Benign Thyroid Nodules and Healthy Controls. <i>Frontiers in Oncology</i> , 2021, 11, 658223.	2.8	7
12	Urinary Proteomics Identifying Novel Biomarkers for the Diagnosis and Phenotyping of Carotid Artery Stenosis. <i>Frontiers in Molecular Biosciences</i> , 2021, 8, 714706.	3.5	7
13	Discovery of potential biomarkers for human atherosclerotic abdominal aortic aneurysm through untargeted metabolomics and transcriptomics. <i>Journal of Zhejiang University: Science B</i> , 2021, 22, 733-745.	2.8	5
14	Parasite dodder enables transfer of bidirectional systemic nitrogen signals between host plants. <i>Plant Physiology</i> , 2021, 185, 1395-1410.	4.8	15
15	A chromosome-scale <i>Gastrodia elata</i> genome and large-scale comparative genomic analysis indicate convergent evolution by gene loss in mycoheterotrophic and parasitic plants. <i>Plant Journal</i> , 2021, 108, 1609-1623.	5.7	38
16	Validation of diagnostic and predictive biomarkers for hereditary angioedema via plasma N-glycomics. <i>Clinical and Translational Allergy</i> , 2021, 11, e12090.	3.2	3
17	Dodder-transmitted mobile signals prime host plants for enhanced salt tolerance. <i>Journal of Experimental Botany</i> , 2020, 71, 1171-1184.	4.8	22
18	Extensive Inter-plant Protein Transfer between <i>Cuscuta</i> Parasites and Their Host Plants. <i>Molecular Plant</i> , 2020, 13, 573-585.	8.3	59

#	ARTICLE	IF	CITATIONS
19	MYC2, MYC3, and MYC4 function additively in wounding-induced jasmonic acid biosynthesis and catabolism. <i>Journal of Integrative Plant Biology</i> , 2020, 62, 1159-1175.	8.5	60
20	<i>Cuscuta australis</i> (dodder) parasite eavesdrops on the host plants' FT signals to flower. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 23125-23130.	7.1	42
21	<i>Cdkn2a</i> Loss in a Model of Neurofibroma Demonstrates Stepwise Tumor Progression to Atypical Neurofibroma and MPNST. <i>Cancer Research</i> , 2020, 80, 4720-4730.	0.9	25
22	Urinary biomarker discovery in gliomas using mass spectrometry-based clinical proteomics. <i>Chinese Neurosurgical Journal</i> , 2020, 6, 11.	0.9	20
23	The oriental armyworm ( <i>Mythimna separata</i> ) feeding induces systemic defence responses within and between maize leaves. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2019, 374, 20180307.	4.0	25
24	An efficient system composed of maize protoplast transfection and HPLC-MS for studying the biosynthesis and regulation of maize benzoxazinoids. <i>Plant Methods</i> , 2019, 15, 144.	4.3	27
25	The host jasmonic acid pathway regulates the transcriptomic changes of dodder and host plant under the scenario of caterpillar feeding on dodder. <i>BMC Plant Biology</i> , 2019, 19, 540.	3.6	10
26	Urinary Proteome Biomarkers for Early Detection of Respiratory Diseases. , 2019, , 135-145.		0
27	miRNAs as a Secret Weapon in the Battlefield of Haustoria, the Interface between Parasites and Host Plants. <i>Molecular Plant</i> , 2018, 11, 354-356.	8.3	4
28	Elevated CO <sub>2</sub> differentially affects tobacco and rice defense against lepidopteran larvae via the jasmonic acid signaling pathway. <i>Journal of Integrative Plant Biology</i> , 2018, 60, 412-431.	8.5	19
29	Ultraviolet-B enhances the resistance of multiple plant species to lepidopteran insect herbivory through the jasmonic acid pathway. <i>Scientific Reports</i> , 2018, 8, 277.	3.3	37
30	Comparative analysis of alfalfa ( <i>Medicago sativa</i> L.) leaf transcriptomes reveals genotype-specific salt tolerance mechanisms. <i>BMC Plant Biology</i> , 2018, 18, 35.	3.6	93
31	Aphid ( <i>Myzus persicae</i> ) feeding on the parasitic plant dodder ( <i>Cuscuta australis</i> ) activates defense responses in both the parasite and soybean host. <i>New Phytologist</i> , 2018, 218, 1586-1596.	7.3	39
32	Urinary candidate biomarkers in an experimental autoimmune myocarditis rat model. <i>Journal of Proteomics</i> , 2018, 179, 71-79.	2.4	34
33	Current understanding of maize and rice defense against insect herbivores. <i>Plant Diversity</i> , 2018, 40, 189-195.	3.7	42
34	Large-scale gene losses underlie the genome evolution of parasitic plant <i>Cuscuta australis</i> . <i>Nature Communications</i> , 2018, 9, 2683.	12.8	145
35	Whole transcriptome analysis of three leaf stages in two cultivars and one of their F1 hybrid of <i>Camellia sinensis</i> L. with differing EGCG content. <i>Tree Genetics and Genomes</i> , 2017, 13, 1.	1.6	10
36	Dynamic changes of urine proteome in a Walker 256 tumor-bearing rat model. <i>Cancer Medicine</i> , 2017, 6, 2713-2722.	2.8	48

#	ARTICLE	IF	CITATIONS
37	Early Detection of Urinary Proteome Biomarkers for Effective Early Treatment of Pulmonary Fibrosis in a Rat Model. <i>Proteomics - Clinical Applications</i> , 2017, 11, 1700103.	1.6	29
38	Stem parasitic plant <i>Cuscuta australis</i> (dodder) transfers herbivory-induced signals among plants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E6703-E6709.	7.1	58
39	Up-regulation of MPK4 increases the feeding efficiency of the green peach aphid under elevated CO <sub>2</sub> in <i>Nicotiana attenuata</i> . <i>Journal of Experimental Botany</i> , 2017, 68, 5923-5935.	4.8	23
40	Transcriptomics and Alternative Splicing Analyses Reveal Large Differences between Maize Lines B73 and Mo17 in Response to Aphid <i>Rhopalosiphum padi</i> Infestation. <i>Frontiers in Plant Science</i> , 2017, 8, 1738.	3.6	47
41	Salt-tolerant and -sensitive alfalfa ( <i>Medicago sativa</i> ) cultivars have large variations in defense responses to the lepidopteran insect <i>Spodoptera litura</i> under normal and salt stress condition. <i>PLoS ONE</i> , 2017, 12, e0181589.	2.5	11
42	Jasmonic acid carboxyl methyltransferase regulates development and herbivory-induced defense response in rice. <i>Journal of Integrative Plant Biology</i> , 2016, 58, 564-576.	8.5	72
43	Oral secretions from <i>Mythimna separata</i> insects specifically induce defence responses in maize as revealed by high-dimensional biological data. <i>Plant, Cell and Environment</i> , 2016, 39, 1749-1766.	5.7	61
44	Genome-wide identification of calcium-dependent protein kinases in soybean and analyses of their transcriptional responses to insect herbivory and drought stress. <i>Scientific Reports</i> , 2016, 6, 18973.	3.3	45
45	Two hAT transposon genes were transferred from Brassicaceae to broomrapes and are actively expressed in some recipients. <i>Scientific Reports</i> , 2016, 6, 30192.	3.3	12
46	COI1-Regulated Hydroxylation of Jasmonoyl-isoleucine Impairs <i>Nicotiana attenuata</i> 's Resistance to the Generalist Herbivore <i>Spodoptera litura</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 2822-2831.	5.2	21
47	The Specific $\alpha$ 1-Adrenergic Receptor Antagonist Prazosin Influences the Urine Proteome. <i>PLoS ONE</i> , 2016, 11, e0164796.	2.5	8
48	Urinary candidate biomarker discovery in a rat unilateral ureteral obstruction model. <i>Scientific Reports</i> , 2015, 5, 9314.	3.3	41
49	Baseline Survey of Root-Associated Microbes of <i>Taxus chinensis</i> (Pilger) Rehd. <i>PLoS ONE</i> , 2015, 10, e0123026.	2.5	14
50	The Parasitic Plant <i>Cuscuta australis</i> Is Highly Insensitive to Abscisic Acid-Induced Suppression of Hypocotyl Elongation and Seed Germination. <i>PLoS ONE</i> , 2015, 10, e0135197.	2.5	19
51	Physiological conditions can be reflected in human urine proteome and metabolome. <i>Expert Review of Proteomics</i> , 2015, 12, 623-636.	3.0	148
52	MAPK signaling: A key element in plant defense response to insects. <i>Insect Science</i> , 2015, 22, 157-164.	3.0	115
53	Scopoletin is a phytoalexin against <i>Alternaria alternata</i> in wild tobacco dependent on jasmonate signalling. <i>Journal of Experimental Botany</i> , 2014, 65, 4305-4315.	4.8	113
54	Fatty acid-amino acid conjugates are essential for systemic activation of salicylic acid-induced protein kinase and accumulation of jasmonic acid in <i>Nicotiana attenuata</i> . <i>BMC Plant Biology</i> , 2014, 14, 326.	3.6	25

#	ARTICLE	IF	CITATIONS
55	Root parasitic plant <i>Orobanche aegyptiaca</i> and shoot parasitic plant <i>Cuscuta australis</i> obtained Brassicaceae-specific strictosidine synthase-like genes by horizontal gene transfer. <i>BMC Plant Biology</i> , 2014, 14, 19.	3.6	57
56	Molecular cloning and characterization of a cytochrome P450 taxoid 9 $\tilde{A}$ -hydroxylase in <i>Ginkgo biloba</i> cells. <i>Biochemical and Biophysical Research Communications</i> , 2014, 443, 938-943.	2.1	22
57	Virus-Induced Gene Silencing in Plant MAPK Research. <i>Methods in Molecular Biology</i> , 2014, 1171, 79-89.	0.9	1
58	High levels of jasmonic acid antagonize the biosynthesis of gibberellins and inhibit the growth of <i>Nicotiana attenuata</i> stems. <i>Plant Journal</i> , 2013, 73, 591-606.	5.7	127
59	The Essential Role of Jasmonic Acid in Plant "Herbivore Interactions" Using the Wild Tobacco <i>Nicotiana attenuata</i> as a Model. <i>Journal of Genetics and Genomics</i> , 2013, 40, 597-606.	3.9	63
60	Silencing Brassinosteroid Receptor <i>BR11</i> Impairs Herbivory-Elicited Accumulation of Jasmonic Acid $\beta$ -oleucine and Diterpene Glycosides, but not Jasmonic Acid and Trypsin Proteinase Inhibitors in <i>Nicotiana attenuata</i> . <i>Journal of Integrative Plant Biology</i> , 2013, 55, 514-526.	8.5	16
61	The Use of VIGS Technology to Study Plant "Herbivore Interactions. <i>Methods in Molecular Biology</i> , 2013, 975, 109-137.	0.9	15
62	<i>Nicotiana attenuata</i> MPK4 suppresses a novel jasmonic acid ( <i>JA</i> ) signaling-independent defense pathway against the specialist insect <i>Manduca sexta</i> , but is not required for the resistance to the generalist <i>Spodoptera littoralis</i> . <i>New Phytologist</i> , 2013, 199, 787-799.	7.3	51
63	Calcium-dependent protein kinases, CDPK4 and CDPK5, affect early steps of jasmonic acid biosynthesis in <i>Nicotiana attenuata</i> . <i>Plant Signaling and Behavior</i> , 2013, 8, e22784.	2.4	25
64	Sugar is an endogenous cue for juvenile-to-adult phase transition in plants. <i>ELife</i> , 2013, 2, e00269.	6.0	279
65	Deep Sequencing Reveals Transcriptome Re-Programming of <i>Taxus</i> Media Cells to the Elicitation with Methyl Jasmonate. <i>PLoS ONE</i> , 2013, 8, e62865.	2.5	71
66	Silencing <i>MPK4</i> in <i>Nicotiana attenuata</i> Enhances Photosynthesis and Seed Production But Compromises Abscisic Acid-Induced Stomatal Closure and Guard Cell-Mediated Resistance to <i>Pseudomonas syringae</i> pv <i>tomato</i> DC3000. <i>Plant Physiology</i> , 2012, 158, 759-776.	4.8	93
67	Silencing <i>Nicotiana attenuata</i> Calcium-Dependent Protein Kinases, <i>CDPK4</i> and <i>CDPK5</i> , Strongly Up-Regulates Wound- and Herbivory-Induced Jasmonic Acid Accumulations. <i>Plant Physiology</i> , 2012, 159, 1591-1607.	4.8	94
68	Three MAPK Kinases, MEK1, SIPKK, and NPK2, are not Involved in Activation of SIPK after Wounding and Herbivore Feeding but Important for Accumulation of Trypsin Proteinase Inhibitors. <i>Plant Molecular Biology Reporter</i> , 2012, 30, 731-740.	1.8	13
69	<i>Arabidopsis</i> Plants Having Defects in Nonsense-Mediated mRNA Decay Factors UPF1, UPF2, and UPF3 Show Photoperiod-Dependent Phenotypes in Development and Stress Responses. <i>Journal of Integrative Plant Biology</i> , 2012, 54, 99-114.	8.5	42
70	SGT1 regulates wounding- and herbivory-induced jasmonic acid accumulation and <i>Nicotiana attenuata</i> 's resistance to the specialist lepidopteran herbivore <i>Manduca sexta</i> . <i>New Phytologist</i> , 2011, 189, 1143-1156.	7.3	36
71	Silencing <i>NOA1</i> Elevates Herbivory-Induced Jasmonic Acid Accumulation and Compromises Most of the Carbon-Based Defense Metabolites in <i>Nicotiana attenuata</i> . <i>Journal of Integrative Plant Biology</i> , 2011, 53, 619-631.	8.5	26
72	For security and stability. <i>Plant Signaling and Behavior</i> , 2011, 6, 1479-1482.	2.4	19

#	ARTICLE	IF	CITATIONS
73	S-Nitrosogluthathione reductase (GSNOR) mediates the biosynthesis of jasmonic acid and ethylene induced by feeding of the insect herbivore <i>Manduca sexta</i> and is important for jasmonate-elicited responses in <i>Nicotiana attenuata</i> . <i>Journal of Experimental Botany</i> , 2011, 62, 4605-4616.	4.8	69
74	Two mitogen-activated protein kinase kinases, MKK1 and MEK2, are involved in wounding- and specialist lepidopteran herbivore <i>Manduca sexta</i> -induced responses in <i>Nicotiana attenuata</i> . <i>Journal of Experimental Botany</i> , 2011, 62, 4355-4365.	4.8	42
75	The multifaceted function of BAK1/SERK3. <i>Plant Signaling and Behavior</i> , 2011, 6, 1322-1324.	2.4	21
76	BAK1 regulates the accumulation of jasmonic acid and the levels of trypsin proteinase inhibitors in <i>Nicotiana attenuata</i> 's responses to herbivory. <i>Journal of Experimental Botany</i> , 2011, 62, 641-652.	4.8	83
77	New Insights into Plant Responses to the Attack from Insect Herbivores. <i>Annual Review of Genetics</i> , 2010, 44, 1-24.	7.6	752
78	Herbivory-induced signalling in plants: perception and action. <i>Plant, Cell and Environment</i> , 2009, 32, 1161-1174.	5.7	221
79	Silencing two herbivory-activated MAP kinases, SIPK and WIPK, does not increase <i>Nicotiana attenuata</i> 's susceptibility to herbivores in the glasshouse and in nature. <i>New Phytologist</i> , 2009, 181, 161-173.	7.3	75
80	<i>PR-13/Thionin</i> But Not <i>PR-1</i> Mediates Bacterial Resistance in <i>Nicotiana attenuata</i> in Nature, and Neither Influences Herbivore Resistance. <i>Molecular Plant-Microbe Interactions</i> , 2008, 21, 988-1000.	2.6	26
81	A Comparison of Two <i>Nicotiana attenuata</i> Accessions Reveals Large Differences in Signaling Induced by Oral Secretions of the Specialist Herbivore <i>Manduca sexta</i> . <i>Plant Physiology</i> , 2008, 146, 927-939.	4.8	68
82	Herbivory Rapidly Activates MAPK Signaling in Attacked and Unattacked Leaf Regions but Not between Leaves of <i>Nicotiana attenuata</i> . <i>Plant Cell</i> , 2007, 19, 1096-1122.	6.6	391
83	Nonsense-mediated mRNA decay (NMD) silences the accumulation of aberrant trypsin proteinase inhibitor mRNA in <i>Nicotiana attenuata</i> . <i>Plant Journal</i> , 2007, 51, 693-706.	5.7	40
84	Evolution of proteinase inhibitor defenses in North American allopolyploid species of <i>Nicotiana</i> . <i>Planta</i> , 2006, 224, 750-760.	3.2	42
85	Differential Elicitation of Two Processing Proteases Controls the Processing Pattern of the Trypsin Proteinase Inhibitor Precursor in <i>Nicotiana attenuata</i> . <i>Plant Physiology</i> , 2005, 139, 375-388.	4.8	34
86	Ageing- and AAA-associated differentially expressed proteins identified by proteomic analysis in mice. <i>PeerJ</i> , 0, 10, e13129.	2.0	2