

Hans-Peter Sinn

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8539224/publications.pdf>

Version: 2024-02-01

266
papers

14,555
citations

28274

55
h-index

24258

110
g-index

310
all docs

310
docs citations

310
times ranked

19873
citing authors

#	ARTICLE	IF	CITATIONS
1	Vacuum-Assisted Breast Biopsy After Neoadjuvant Systemic Treatment for Reliable Exclusion of Residual Cancer in Breast Cancer Patients. <i>Annals of Surgical Oncology</i> , 2022, 29, 1076-1084.	1.5	15
2	Diagnosing Pathologic Complete Response in the Breast After Neoadjuvant Systemic Treatment of Breast Cancer Patients by Minimal Invasive Biopsy. <i>Annals of Surgery</i> , 2022, 275, 576-581.	4.2	38
3	ASO Visual Abstract: Vacuum-Assisted Breast Biopsy After Neoadjuvant Systemic Treatment to Reliably Exclude Residual Cancer in Breast Cancer Patients. <i>Annals of Surgical Oncology</i> , 2022, 29, 1085-1086.	1.5	0
4	MCM3 is a novel proliferation marker associated with longer survival for patients with tubo-ovarian high-grade serous carcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2022, 480, 855-871.	2.8	8
5	Validated biomarker assays confirm that <i>ARID1A</i> loss is confounded with <i>MMR</i> deficiency, <i>CD8</i> ⁺ TIL infiltration, and provides no independent prognostic value in endometriosis-associated ovarian carcinomas. <i>Journal of Pathology</i> , 2022, 256, 388-401.	4.5	15
6	Perivascular tenascin C triggers sequential activation of macrophages and endothelial cells to generate a pro-metastatic vascular niche in the lungs. <i>Nature Cancer</i> , 2022, 3, 486-504.	13.2	35
7	Sex chromosome DSD individuals with mosaic 45,X0 and aberrant Y chromosomes in 46,XY cells: distinct gender phenotypes and germ cell tumour risks. <i>Systems Biology in Reproductive Medicine</i> , 2022, 68, 247-257.	2.1	3
8	AGO Recommendations for the Diagnosis and Treatment of Patients with Early Breast Cancer: Update 2022. <i>Breast Care</i> , 2022, 17, 403-420.	1.4	43
9	AGO Recommendations for the Diagnosis and Treatment of Patients with Locally Advanced and Metastatic Breast Cancer: Update 2022. <i>Breast Care</i> , 2022, 17, 421-429.	1.4	9
10	A multicentre analytical comparison study of inter-reader and inter-assay agreement of four programmed death-ligand 1 immunohistochemistry assays for scoring in triple-negative breast cancer. <i>Histopathology</i> , 2021, 78, 567-577.	2.9	23
11	CaM Kinase II- γ Is Required for Diabetic Hyperglycemia and Retinopathy but Not Nephropathy. <i>Diabetes</i> , 2021, 70, 616-626.	0.6	9
12	Diagnostic accuracy of axillary staging by ultrasound in early breast cancer patients. <i>European Journal of Radiology</i> , 2021, 135, 109468.	2.6	23
13	Immune-related Gene Expression Predicts Response to Neoadjuvant Chemotherapy but not Additional Benefit from PD-L1 Inhibition in Women with Early Triple-negative Breast Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 2584-2591.	7.0	27
14	CATCH: A Prospective Precision Oncology Trial in Metastatic Breast Cancer. <i>JCO Precision Oncology</i> , 2021, 5, 676-686.	3.0	20
15	Reconstructing tumor history in breast cancer: signatures of mutational processes and response to neoadjuvant chemotherapy. <i>Annals of Oncology</i> , 2021, 32, 500-511.	1.2	9
16	Clinicopathologic and molecular analysis of embryonal rhabdomyosarcoma of the genitourinary tract: evidence for a distinct DICER1-associated subgroup. <i>Modern Pathology</i> , 2021, 34, 1558-1569.	5.5	28
17	uPA heteromerization promotes breast cancer progression by attracting tumorigenic neutrophils. <i>EMBO Molecular Medicine</i> , 2021, 13, e13110.	6.9	5
18	66P Baseline menopausal status, Ki-67 and stromal tumour-infiltrating lymphocytes (TILs) and association with outcome in triple-negative breast cancer (TNBC): Exploratory analysis in GeparSixto. <i>Annals of Oncology</i> , 2021, 32, S49-S50.	1.2	0

#	ARTICLE	IF	CITATIONS
19	Therapy response and prognosis of patients with early breast cancer with low positivity for hormone receptors – An analysis of 2765 patients from neoadjuvant clinical trials. <i>European Journal of Cancer</i> , 2021, 148, 159-170.	2.8	41
20	JUNB suppresses distant metastasis by influencing the initial metastatic stage. <i>Clinical and Experimental Metastasis</i> , 2021, 38, 411-423.	3.3	5
21	Breast cancer characteristics and surgery among women with LiéFraumeni syndrome in Germany – A retrospective cohort study. <i>Cancer Medicine</i> , 2021, 10, 7747-7758.	2.8	7
22	AGO Recommendations for the Diagnosis and Treatment of Patients with Locally Advanced and Metastatic Breast Cancer: Update 2021. <i>Breast Care</i> , 2021, 16, 228-235.	1.4	20
23	AGO Recommendations for the Diagnosis and Treatment of Patients with Early Breast Cancer: Update 2021. <i>Breast Care</i> , 2021, 16, 214-227.	1.4	51
24	AGO Recommendations for the Surgical Therapy of the Axilla After Neoadjuvant Chemotherapy: 2021 Update. <i>Geburtshilfe Und Frauenheilkunde</i> , 2021, 81, 1112-1120.	1.8	17
25	Efficacy of intraoperative specimen radiography as margin assessment tool in breast conserving surgery. <i>Breast Cancer Research and Treatment</i> , 2020, 179, 425-433.	2.5	16
26	Eliminating the breast cancer surgery paradigm after neoadjuvant systemic therapy: current evidence and future challenges. <i>Annals of Oncology</i> , 2020, 31, 61-71.	1.2	119
27	DNA methylation-based profiling of uterine neoplasms: a novel tool to improve gynecologic cancer diagnostics. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 97-104.	2.5	29
28	Chromothripsis in Human Breast Cancer. <i>Cancer Research</i> , 2020, 80, 4918-4931.	0.9	11
29	Endometrial Cancer Molecular Risk Stratification is Equally Prognostic for Endometrioid Ovarian Carcinoma. <i>Clinical Cancer Research</i> , 2020, 26, 5400-5410.	7.0	41
30	Frequent Molecular Subtype Switching and Gene Expression Alterations in Lung and Pleural Metastasis From Luminal A – Type Breast Cancer. <i>JCO Precision Oncology</i> , 2020, 4, 848-859.	3.0	7
31	MGMT promoter methylation in triple negative breast cancer of the GeparSixto trial. <i>PLoS ONE</i> , 2020, 15, e0238021.	2.5	8
32	Immunohistological Expression of SOX-10 in Triple-Negative Breast Cancer: A Descriptive Analysis of 113 Samples. <i>International Journal of Molecular Sciences</i> , 2020, 21, 6407.	4.1	18
33	Prediction of pathological complete response in breast cancer patients during neoadjuvant chemotherapy: Is shear wave elastography a useful tool in clinical routine?. <i>European Journal of Radiology</i> , 2020, 128, 109025.	2.6	14
34	Prognostic gene expression signature for high-grade serous ovarian cancer. <i>Annals of Oncology</i> , 2020, 31, 1240-1250.	1.2	85
35	Ultra-High-Value Kurtosis Imaging for Noninvasive Tissue Characterization of Ovarian Lesions. <i>Radiology</i> , 2020, 296, 358-369.	7.3	10
36	Development and Validation of the Gene Expression Predictor of High-grade Serous Ovarian Carcinoma Molecular SubTYPE (PrOTYPE). <i>Clinical Cancer Research</i> , 2020, 26, 5411-5423.	7.0	43

#	ARTICLE	IF	CITATIONS
37	Deep Learning for the Classification of Small-Cell and Non-Small-Cell Lung Cancer. <i>Cancers</i> , 2020, 12, 1604.	3.7	63
38	Clinical and pathological associations of PTEN expression in ovarian cancer: a multicentre study from the Ovarian Tumour Tissue Analysis Consortium. <i>British Journal of Cancer</i> , 2020, 123, 793-802.	6.4	35
39	Metastasis-initiating cells induce and exploit a fibroblast niche to fuel malignant colonization of the lungs. <i>Nature Communications</i> , 2020, 11, 1494.	12.8	115
40	Estrogen, progesterone, and human epidermal growth factor receptor 2 discordance between primary and metastatic breast cancer. <i>Breast Cancer Research and Treatment</i> , 2020, 183, 137-144.	2.5	19
41	Impact of mRNA-Assessed Molecular Subtype Conversion, Intact and Apoptotic Circulating Tumor Cells on Survival of Metastatic Breast Cancer Patients: Proof of Principle. <i>Diagnostics</i> , 2020, 10, 369.	2.6	2
42	Immunohistological expression of oestrogen receptor, progesterone receptor, mammaglobin, human epidermal growth factor receptor 2 and GATA-binding protein 3 in non-small-cell lung cancer. <i>Histopathology</i> , 2020, 77, 900-914.	2.9	6
43	Tumor mutational burden and immune infiltration as independent predictors of response to neoadjuvant immune checkpoint inhibition in early TNBC in GeparNuevo. <i>Annals of Oncology</i> , 2020, 31, 1216-1222.	1.2	128
44	Statistical modelling of HER2-positivity in breast cancer: Final analyses from two large, multicentre, non-interventional studies in Germany. <i>Breast</i> , 2020, 49, 246-253.	2.2	2
45	Endometrial stromal sarcomas with <i>BCOR</i> rearrangement harbor <i>MDM2</i> amplifications. <i>Journal of Pathology: Clinical Research</i> , 2020, 6, 178-184.	3.0	32
46	The Lack of Evidence for an Association between Cancer Biomarker Conversion Patterns and CTC-Status in Patients with Metastatic Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2161.	4.1	6
47	Abstract GS5-03: Diagnosing residual disease and pathologic complete response after neoadjuvant chemotherapy in breast cancer patients by image-guided vacuum-assisted breast biopsy: Results of a prospective multicenter trial. <i>Cancer Research</i> , 2020, 80, GS5-03-GS5-03.	0.9	9
48	Abstract PD5-08: Tumor immune-cell activity assessed by RNAseq is an independent predictor of therapy response and prognosis after neoadjuvant chemotherapy in HER2 negative breast cancer patients - An analysis of the GeparSepto trial. , 2020, , .		0
49	Abstract P6-10-04: Landscape of immune-cell signatures in early high-risk breast cancer (BC) reveals clinically-relevant enrichment of immune subpopulations. , 2020, , .		0
50	Impact of mRNA-assessed intrinsic subtype conversion between primary and metastatic breast cancer on survival. , 2020, 80, .		0
51	Diagnostic accuracy and clinical utility of axillary ultrasound in the evaluation of axillary lymph node status in early breast cancer patients. , 2020, 80, .		0
52	Receptor discordance between primary tumor and metastasis influences CTC-status. , 2020, 80, .		0
53	IgG4-related sclerosing mastitis in a 49-year-old patient with multiple, tumor-like nodules” Diagnostic accuracy of core needle biopsy. <i>Breast Journal</i> , 2019, 25, 1251-1253.	1.0	9
54	Heterogeneous Responses of Axillary Lymph Node Metastases to Neoadjuvant Chemotherapy are Common and Depend on Breast Cancer Subtype. <i>Annals of Surgical Oncology</i> , 2019, 26, 4381-4389.	1.5	18

#	ARTICLE	IF	CITATIONS
55	AGO Recommendations for the Diagnosis and Treatment of Patients with Locally Advanced and Metastatic Breast Cancer: Update 2019. <i>Breast Care</i> , 2019, 14, 247-255.	1.4	32
56	AGO Recommendations for the Diagnosis and Treatment of Patients with Early Breast Cancer: Update 2019. <i>Breast Care</i> , 2019, 14, 224-245.	1.4	72
57	A combination of the immunohistochemical markers CK7 and SATB2 is highly sensitive and specific for distinguishing primary ovarian mucinous tumors from colorectal and appendiceal metastases. <i>Modern Pathology</i> , 2019, 32, 1834-1846.	5.5	54
58	Author's reply to: Comparing the performance of gene expression assays in breast cancer. <i>International Journal of Cancer</i> , 2019, 145, 1163-1164.	5.1	0
59	Molecular Subtype Conversion between Primary and Metastatic Breast Cancer Corresponding to the Dynamics of Apoptotic and Intact Circulating Tumor Cells. <i>Cancers</i> , 2019, 11, 342.	3.7	8
60	Gonadoblastoma Y locus genes expressed in germ cells of individuals with dysgenetic gonads and a Y chromosome in their karyotypes include <i>DDX3Y</i> and <i>TSPY</i> . <i>Human Reproduction</i> , 2019, 34, 770-779.	0.9	21
61	Renal Tubular Dysgenesis in a Case of Fetus Acardius Amorphus. <i>Case Reports in Pathology</i> , 2019, 2019, 1-11.	0.3	1
62	Mismatch Repair Deficiency Drives Durable Complete Remission by Targeting Programmed Death Receptor 1 in a Metastatic Luminal Breast Cancer Patient. <i>Breast Care</i> , 2019, 14, 53-59.	1.4	13
63	Summary of head-to-head comparisons of patient risk classifications by the 21-gene Recurrence Score® (RS) assay and other genomic assays for early breast cancer. <i>International Journal of Cancer</i> , 2019, 145, 882-893.	5.1	32
64	Sustained prognostic impact of circulating tumor cell status and kinetics upon further progression of metastatic breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019, 173, 155-165.	2.5	11
65	Exome analysis of oncogenic pathways and tumor mutational burden (TMB) in triple-negative breast cancer (TNBC): Results of the translational biomarker program of the neoadjuvant double-blind placebo controlled GeparNuevo trial. <i>Journal of Clinical Oncology</i> , 2019, 37, 509-509.	1.6	10
66	Abstract PD2-07: mRNA signatures predict response to durvalumab therapy in triple negative breast cancer (TNBC) – Results of the translational biomarker programme of the neoadjuvant double-blind placebo controlled GeparNuevo trial. <i>Cancer Research</i> , 2019, 79, PD2-07-PD2-07.	0.9	7
67	MyD88 and TLR4 Expression in Epithelial Ovarian Cancer. <i>Mayo Clinic Proceedings</i> , 2018, 93, 307-320.	3.0	22
68	Etiology of hormone receptor positive breast cancer differs by levels of histologic grade and proliferation. <i>International Journal of Cancer</i> , 2018, 143, 746-757.	5.1	19
69	Novel, improved grading system(s) for IDH-mutant astrocytic gliomas. <i>Acta Neuropathologica</i> , 2018, 136, 153-166.	7.7	298
70	Clinical relevance and concordance of HER2 status in local and central testing – an analysis of 1581 HER2-positive breast carcinomas over 12 years. <i>Modern Pathology</i> , 2018, 31, 607-615.	5.5	25
71	Complement Activation in Peritoneal Dialysis-Induced Arteriopathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2018, 29, 268-282.	6.1	45
72	Vacuum-Assisted Biopsy to Diagnose a Pathological Complete Response in Breast Cancer Patients After Neoadjuvant Systemic Therapy. <i>Annals of Surgery</i> , 2018, 268, e60-e61.	4.2	3

#	ARTICLE	IF	CITATIONS
73	Tumour-infiltrating lymphocytes and prognosis in different subtypes of breast cancer: a pooled analysis of 3771 patients treated with neoadjuvant therapy. <i>Lancet Oncology</i> , The, 2018, 19, 40-50.	10.7	1,327
74	Interdisciplinary Screening, Diagnosis, Therapy and Follow-up of Breast Cancer. Guideline of the DGGG and the DKG (S3-Level, AWMF Registry Number 032/045OL, December 2017) – Part 2 with Recommendations for the Therapy of Primary, Recurrent and Advanced Breast Cancer. <i>Geburtshilfe Und Frauenheilkunde</i> , 2018, 78, 1056-1088.	1.8	69
75	Interdisciplinary Screening, Diagnosis, Therapy and Follow-up of Breast Cancer. Guideline of the DGGG and the DKG (S3-Level, AWMF Registry Number 032/045OL, December 2017) – Part 1 with Recommendations for the Screening, Diagnosis and Therapy of Breast Cancer. <i>Geburtshilfe Und Frauenheilkunde</i> , 2018, 78, 927-948.	1.8	59
76	Stress signaling in breast cancer cells induces matrix components that promote chemoresistant metastasis. <i>EMBO Molecular Medicine</i> , 2018, 10, .	6.9	77
77	The cancer-associated microprotein CASIMO1 controls cell proliferation and interacts with squalene epoxidase modulating lipid droplet formation. <i>Oncogene</i> , 2018, 37, 4750-4768.	5.9	111
78	Association of p16 expression with prognosis varies across ovarian carcinoma histotypes: an Ovarian Tumor Tissue Analysis consortium study. <i>Journal of Pathology: Clinical Research</i> , 2018, 4, 250-261.	3.0	70
79	Obesity as risk factor for subtypes of breast cancer: results from a prospective cohort study. <i>BMC Cancer</i> , 2018, 18, 616.	2.6	47
80	RESPONDER – diagnosis of pathological complete response by vacuum-assisted biopsy after neoadjuvant chemotherapy in breast Cancer - a multicenter, confirmative, one-armed, intra-individually-controlled, open, diagnostic trial. <i>BMC Cancer</i> , 2018, 18, 851.	2.6	32
81	Comparison of immunohistochemistry with PCR for assessment of ER, PR, and Ki-67 and prediction of pathological complete response in breast cancer. <i>BMC Cancer</i> , 2017, 17, 124.	2.6	62
82	Intrinsic subtypes and risk scores in ER+/HER2-Breast Cancer: a comparison of Prosigna and OncotypeDX risk categories with Ki67. <i>Breast</i> , 2017, 32, S105.	2.2	2
83	The branched-chain amino acid transaminase 1 sustains growth of antiestrogen-resistant and ER±-negative breast cancer. <i>Oncogene</i> , 2017, 36, 4124-4134.	5.9	60
84	Dose-Response Association of CD8 ⁺ Tumor-Infiltrating Lymphocytes and Survival Time in High-Grade Serous Ovarian Cancer. <i>JAMA Oncology</i> , 2017, 3, e173290.	7.1	260
85	Genistein and enterolactone in relation to Ki67 expression and HER2 status in postmenopausal breast cancer patients. <i>Molecular Nutrition and Food Research</i> , 2017, 61, 1700449.	3.3	13
86	Initial Treatment of Patients with Primary Breast Cancer: Evidence, Controversies, Consensus. <i>Geburtshilfe Und Frauenheilkunde</i> , 2017, 77, 633-644.	1.8	28
87	AGO Recommendations for the Diagnosis and Treatment of Patients with Early Breast Cancer: Update 2017. <i>Breast Care</i> , 2017, 12, 172-183.	1.4	17
88	AGO Recommendations for the Diagnosis and Treatment of Patients with Advanced and Metastatic Breast Cancer: Update 2017. <i>Breast Care</i> , 2017, 12, 184-191.	1.4	11
89	An international reproducibility study validating quantitative determination of ERBB2, ESR1, PGR, and MKI67 mRNA in breast cancer using MammaTyper®. <i>Breast Cancer Research</i> , 2017, 19, 55.	5.0	29
90	Targeted next-generation sequencing enables reliable detection of HER2 (ERBB2) status in breast cancer and provides ancillary information of clinical relevance. <i>Genes Chromosomes and Cancer</i> , 2017, 56, 255-265.	2.8	21

#	ARTICLE	IF	CITATIONS
91	Assessing HER2 testing quality in breast cancer: variables that influence HER2 positivity rate from a large, multicenter, observational study in Germany. <i>Modern Pathology</i> , 2017, 30, 217-226.	5.5	29
92	Identification of a characteristic vascular belt zone in human colorectal cancer. <i>PLoS ONE</i> , 2017, 12, e0171378.	2.5	14
93	Tumor biomarker conversion between primary and metastatic breast cancer: mRNA assessment and its concordance with immunohistochemistry. <i>Oncotarget</i> , 2017, 8, 51416-51428.	1.8	16
94	Abstract P2-05-25: Predictive value of ultra-high ESR1 mRNA expression in early breast cancer. , 2017, , .		0
95	Mutations in genes encoding <sc>PI3K</sc> and <sc>MAPK</sc> signaling define anogenital papillary hidradenoma. <i>Genes Chromosomes and Cancer</i> , 2016, 55, 113-119.	2.8	29
96	Standardized evaluation of tumor-infiltrating lymphocytes in breast cancer: results of the ring studies of the international immuno-oncology biomarker working group. <i>Modern Pathology</i> , 2016, 29, 1155-1164.	5.5	230
97	Lean Umbilical Cord â€“ a Case Report. <i>Geburtshilfe Und Frauenheilkunde</i> , 2016, 76, 1186-1188.	1.8	1
98	Prognosis of breast cancer molecular subtypes in routine clinical care: A large prospective cohort study. <i>BMC Cancer</i> , 2016, 16, 734.	2.6	126
99	Changes in chemotherapy usage and outcome of early breast cancer patients in the last decade. <i>Breast Cancer Research and Treatment</i> , 2016, 160, 491-499.	2.5	54
100	Can a pathological complete response of breast cancer after neoadjuvant chemotherapy be diagnosed by minimal invasive biopsy?. <i>European Journal of Cancer</i> , 2016, 69, 142-150.	2.8	59
101	Prognostic value of automated Ki67 scoring in breast cancer: a centralised evaluation of 8088 patients from 10 study groups. <i>Breast Cancer Research</i> , 2016, 18, 104.	5.0	56
102	High-throughput automated scoring of Ki67 in breast cancer tissue microarrays from the Breast Cancer Association Consortium. <i>Journal of Pathology: Clinical Research</i> , 2016, 2, 138-153.	3.0	19
103	Disseminated Tumor Cells in the Bone Marrow of Patients with Operable Primary Breast Cancer: Prognostic Impact in Immunophenotypic Subgroups and Clinical Implication for Bisphosphonate Treatment. <i>Annals of Surgical Oncology</i> , 2016, 23, 757-766.	1.5	15
104	Do Patients After Reexcision Due to Involved or Close Margins Have the Same Risk of Local Recurrence as Those After One-Step Breast-Conserving Surgery?. <i>Annals of Surgical Oncology</i> , 2016, 23, 1831-1837.	1.5	25
105	Can Routine Imaging After Neoadjuvant Chemotherapy in Breast Cancer Predict Pathologic Complete Response?. <i>Annals of Surgical Oncology</i> , 2016, 23, 789-795.	1.5	84
106	Role of TP53 mutations in triple negative and HER2-positive breast cancer treated with neoadjuvant anthracycline/taxane-based chemotherapy. <i>Oncotarget</i> , 2016, 7, 67686-67698.	1.8	50
107	Premalignant and Malignant Breast Pathology. , 2016, , 179-194.		0
108	Abstract P1-07-12: Prognosis of clinico-pathological breast cancer subtypes in routine clinical care. , 2016, , .		0

#	ARTICLE	IF	CITATIONS
109	Abstract P1-03-05: SPARC expression in primary metastatic breast cancer. , 2016, , .		0
110	Predictive value of ultra-high ESR1 mRNA expression in early breast cancer.. Journal of Clinical Oncology, 2016, 34, e12045-e12045.	1.6	0
111	Distribution of <i>MED12</i> mutations in fibroadenomas and phyllodes tumors of the breast—implications for tumor biology and pathological diagnosis. Genes Chromosomes and Cancer, 2015, 54, 444-452.	2.8	55
112	Tumor-Infiltrating Lymphocytes and Response to Neoadjuvant Chemotherapy With or Without Carboplatin in Human Epidermal Growth Factor Receptor 2-Positive and Triple-Negative Primary Breast Cancers. Journal of Clinical Oncology, 2015, 33, 983-991.	1.6	863
113	WHO grade related expression of TRAIL-receptors and apoptosis regulators in meningioma. Pathology Research and Practice, 2015, 211, 109-116.	2.3	11
114	Reprogramming of the ER α and ER β Target Gene Landscape Triggers Tamoxifen Resistance in Breast Cancer. Cancer Research, 2015, 75, 720-731.	0.9	36
115	S100P and HYAL2 as prognostic markers for patients with triple-negative breast cancer. Experimental and Molecular Pathology, 2015, 99, 180-187.	2.1	21
116	Breast Cancers with a <i>BRCA1</i> -like DNA Copy Number Profile Recur Less Often Than Expected after High-Dose Alkylating Chemotherapy. Clinical Cancer Research, 2015, 21, 763-770.	7.0	34
117	Predictors of Residual Tumor in Breast-Conserving Therapy. Annals of Surgical Oncology, 2015, 22, 451-458.	1.5	12
118	Abstract P5-10-17: Evaluation of PgR status by immunohistochemistry may be inferior to PgR results by Oncotype DX for assessing the recurrence risk in ER+/HER2- breast cancer with low or intermediate tumor proliferation. , 2015, , .		0
119	Assessing HER2 testing quality in breast cancer (BC): Variables that influence HER2-positivity from a large, multicenter, observational study in Germany.. Journal of Clinical Oncology, 2015, 33, 11062-11062.	1.6	1
120	Differentiation and histogenesis of syringomatous tumour of the nipple and low-grade adenosquamous carcinoma: evidence for a common origin. Histopathology, 2014, 65, 9-23.	2.9	41
121	Bortezomib Sensitizes Primary Meningioma Cells to TRAIL-Induced Apoptosis by Enhancing Formation of the Death-Inducing Signaling Complex. Journal of Neuropathology and Experimental Neurology, 2014, 73, 1034-1046.	1.7	18
122	Ki-67 and p53 expression of the fallopian tube mucosa in breast cancer patients with hereditary risk. Archives of Gynecology and Obstetrics, 2014, 289, 1079-1085.	1.7	2
123	Comparison of molecular abnormalities in vulvar and vaginal melanomas. Modern Pathology, 2014, 27, 1386-1393.	5.5	70
124	Pre-diagnostic smoking behaviour and poorer prognosis in a German breast cancer patient cohort — Differential effects by tumour subtype, NAT2 status, BMI and alcohol intake. Cancer Epidemiology, 2014, 38, 419-426.	1.9	19
125	<i>PIK3CA</i> Mutations Are Associated With Lower Rates of Pathologic Complete Response to Anti-Human Epidermal Growth Factor Receptor 2 (HER2) Therapy in Primary HER2-Overexpressing Breast Cancer. Journal of Clinical Oncology, 2014, 32, 3212-3220.	1.6	231
126	Reverse phase protein array based tumor profiling identifies a biomarker signature for risk classification of hormone receptor-positive breast cancer. Translational Proteomics, 2014, 2, 52-59.	1.2	19

#	ARTICLE	IF	CITATIONS
127	Co-expression of MET and CD47 is a novel prognosticator for survival of luminal-type breast cancer patients. <i>Oncotarget</i> , 2014, 5, 8147-8160.	1.8	83
128	Mutational profiles in triple-negative breast cancer defined by ultradeep multigene sequencing show high rates of PI3K pathway alterations and clinically relevant entity subgroup specific differences. <i>Oncotarget</i> , 2014, 5, 9952-9965.	1.8	58
129	Expression of SPARC and response to nab-paclitaxel (nab-p) in patients (pts) with metastatic breast cancer (MBC).. <i>Journal of Clinical Oncology</i> , 2014, 32, e12009-e12009.	1.6	0
130	Efficacy of nab-paclitaxel does not seem to be associated with SPARC expression in metastatic breast cancer. <i>Anticancer Research</i> , 2014, 34, 6609-15.	1.1	32
131	Giant struma ovarii. <i>Archives of Gynecology and Obstetrics</i> , 2013, 287, 399-400.	1.7	2
132	Interobserver agreement of proliferation index (Ki-67) outperforms mitotic count in pulmonary carcinoids. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2013, 462, 507-513.	2.8	63
133	Intrinsic breast cancer subtypes defined by estrogen receptor signallingâ€”prognostic relevance of progesterone receptor loss. <i>Modern Pathology</i> , 2013, 26, 1161-1171.	5.5	47
134	Omission of Axillary Dissection According to ACOSOG Z0011: Impact on Adjuvant Treatment Recommendations. <i>Annals of Surgical Oncology</i> , 2013, 20, 1538-1544.	1.5	23
135	Circulating Fibronectin Controls Tumor Growth. <i>Neoplasia</i> , 2013, 15, 925-IN24.	5.3	55
136	Biological subtypes of triple-negative breast cancer are associated with distinct morphological changes and clinical behaviour. <i>Breast</i> , 2013, 22, 986-992.	2.2	35
137	Prediction of underestimated invasiveness in patients with ductal carcinoma inÂsitu of the breast on percutaneous biopsy as rationale for recommending concurrent sentinel lymph node biopsy. <i>Breast</i> , 2013, 22, 537-542.	2.2	48
138	Identification of a population of blood circulating tumor cells from breast cancer patients that initiates metastasis in a xenograft assay. <i>Nature Biotechnology</i> , 2013, 31, 539-544.	17.5	920
139	A Brief Overview of the WHO Classification of Breast Tumors, 4th Edition, Focusing on Issues and Updates from the 3rd Edition. <i>Breast Care</i> , 2013, 8, 149-154.	1.4	280
140	Effects of mTOR Inhibition On IR/IGF-1R Signalling in PIK3CA-Mutated, Tamoxifen Resistant Breast Cancer. <i>Annals of Oncology</i> , 2013, 24, iii25.	1.2	0
141	Multigene Assays for Classification, Prognosis, and Prediction in Breast Cancer: a Critical Review on the Background and Clinical Utility. <i>Geburtshilfe Und Frauenheilkunde</i> , 2013, 73, 932-940.	1.8	55
142	Protein phosphatase 1, regulatory subunit 15B is a survival factor for ERÎ±-positive breast cancer. <i>International Journal of Cancer</i> , 2013, 132, 2714-2719.	5.1	7
143	Neoadjuvant epirubicin, gemcitabine and docetaxel for primary breast cancer: Long-term survival data and major prognostic factors based on two consecutive neoadjuvant phase I/II trials. <i>International Journal of Cancer</i> , 2013, 133, 1006-1015.	5.1	12
144	Comparison of EndoPredict and Oncotype DX Test Results in Hormone Receptor Positive Invasive Breast Cancer. <i>PLoS ONE</i> , 2013, 8, e58483.	2.5	59

#	ARTICLE	IF	CITATIONS
145	Plasma MicroRNA Panel for Minimally Invasive Detection of Breast Cancer. PLoS ONE, 2013, 8, e76729.	2.5	112
146	Abstract S1-06: Increased tumor-associated lymphocytes predict benefit from addition of carboplatin to neoadjuvant therapy for triple-negative and HER2-positive early breast cancer in the GeparSixto trial (GBG 66). , 2013, , .		16
147	Abstract S4-06: PIK3CA mutation predicts resistance to anti-HER2/chemotherapy in primary HER2-positive/hormone-receptor-positive breast cancer – Prospective analysis of 737 participants of the GeparSixto and GeparQuinto studies. Cancer Research, 2013, 73, S4-06-S4-06.	0.9	8
148	Abstract P1-08-07: High tumor CD68 mRNA content (intratumoral macrophages) predicts response to neoadjuvant chemotherapy. , 2013, , .		0
149	Abstract P1-08-15: Prognostic value of five different histologic scoring systems for tumor regression after neoadjuvant chemotherapy of breast cancer in luminal type breast cancer. , 2013, , .		0
150	19p13.1 Is a Triple-Negative – Specific Breast Cancer Susceptibility Locus. Cancer Research, 2012, 72, 1795-1803.	0.9	100
151	Interdisciplinary Consensus Recommendations for the use of Vacuum-Assisted Breast Biopsy under Sonographic Guidance: First update 2012. Ultraschall in Der Medizin, 2012, 33, 366-371.	1.5	51
152	Deubiquitination of EGFR by Cezanne-1 contributes to cancer progression. Oncogene, 2012, 31, 4599-4608.	5.9	84
153	Extent of Primary Breast Cancer Surgery: Standards and Individualized Concepts. Breast Care, 2012, 7, 364-369.	1.4	11
154	Transitions Between Flat Epithelial Atypia and Low-grade Ductal Carcinoma In Situ of the Breast. American Journal of Surgical Pathology, 2012, 36, 1247-1252.	3.7	13
155	847 Identification of a Biomarker Signature to Predict the Need for Chemotherapy in Patients With Hormone Receptor Positive Breast Cancer. European Journal of Cancer, 2012, 48, S203-S204.	2.8	0
156	Genome-wide methylation screen in low-grade breast cancer identifies novel epigenetically altered genes as potential biomarkers for tumor diagnosis. FASEB Journal, 2012, 26, 4937-4950.	0.5	84
157	Predictors of resectability in breast-conserving therapy. Archives of Gynecology and Obstetrics, 2012, 286, 1023-1031.	1.7	7
158	A meta-analysis of genome-wide association studies of breast cancer identifies two novel susceptibility loci at 6q14 and 20q11. Human Molecular Genetics, 2012, 21, 5373-5384.	2.9	168
159	CD24 Ala57Val polymorphism predicts pathologic complete response to sequential anthracycline- and taxane-based neoadjuvant chemotherapy for primary breast cancer. Breast Cancer Research and Treatment, 2012, 132, 819-831.	2.5	21
160	Outcome analysis of patients with primary breast cancer initially treated at a certified academic breast unit. Breast, 2012, 21, 303-308.	2.2	42
161	Abstract P3-06-08: Ki-67 mRNA as a predictor for response to neoadjuvant chemotherapy in primary breast cancer. , 2012, , .		2
162	Associations of Breast Cancer Risk Factors With Tumor Subtypes: A Pooled Analysis From the Breast Cancer Association Consortium Studies. Journal of the National Cancer Institute, 2011, 103, 250-263.	6.3	596

#	ARTICLE	IF	CITATIONS
163	Low penetrance breast cancer susceptibility loci are associated with specific breast tumor subtypes: findings from the Breast Cancer Association Consortium. <i>Human Molecular Genetics</i> , 2011, 20, 3289-3303.	2.9	152
164	A common variant at the TERT-CLPTM1L locus is associated with estrogen receptor-negative breast cancer. <i>Nature Genetics</i> , 2011, 43, 1210-1214.	21.4	279
165	Automatic Tumor-Stroma Separation in Fluorescence TMAs Enables the Quantitative High-Throughput Analysis of Multiple Cancer Biomarkers. <i>PLoS ONE</i> , 2011, 6, e28048.	2.5	13
166	Triple-Negative Breast Cancer: Clinical and Histological Correlations. <i>Breast Care</i> , 2011, 6, 273-278.	1.4	39
167	Determination of paraneoplastic autoimmune responses by tumor cell biology and intratumoral IFN-alpha/IL-12 in breast cancer patients. <i>Cancer Immunology, Immunotherapy</i> , 2011, 60, 401-411.	4.2	4
168	Synthetic Antitumor Vaccines Containing MUC1 Glycopeptides with Two Immunodominant Domains Induction of a Strong Immune Response against Breast Tumor Tissues. <i>Angewandte Chemie - International Edition</i> , 2011, 50, 9977-9981.	13.8	90
169	Variable expression of the Fragile X Mental Retardation 1 (FMR1) gene in patients with premature ovarian failure syndrome is not dependent on number of (CGG) _n triplets in exon 1. <i>Human Reproduction</i> , 2011, 26, 1241-1251.	0.9	36
170	High Nuclear Poly(Adenosine Diphosphate-Ribose) Polymerase Expression Is Predictive for BRCA1- and BRCA2-Deficient Breast Cancer. <i>Journal of Clinical Oncology</i> , 2011, 29, 4586-4588.	1.6	17
171	Common Breast Cancer Susceptibility Loci Are Associated with Triple-Negative Breast Cancer. <i>Cancer Research</i> , 2011, 71, 6240-6249.	0.9	109
172	Reply to L. Ozretić et al. <i>Journal of Clinical Oncology</i> , 2011, 29, 4588-4589.	1.6	1
173	A randomized phase II trial of doxorubicin plus pemetrexed followed by docetaxel versus doxorubicin plus cyclophosphamide followed by docetaxel as neoadjuvant treatment of early breast cancer. <i>Annals of Oncology</i> , 2011, 22, 609-617.	1.2	18
174	Basoluminal and luminal phenotypes in triple-negative breast cancer: Immunohistochemical profiling and survival. <i>Journal of Clinical Oncology</i> , 2011, 29, 1098-1098.	1.6	0
175	Quality management and accreditation of research tissue banks: experience of the National Center for Tumor Diseases (NCT) Heidelberg. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2010, 457, 741-747.	2.8	31
176	Robust gridding of TMAs after whole-slide imaging using template matching. <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2010, 77A, 1169-1176.	1.5	11
177	Simulation of heat transfer in zone plate optics irradiated by X-ray free electron laser radiation. <i>Nuclear Instruments and Methods in Physics Research, Section A: Accelerators, Spectrometers, Detectors and Associated Equipment</i> , 2010, 621, 620-626.	1.6	11
178	FGFR4 Arg388 genotype is associated with pathological complete response to neoadjuvant chemotherapy for primary breast cancer. <i>Annals of Oncology</i> , 2010, 21, 1636-1642.	1.2	31
179	Early Breast Cancer Precursor Lesions: Lessons Learned from Molecular and Clinical Studies. <i>Breast Care</i> , 2010, 5, 218-226.	1.4	24
180	Invasive ductal breast cancer within a malignant phyllodes tumor: case report and assessment of clonality. <i>Human Pathology</i> , 2010, 41, 293-296.	2.0	21

#	ARTICLE	IF	CITATIONS
181	Reduction of CD44+/CD24 ^{low} breast cancer cells by conventional cytotoxic chemotherapy. <i>Human Pathology</i> , 2010, 41, 574-581.	2.0	32
182	Predictive value of CD24 ala/val polymorphism for pathologic complete response to sequential anthracycline- and taxane-based neoadjuvant chemotherapy for primary breast cancer.. <i>Journal of Clinical Oncology</i> , 2010, 28, 543-543.	1.6	9
183	Invasive Tubular Carcinoma of the Breast Frequently is Clonally Related to Flat Epithelial Atypia and Low-grade Ductal Carcinoma In Situ. <i>American Journal of Surgical Pathology</i> , 2009, 33, 1646-1653.	3.7	48
184	Intratumoral Cytokines and Tumor Cell Biology Determine Spontaneous Breast Cancer ^{â€} Specific Immune Responses and Their Correlation to Prognosis. <i>Cancer Research</i> , 2009, 69, 8420-8428.	0.9	55
185	Estrogen-Related Receptor β Expression and Function Is Associated with the Transcriptional Coregulator AIB1 in Breast Carcinoma. <i>Cancer Research</i> , 2009, 69, 5186-5193.	0.9	28
186	0178 Results of clinical endpoints of a randomized phase II trial with doxorubicin + pemetrexed followed by docetaxel versus doxorubicin + cyclophosphamide followed by docetaxel as primary systemic therapy for early breast cancer. <i>Breast</i> , 2009, 18, S63.	2.2	0
187	Breast cancer precursors: lessons learned from molecular genetics. <i>Journal of Molecular Medicine</i> , 2009, 87, 113-115.	3.9	14
188	Prognostic significance of tumour necrosis factor-related apoptosis-inducing ligand (TRAIL) receptor expression in patients with breast cancer. <i>Journal of Molecular Medicine</i> , 2009, 87, 995-1007.	3.9	72
189	Treatment of advanced metastasized breast cancer with bone marrow-derived tumour-reactive memory T cells: a pilot clinical study. <i>Cancer Immunology, Immunotherapy</i> , 2009, 58, 887-900.	4.2	46
190	The role of molecular analysis in breast cancer. <i>Pathology</i> , 2009, 41, 77-88.	0.6	44
191	Clonality of lobular carcinoma in situ (LCIS) and metachronous invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2008, 107, 331-335.	2.5	50
192	Clinical response after two cycles compared to HER2, Ki-67, p53, and bcl-2 in independently predicting a pathological complete response after preoperative chemotherapy in patients with operable carcinoma of the breast. <i>Breast Cancer Research</i> , 2008, 10, R30.	5.0	96
193	The histone acetyltransferase hMOF is frequently downregulated in primary breast carcinoma and medulloblastoma and constitutes a biomarker for clinical outcome in medulloblastoma. <i>International Journal of Cancer</i> , 2008, 122, 1207-1213.	5.1	146
194	Oblimersen combined with docetaxel, adriamycin and cyclophosphamide as neo-adjuvant systemic treatment in primary breast cancer: final results of a multicentric phase I study. <i>Annals of Oncology</i> , 2008, 19, 1698-1705.	1.2	11
195	Gains of chromosome region 3q26 in intraepithelial neoplasia and invasive squamous cell carcinoma of the vulva are frequent and independent of HPV status. <i>Journal of Clinical Pathology</i> , 2008, 61, 1034-1037.	2.0	24
196	Safety and pharmacokinetics of bivatuzumab mertansine in patients with CD44v6-positive metastatic breast cancer: final results of a phase I study. <i>Anti-Cancer Drugs</i> , 2007, 18, 477-485.	1.4	75
197	35 Tumor endothelial marker 1 (TEM1, endosialin) is expressed in functional and morphological distinct trophoblast cells in human and mouse placenta. <i>Thrombosis Research</i> , 2007, 119, S106.	1.7	0
198	Doxorubicin/Pemetrexed Followed by Docetaxel Versus Doxorubicin/Cyclophosphamide Followed by Docetaxel as Neoadjuvant Treatment for Early-Stage Breast Cancer: A Randomized Phase II Trial. <i>Clinical Breast Cancer</i> , 2007, 7, 555-558.	2.4	6

#	ARTICLE	IF	CITATIONS
199	Reduced incidence of distant metastases and lower mortality in 1072 patients with breast cancer with a history of hormone replacement therapy. <i>American Journal of Obstetrics and Gynecology</i> , 2007, 196, 342.e1-342.e9.	1.3	35
200	HER2 overexpression is associated with shorter disease-free survival (DFS) following neoadjuvant systemic therapy (NST) of primary breast cancer (PBC) patients with gemcitabine (G), epirubicin (E) and docetaxel (Doc). <i>Journal of Clinical Oncology</i> , 2007, 25, 11087-11087.	1.6	0
201	Oblimersen (O) in combination with docetaxel (T), adriamycin (A), and cyclophosphamide (C) as neoadjuvant systemic therapy (NST) in primary breast cancer (PBC): Final results of a multicentric phase I study. <i>Journal of Clinical Oncology</i> , 2007, 25, 11047-11047.	1.6	0
202	Comparison of immunohistological parameters in primary breast cancers and corresponding locoregional recurrences. <i>Pathology Research and Practice</i> , 2006, 202, 125-130.	2.3	2
203	Consensus recommendations for the use of vacuum-assisted breast biopsy under sonographic guidance. <i>Gynecological Surgery</i> , 2006, 3, 309-314.	0.9	2
204	Gene Expression Signature Predicting Pathologic Complete Response With Gemcitabine, Epirubicin, and Docetaxel in Primary Breast Cancer. <i>Journal of Clinical Oncology</i> , 2006, 24, 1839-1845.	1.6	146
205	c-myc Amplifications in primary breast carcinomas and their local recurrences. <i>Journal of Clinical Pathology</i> , 2006, 59, 424-4248.	2.0	49
206	Dose-dense primary systemic chemotherapy with gemcitabine plus epirubicin sequentially followed by docetaxel for early breast cancer: final results of a phase I/II trial. <i>Anti-Cancer Drugs</i> , 2005, 16, 1023-1028.	1.4	11
207	Immunohistochemical evaluation of endothelial nitric oxide synthase expression in primary breast cancer. <i>Breast</i> , 2005, 14, 230-235.	2.2	10
208	Immunohistochemical and cytogenetic characterization of acantholytic squamous cell carcinoma of the breast. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2005, 446, 305-309.	2.8	12
209	Intraoperative venous blood sampling to localize a small androgen-producing ovarian tumor. <i>Gynecological Endocrinology</i> , 2005, 21, 138-141.	1.7	17
210	The role of early expression of inducible nitric oxide synthase in human breast cancer. <i>European Journal of Cancer</i> , 2005, 41, 265-271.	2.8	88
211	Micrometastatic bone marrow cells at diagnosis have no impact on survival of primary breast cancer patients with extensive axillary lymph node involvement treated with stem cell-supported high-dose chemotherapy. <i>Annals of Oncology</i> , 2004, 15, 1627-1632.	1.2	1
212	Prediction of the axillary lymph node status in mammary cancer on the basis of clinicopathological data and flow cytometry. <i>Medical and Biological Engineering and Computing</i> , 2004, 42, 733-739.	2.8	16
213	Gemcitabine, epirubicin and docetaxel as primary systemic therapy in patients with early breast cancer: results of a multicentre phase I/II study. <i>European Journal of Cancer</i> , 2004, 40, 2432-2438.	2.8	22
214	Only grading has independent impact on breast cancer survival after adjustment for pathological response to preoperative chemotherapy. <i>Anti-Cancer Drugs</i> , 2004, 15, 127-135.	1.4	23
215	Intronic TP53 Germline Sequence Variants Modify the Risk in German Breast/Ovarian Cancer Families. <i>Hereditary Cancer in Clinical Practice</i> , 2004, 2, 139.	1.5	2
216	CTCF Gene Mutations in Invasive Ductal Breast Cancer. <i>Breast Cancer Research and Treatment</i> , 2003, 80, 347-352.	2.5	38

#	ARTICLE	IF	CITATIONS
217	Evaluation of neoadjuvant chemotherapeutic response of breast cancer using dynamic MRI with high temporal resolution. <i>European Radiology</i> , 2003, 13, 80-87.	4.5	84
218	Efficient engraftment of human primary breast cancer transplants in nonconditioned NOD/Scid mice. <i>International Journal of Cancer</i> , 2003, 105, 444-453.	5.1	44
219	Centrosomal aberrations in primary invasive breast cancer are associated with nodal status and hormone receptor expression. <i>International Journal of Cancer</i> , 2003, 107, 346-352.	5.1	55
220	Breast cancer in young women (≤35 years): Genomic aberrations detected by comparative genomic hybridization. <i>International Journal of Cancer</i> , 2003, 107, 583-592.	5.1	54
221	Similar contributions of BRCA1 and BRCA2 germline mutations to early-onset breast cancer in Germany. <i>European Journal of Human Genetics</i> , 2003, 11, 464-467.	2.8	21
222	Accuracy of tumor size measurement in breast cancer using MRI is influenced by histological regression induced by neoadjuvant chemotherapy. <i>European Radiology</i> , 2003, 13, 1213-1223.	4.5	112
223	International Expert Panel on the Use of Primary (Preoperative) Systemic Treatment of Operable Breast Cancer: Review and Recommendations. <i>Journal of Clinical Oncology</i> , 2003, 21, 2600-2608.	1.6	322
224	Prognosefaktoren bei der Planung organübergreifender Debulkingoperation und intraoperativer Strahlentherapie bei Beckenwandrezidiven gynäkologischer Karzinome. <i>Geburtshilfe Und Frauenheilkunde</i> , 2002, 62, 554-559.	1.8	1
225	Magnetic Resonance Imaging of Nude Mice With Heterotransplanted High-Grade Squamous Cell Carcinomas. <i>Investigative Radiology</i> , 2002, 37, 193-198.	6.2	25
226	Characteristics associated with long-term progression-free survival following high-dose chemotherapy in metastatic breast cancer and influence of chemotherapy dose. <i>Annals of Oncology</i> , 2002, 13, 679-688.	1.2	10
227	Anti-p53 in Breast Cancer: Concordance of Different Assay Procedures and Association with p53 Antigen Expression. <i>Oncology</i> , 2002, 63, 297-305.	1.9	19
228	P53 is the strongest predictor of survival in high-risk primary breast cancer patients undergoing high-dose chemotherapy with autologous blood stem cell support. <i>International Journal of Cancer</i> , 2002, 100, 290-296.	5.1	29
229	Expression of endothelial and inducible nitric oxide synthase in benign and malignant lesions of the breast and measurement of nitric oxide using electron paramagnetic resonance spectroscopy. <i>Cancer</i> , 2002, 95, 1191-1198.	4.1	70
230	Stem Cell Dose and Tumorbiologic Parameters as Prognostic Markers for Patients with Metastatic Breast Cancer Undergoing High-Dose Chemotherapy with Autologous Blood Stem Cell Support. <i>Stem Cells</i> , 2002, 20, 32-40.	3.2	8
231	C-myc oncogene amplification in ductal carcinoma in situ of the breast. <i>Breast Cancer Research and Treatment</i> , 2002, 74, 25-31.	2.5	41
232	Clinical response after two cycles is superior to HER2, Ki-67, p53, and bcl-2 in independently predicting a pathological complete response after preoperative chemotherapy in patients with operable carcinoma of the breast. <i>European Journal of Cancer</i> , 2002, 38, S107.	2.8	0
233	Tandem high-dose chemotherapy in high-risk primary breast cancer: A multivariate analysis and a matched-pair comparison with standard-dose chemotherapy. <i>Biology of Blood and Marrow Transplantation</i> , 2001, 7, 332-342.	2.0	11
234	Ductal epithelial proliferations of the breast: a biological continuum? Comparative genomic hybridization and high-molecular-weight cytokeratin expression patterns. <i>Journal of Pathology</i> , 2001, 195, 415-421.	4.5	112

#	ARTICLE	IF	CITATIONS
235	Prognostic value of codon 918 (ATC?ACG)RET proto-oncogene mutations in sporadic medullary thyroid carcinoma. <i>International Journal of Cancer</i> , 2001, 95, 62-66.	5.1	127
236	Clinical relevance of genomic aberrations in homogeneously treated high-risk stage II/III breast cancer patients. <i>International Journal of Cancer</i> , 2001, 93, 80-84.	5.1	34
237	Survival and tumor characteristics of German hereditary breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2000, 59, 185-192.	2.5	48
238	Stage IIa Cervix Carcinoma with Metastasis to the Heart: Report of a Case with Immunohistochemistry, Flow Cytometry, and Virology Findings. <i>Gynecologic Oncology</i> , 2000, 76, 133-138.	1.4	5
239	Prognostic value of cathepsin D in breast cancer. <i>British Journal of Cancer</i> , 1999, 79, 189-190.	6.4	35
240	Trypsinogen activation peptides (TAP) in peritoneal fluid as predictors of late histopathologic injury in necrotizing pancreatitis of the rat. <i>Digestive Diseases and Sciences</i> , 1999, 44, 823-829.	2.3	8
241	Pathophysiologic basis of contrast enhancement in breast tumors. <i>Journal of Magnetic Resonance Imaging</i> , 1999, 10, 260-266.	3.4	360
242	Expression of endothelial and inducible nitric oxide synthase in benign and malignant lesions of the breast. <i>European Journal of Cancer</i> , 1999, 35, S86.	2.8	0
243	Aberrant cytoplasmic expression of the p16 protein in breast cancer is associated with accelerated tumour proliferation. <i>British Journal of Cancer</i> , 1998, 78, 1661-1668.	6.4	88
244	Extensive and predominant in situ component in breast carcinoma: their influence on treatment results after breast-conserving therapy. <i>European Journal of Cancer</i> , 1998, 34, 646-653.	2.8	53
245	Cyclins and Breast Cancer. <i>Oncology Research and Treatment</i> , 1998, 21, 28-34.	1.2	1
246	Response to primary chemotherapy of breast cancer. <i>European Journal of Cancer</i> , 1997, 33, S147.	2.8	0
247	Benefits of various dextrans after delayed therapy in necrotizing pancreatitis of the rat. <i>Intensive Care Medicine</i> , 1996, 22, 1207-1213.	8.2	2
248	Laser-Induced Fluorescence Diagnosis and Photodynamic Therapy of Human Renal Cell Carcinoma. <i>Urologia Internationalis</i> , 1995, 55, 197-201.	1.3	6
249	Comparison of immunohistochemistry and RT-PCR for detection of CD44 expression, a new prognostic factor in human breast cancer. <i>International Journal of Cancer</i> , 1995, 60, 471-477.	5.1	149
250	Human mammary carcinomas express homologues of rat metastasis-associated variants of CD44. <i>Breast Cancer Research and Treatment</i> , 1995, 36, 307-313.	2.5	64
251	Hyperoncotic Dextran and Systemic Aprotinin in Necrotizing Rodent Pancreatitis. <i>Scandinavian Journal of Gastroenterology</i> , 1995, 30, 812-816.	1.5	21
252	CD44 variant exon epitopes in primary breast cancer and length of survival. <i>Lancet, The</i> , 1995, 345, 615-619.	13.7	344

#	ARTICLE	IF	CITATIONS
253	CD44 isoforms in prognosis of breast cancer. <i>Lancet, The</i> , 1995, 346, 502.	13.7	14
254	Effects of passive immunization against parathyroid hormone-related protein: PTHrP is the responsible factor in mediating hypercalcemia in the walker carcinosarcoma 256 rat model. <i>Journal of Bone and Mineral Research</i> , 1995, 10, 7-16.	2.8	9
255	Relevance of CD44 Variant Epitopes for Diagnosis and Therapy of Human Breast Cancer. <i>Contributions To Oncology / Beitrage Zur Onkologie</i> , 1995, , 36-50.	0.1	0
256	Diagnosis and Surgical Treatment of Gastric Sarcoma. <i>Oncology Research and Treatment</i> , 1994, 17, 391-396.	1.2	2
257	Metastatic potential of small and minimally invasive breast carcinomas. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 1994, 425, 237-41.	2.8	14
258	Effect of N-methyl-N ² -nitro-N-nitrosoguanidine on carbohydrate profiles of non-metaplastic rat gastric mucosa. <i>Journal of Cancer Research and Clinical Oncology</i> , 1993, 119, 155-159.	2.5	0
259	Characterization of New Monoclonal Antibodies Directed Against Normal Human Exocrine Pancreas and Pancreatic Adenocarcinomas. <i>Pancreas</i> , 1993, 8, 279-288.	1.1	2
260	ROLE OF NATURAL KILLER CELLS IN THE PATHOGENESIS OF HUMAN ACUTE GRAFT-VERSUS-HOST DISEASE. <i>Transplantation</i> , 1993, 56, 113-119.	1.0	27
261	Chromogranin A as Tumor Marker in Medullary Thyroid Carcinoma. <i>Thyroid</i> , 1992, 2, 5-10.	4.5	41
262	Analysis of the lewisx epitope in human pancreas and pancreatic adenocarcinomas. <i>International Journal of Gastrointestinal Cancer</i> , 1992, 11, 125-35.	0.4	15
263	Nucleolar organizer regions in myogenic stromal tumours of the stomach. <i>Virchows Archiv A, Pathological Anatomy and Histopathology</i> , 1989, 415, 317-321.	1.4	20
264	Early myocardial lesions in mice with EMC virus myocarditis. <i>Electrocardiographic and morphological findings. European Heart Journal</i> , 1987, 8, 411-415.	2.2	3
265	Histological Alterations of the Preampullary Common Bile and Pancreatic Duct in Acute Biliary and Nonbiliary Pancreatitis. <i>Digestion</i> , 1986, 34, 93-100.	2.3	6
266	A program nucleus for encoding and decoding calendar dates. <i>Computer Methods and Programs in Biomedicine</i> , 1986, 23, 317-318.	4.7	0