
## Tim J Peters

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8560789/publications.pdf Version: 2024-02-01



TIM I DETEDS

#	Article	IF	CITATIONS
1	10-Year Outcomes after Monitoring, Surgery, or Radiotherapy for Localized Prostate Cancer. New England Journal of Medicine, 2016, 375, 1415-1424.	13.9	2,101
2	ICIQ: A brief and robust measure for evaluating the symptoms and impact of urinary incontinence. Neurourology and Urodynamics, 2004, 23, 322-330.	0.8	1,478
3	Patient-Reported Outcomes after Monitoring, Surgery, or Radiotherapy for Prostate Cancer. New England Journal of Medicine, 2016, 375, 1425-1437.	13.9	962
4	Quality improvement report: Improving design and conduct of randomised trials by embedding them in qualitative research: ProtecT (prostate testing for cancer and treatment) study * Commentary: presenting unbiased information to patients can be difficult. BMJ: British Medical Journal, 2002, 325, 766-770.	2.4	461
5	Valuing the ICECAP capability index for older people. Social Science and Medicine, 2008, 67, 874-882.	1.8	422
6	The International Continence Society "Benign Prostatic Hyperplasia―Study: The Bothersomeness of Urinary Symptoms. Journal of Urology, 1997, 157, 885-889.	0.2	281
7	Cognitive behavioural therapy as an adjunct to pharmacotherapy for primary care based patients with treatment resistant depression: results of the CoBalT randomised controlled trial. Lancet, The, 2013, 381, 375-384.	6.3	278
8	Evaluation of computer based clinical decision support system and risk chart for management of hypertension in primary care: randomised controlled trial. BMJ: British Medical Journal, 2000, 320, 686-690.	2.4	225
9	SCORING THE SHORT FORM ICSmaleSF QUESTIONNAIRE. Journal of Urology, 2000, 164, 1948-1955.	0.2	208
10	Screening in the community to reduce fractures in older women (SCOOP): a randomised controlled trial. Lancet, The, 2018, 391, 741-747.	6.3	206
11	Active monitoring, radical prostatectomy, or radiotherapy for localised prostate cancer: study design and diagnostic and baseline results of the ProtecT randomised phase 3 trial. Lancet Oncology, The, 2014, 15, 1109-1118.	5.1	205
12	Relation between parasuicide, suicide, psychiatric admissions, and socioeconomic deprivation. BMJ: British Medical Journal, 1995, 311, 226-230.	2.4	187
13	Childhood energy intake and adult mortality from cancer: the boyd orr cohort study. BMJ: British Medical Journal, 1998, 316, 499-504.	2.4	155
14	Scoring the Icecapâ€a Capability Instrument. Estimation of a UK General Population Tariff. Health Economics (United Kingdom), 2015, 24, 258-269.	0.8	153
15	Socioâ€economic and dietary influences on leg length and trunk length in childhood: a reanalysis of the Carnegie (Boyd Orr) survey of diet and health in prewar Britain (1937–39). Paediatric and Perinatal Epidemiology, 1998, 12, 96-113.	0.8	138
16	Validity of the Short Mood and Feelings Questionnaire in late adolescence Psychological Assessment, 2014, 26, 752-762.	1.2	137
17	Development of a complex intervention improved randomization and informed consent in a randomized controlled trial. Journal of Clinical Epidemiology, 2009, 62, 29-36.	2.4	133
18	Randomised controlled trial comparing effectiveness and acceptability of an early discharge, hospital at home scheme with acute hospital care. BMJ: British Medical Journal, 1998, 316, 1796-1801.	2.4	129

Tim J Peters

#	Article	IF	CITATIONS
19	Hospital at home or acute hospital care? A cost minimisation analysis. BMJ: British Medical Journal, 1998, 316, 1802-1806.	2.4	119
20	Prevalence of treatment-resistant depression in primary care: cross-sectional data. British Journal of General Practice, 2013, 63, e852-e858.	0.7	115
21	What do people with MS want and expect from health-care services?. Health Expectations, 2001, 4, 29-37.	1.1	110
22	Relationships between lower urinary tract symptoms and bladder outlet obstruction: Results from the ICS-"BPH―Study. Neurourology and Urodynamics, 1998, 17, 99-108.	0.8	107
23	Ten-year Mortality, Disease Progression, and Treatment-related Side Effects in Men with Localised Prostate Cancer from the ProtecT Randomised Controlled Trial According to Treatment Received. European Urology, 2020, 77, 320-330.	0.9	107
24	Prevalence and Predictors of Persistent Speech Sound Disorder at Eight Years Old: Findings From a Population Cohort Study. Journal of Speech, Language, and Hearing Research, 2016, 59, 647-673.	0.7	104
25	Long-term effectiveness and cost-effectiveness of cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: follow-up of the CoBalT randomised controlled trial. Lancet Psychiatry,the, 2016, 3, 137-144.	3.7	103
26	GPs' experiences of primary care mental health research: a qualitative study of the barriers to recruitment. Family Practice, 2007, 24, 518-525.	0.8	84
27	A RANDOMIZED TRIAL COMPARING TRANSURETHRAL RESECTION OF THE PROSTATE, LASER THERAPY AND CONSERVATIVE TREATMENT OF MEN WITH SYMPTOMS ASSOCIATED WITH BENIGN PROSTATIC ENLARGEMENT: THE CLasP STUDY. Journal of Urology, 2000, 164, 65-70.	0.2	81
28	Using the ICSQoL to measure the impact of lower urinary tract symptoms on quality of life: evidence from the ICS-'BPH' study. BJU International, 1997, 80, 712-721.	1.3	78
29	The clinical effectiveness of sertraline in primary care and the role of depression severity and duration (PANDA): a pragmatic, double-blind, placebo-controlled randomised trial. Lancet Psychiatry,the, 2019, 6, 903-914.	3.7	78
30	Factors associated with change in pain and disability over time: a community-based prospective observational study of hip and knee osteoarthritis. British Journal of General Practice, 2005, 55, 205-11.	0.7	73
31	The process and delivery of CBT for depression in adults: a systematic review and network meta-analysis. Psychological Medicine, 2019, 49, 1937-1947.	2.7	70
32	Clinical effectiveness and cost-effectiveness of cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: the CoBalT randomised controlled trial. Health Technology Assessment, 2014, 18, 1-167, vii-viii.	1.3	69
33	Domestic violence and mental health: a cross-sectional survey of women seeking help from domestic violence support services. Global Health Action, 2016, 9, 29890.	0.7	66
34	Patientâ€reported outcomes in the ProtecT randomized trial of clinically localized prostate cancer treatments: study design, and baseline urinary, bowel and sexual function and quality of life. BJU International, 2016, 118, 869-879.	1.3	52
35	Factors associated with term low birthweight in Cardiff, Wales. Paediatric and Perinatal Epidemiology, 1997, 11, 287-297.	0.8	51
36	Screen-detected prostate cancer and the insulin-like growth factor axis: Results of a population-based case-control study. International Journal of Cancer, 2004, 108, 887-892.	2.3	51

Tim J Peters

#	Article	IF	CITATIONS
37	How much change is enough? Evidence from a longitudinal study on depression in UK primary care. Psychological Medicine, 2022, 52, 1875-1882.	2.7	51
38	Preferences for aspects of a dermatology consultation. British Journal of Dermatology, 2006, 155, 387-392.	1.4	49
39	Management of Patients With High Baseline Hip Fracture Risk by FRAX Reduces Hip Fractures—A Post Hoc Analysis of the SCOOP Study. Journal of Bone and Mineral Research, 2018, 33, 1020-1026.	3.1	45
40	Determination of Prostatic Volume with Transrectal Ultrasound: A Study of Intra-Observer and Interobserver Variation. Journal of Urology, 1996, 155, 1299-1300.	0.2	44
41	Mirtazapine added to SSRIs or SNRIs for treatment resistant depression in primary care: phase III randomised placebo controlled trial (MIR). BMJ: British Medical Journal, 2018, 363, k4218.	2.4	44
42	Cluster randomised controlled trial comparing the effectiveness and cost-effectiveness of two primary care interventions aimed at improving attendance for breast screening. Journal of Medical Screening, 2001, 8, 91-98.	1.1	42
43	Measuring the psychosocial impact of population-based prostate-specific antigen testing for prostate cancer in the UK. BJU International, 2006, 98, 777-782.	1.3	40
44	Domestic violence and mental health: a cross-sectional survey of women seeking help from domestic violence support services. Global Health Action, 2014, 7, 25519.	0.7	40
45	Effect of a Digital Intervention on Depressive Symptoms in Patients With Comorbid Hypertension or Diabetes in Brazil and Peru. JAMA - Journal of the American Medical Association, 2021, 325, 1852.	3.8	40
46	Vision-related quality of life impairment in an elderly UK population: Associations with age, sex, social class and material deprivation. Eye, 2001, 15, 739-744.	1.1	38
47	Improving attendance for breast screening among recent non-attenders: a randomised controlled trial of two interventions in primary care. Journal of Medical Screening, 2001, 8, 99-105.	1.1	37
48	The epidemiology of childhood eczema: II. Statistical analyses to identify independent early predictors. Paediatric and Perinatal Epidemiology, 1987, 1, 80-94.	0.8	36
49	Cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment resistant depression in primary care: The CoBalT randomised controlled trial protocol. Contemporary Clinical Trials, 2012, 33, 312-319.	0.8	32
50	Prediction of sudden infant death syndrome: An independent evaluation of four scoring methods. Statistics in Medicine, 1986, 5, 113-126.	0.8	27
51	Guidelines and educational outreach visits from community pharmacists to improve prescribing in general practice: a randomised controlled trial. Journal of Health Services Research and Policy, 2001, 6, 207-213.	0.8	27
52	Long-term effects of the Active for Life Year 5 (AFLY5) school-based cluster-randomised controlled trial. BMJ Open, 2016, 6, e010957.	0.8	27
53	Multifarious terminology: multivariable or multivariate? univariable or univariate?. Paediatric and Perinatal Epidemiology, 2008, 22, 506-506.	0.8	25
54	Quantifying response shift or adaptation effects in quality of life by synthesising best-worst scaling and discrete choice data. Journal of Choice Modelling, 2013, 6, 34-43.	1.2	25

#	Article	IF	CITATIONS
55	Systematic screening using FRAX® leads to increased use of, and adherence to, anti-osteoporosis medications: an analysis of the UK SCOOP trial. Osteoporosis International, 2020, 31, 67-75.	1.3	25
56	Intervening to reduce anxiety for women with mild dyskaryosis: do we know what works and why?. Journal of Advanced Nursing, 1998, 28, 563-570.	1.5	24
57	Management of treatment-resistant depression in primary care: a mixed-methods study. British Journal of General Practice, 2018, 68, e673-e681.	0.7	23
58	Functional and quality of life outcomes of localised prostate cancer treatments (Prostate Testing) Tj ETQq0 0 0 r	gBT /Over 1.3	ock 10 Tf 50
59	Pilot study of a two-arm non-randomized controlled cluster trial of a psychosocial intervention to improve late life depression in socioeconomically deprived areas of SA£o Paulo, Brazil (PROACTIVE):	1.2	22

	feasibility study of a psychosocial intervention for late life depression in SAŁo Paulo. BMC Public Health. 2019. 19. 1152.		
60	Active monitoring, radical prostatectomy and radical radiotherapy in PSA-detected clinically localised prostate cancer: the ProtecT three-arm RCT. Health Technology Assessment, 2020, 24, 1-176.	1.3	22
61	Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. Health Technology Assessment, 2018, 22, 1-136.	1.3	21
62	Clinical- and cost-effectiveness of the STAR care pathway compared to usual care for patients with chronic pain after total knee replacement: study protocol for a UK randomised controlled trial. Trials, 2018, 19, 132.	0.7	20
63	Impact of antibiotics for children presenting to general practice with cough on adverse outcomes: secondary analysis from a multicentre prospective cohort study. British Journal of General Practice, 2018, 68, e682-e693.	0.7	19
64	A collaborative care psychosocial intervention to improve late life depression in socioeconomically deprived areas of Guarulhos, Brazil: the PROACTIVE cluster randomised controlled trial protocol. Trials, 2020, 21, 914.	0.7	19
65	The population requirement for cataract extraction: A cross-sectional study. Eye, 2001, 15, 745-752.	1.1	18
66	Psychological advocacy toward healing (PATH): study protocol for a randomized controlled trial. Trials, 2013, 14, 221.	0.7	17
67	Psychological advocacy towards healing (PATH): A randomized controlled trial of a psychological intervention in a domestic violence service setting. PLoS ONE, 2018, 13, e0205485.	1.1	17
68	Transurethral Prostate Resection, Noncontact Laser Therapy or Conservative Management in Men With Symptoms of Benign Prostatic Enlargement? An Economic Evaluation. Journal of Urology, 2002, 168, 2476-2482.	0.2	16
69	A prospective cohort and extended comprehensive-cohort design provided insights about the generalizability of a pragmatic trial: the ProtecT prostate cancer trial. Journal of Clinical Epidemiology, 2018, 96, 35-46.	2.4	16
70	The STAR care pathway for patients with pain at 3 months after total knee replacement: a multicentre, pragmatic, randomised, controlled trial. Lancet Rheumatology, The, 2022, 4, e188-e197.	2.2	16
71	A randomised controlled trial assessing the severity and duration of depressive symptoms associated with a clinically significant response to sertraline versus placebo, in people presenting to primary care with depression (PANDA trial): study protocol for a randomised controlled trial. Trials, 2017, 18, 496.	0.7	15
72	The ProtecT randomised trial cost-effectiveness analysis comparing active monitoring, surgery, or radiotherapy for prostate cancer. British Journal of Cancer, 2020, 123, 1063-1070.	2.9	15

#	Article	IF	CITATIONS
73	Educational outcomes associated with persistent speech disorder. International Journal of Language and Communication Disorders, 2021, 56, 299-312.	0.7	14
74	Comparison between self-administered depression questionnaires and patients' own views of changes in their mood: a prospective cohort study in primary care. Psychological Medicine, 2021, 51, 853-860.	2.7	13
75	Developing a scale to measure parental attitudes towards preschool speech and language therapy services. International Journal of Language and Communication Disorders, 2001, 36, 503-513.	0.7	11
76	Identifying Scoliosis in Population-Based Cohorts: Automation of a Validated Method Based on Total Body Dual Energy X-ray Absorptiometry Scans. Calcified Tissue International, 2020, 106, 378-385.	1.5	11
77	Does father's unemployment put the fetus at risk?. BJOG: an International Journal of Obstetrics and Gynaecology, 1986, 93, 704-710.	1.1	10
78	Mirtazapine added to selective serotonin reuptake inhibitors for treatment-resistant depression in primary care (MIR trial): study protocol for a randomised controlled trial. Trials, 2016, 17, 66.	0.7	10
79	Psychometric properties of the PHQ-9 measure of depression among Brazilian older adults. Aging and Mental Health, 2022, 26, 2285-2290.	1.5	10
80	The Relationship between LUTS and sexual function. Prostate Cancer and Prostatic Diseases, 2001, 4, S2-S6.	2.0	9
81	Assessing the validity, responsiveness and reliability of the Recap measure of eczema control. British Journal of Dermatology, 2021, 184, 955-957.	1.4	9
82	Investigating Software Requirements for Systems Supporting Task-Shifted Interventions: Usability Study. Journal of Medical Internet Research, 2019, 21, e11346.	2.1	9
83	Development of guidelines to reduce, handle and report missing data in palliative care trials: A multi-stakeholder modified nominal group technique. Palliative Medicine, 2022, 36, 59-70.	1.3	9
84	The design and analysis of randomized controlled trials of treatments for lower urinary tract symptoms. BJU International, 2000, 85, 3-9.	1.3	8
85	An expert consensus on the most effective components of cognitive behavioural therapy for adults with depression: a modified Delphi study. Cognitive Behaviour Therapy, 2020, 49, 242-255.	1.9	8
86	Evaluating an enhanced quality improvement intervention in maternity units: PReCePT trial protocol. BMJ Open Quality, 2021, 10, e001204.	0.4	8
87	Challenges in Using Recommended QualityÂofÂLife Measures to Assess Fluctuating Health: A Think-Aloud Study to Understand How Recall and Timing of Assessment Influence Patient Responses. Patient, 2022, 15, 445-457.	1.1	8
88	Radiotherapy for Prostate Cancer: is it â€~what you do' or â€~the way that you do it'? A UK Perspective on Technique and Quality Assurance. Clinical Oncology, 2016, 28, e92-e100.	0.6	7
89	Post-consultation illness trajectories in children with acute cough and respiratory tract infection: prospective cohort study. Family Practice, 2018, 35, 676-683.	0.8	7
90	Screening to Identify Postoperative Pain and <scp>Crossâ€5ectional</scp> Associations Between Factors Identified in This Process With Pain and Function, Three Months After Total Knee Replacement. Arthritis Care and Research, 2022, 74, 790-798.	1.5	7

#	Article	IF	CITATIONS
91	Cost-Effectiveness of Different Formats for Delivery of Cognitive Behavioral Therapy for Depression: A Systematic Review Based Economic Model. Value in Health, 2020, 23, 1662-1670.	0.1	7
92	Strategies adopted by men to deal with uncertainty and anxiety when following an active surveillance/monitoring protocol for localised prostate cancer and implications for care: a longitudinal qualitative study embedded within the ProtecT trial. BMJ Open, 2020, 10, e036024.	0.8	7
93	Risk factors associated with poor pain outcomes following primary knee replacement surgery: Analysis of data from the clinical practice research datalink, hospital episode statistics and patient reported outcomes as part of the STAR research programme. PLoS ONE, 2021, 16, e0261850.	1.1	7
94	Day in, Day Out: The Everyday Therapy of Community Clinics. International Journal of Language and Communication Disorders, 2001, 36, 435-440.	0.7	6
95	Factors associated with trial recruitment, preferences, and treatments received were elucidated in a comprehensive cohort study. Journal of Clinical Epidemiology, 2019, 113, 200-213.	2.4	6
96	Materials used to support cognitive behavioural therapy for depression: a survey of therapists' clinical practice and views. Cognitive Behaviour Therapy, 2019, 48, 463-481.	1.9	6
97	Post-operative determinants of chronic pain after primary knee replacement surgery: Analysis of data on 258,386 patients from the National Joint Registry for England, Wales, Northern Ireland and the Isle of Man (NJR). Osteoarthritis and Cartilage Open, 2021, 3, 100139.	0.9	6
98	Challenges in developing capability measures for children and young people for use in the economic evaluation of health and care interventions. Health Economics (United Kingdom), 2021, 30, 1990-2003.	0.8	6
99	Antidepressant treatment with sertraline for adults with depressive symptoms in primary care: the PANDA research programme including RCT. Programme Grants for Applied Research, 2019, 7, 1-108.	0.4	6
100	Depressive and subthreshold depressive symptomatology among older adults in a socioeconomically deprived area in Brazil. International Journal of Geriatric Psychiatry, 2022, 37, .	1.3	5
101	Locally developed guidelines for the aftercare of deliberate self-harm patients in general practice. Primary Health Care Research and Development, 2003, 4, 21-28.	0.5	4
102	Sex differences in depressive symptoms and tolerability after treatment with selective serotonin reuptake inhibitor antidepressants: Secondary analyses of the GENPOD trial. Journal of Psychopharmacology, 2021, 35, 919-927.	2.0	4
103	Does father's unemployment put the fetus at risk?. BJOC: an International Journal of Obstetrics and Gynaecology, 1986, 93, 704-710.	1.1	3
104	PROTO (Prostate Trials Office). BJU International, 2000, 85, 1-2.	1.3	3
105	Early speech- and language-impaired children: linguistic, literacy, and social outcomes. Developmental Medicine and Child Neurology, 2007, 48, 489-494.	1.1	3
106	Parental and clinician agreement of illness severity in children with RTIs: secondary analysis of data from a prospective cohort study. British Journal of General Practice, 2019, 69, e236-e245.	0.7	3
107	Cost-Effectiveness of Sertraline in Primary Care According to Initial Severity and Duration of Depressive Symptoms: Findings from the PANDA RCT. PharmacoEconomics - Open, 2020, 4, 427-438.	0.9	3
108	A clinical tool to identify older women with back pain at high risk of osteoporotic vertebral fractures (Vfrac): a population-based cohort study with exploratory economic evaluation. Age and Ageing, 2022, 51, .	0.7	3

#	Article	IF	CITATIONS
109	The Children and Young People Quality of Life Study: A protocol for the qualitative development of attributes for capability wellbeing measures for use in health economic evaluation with children and young people. Wellcome Open Research, 0, 7, 117.	0.9	3
110	Acceptability and fidelity of a psychosocial intervention (PROACTIVE) for older adults with depression in a basic health unit in São Paulo, Brazil: a qualitative study. BMC Public Health, 2021, 21, 2278.	1.2	3
111	Does anxiety moderate the effectiveness of mirtazapine in patients with treatment-resistant depression? A secondary analysis of the MIR trial. Journal of Psychopharmacology, 2020, 34, 1342-1349.	2.0	2
112	A Validation Study of the EQ-5D-5L and ICEpop Capability Measure for Older People Among Older Individuals With Depressive Symptoms in Brazil. Value in Health Regional Issues, 2022, 30, 91-99.	0.5	2
113	Can We Trust the Quality of Routine Hospital Outpatient Information in the UK?. Journal of Health Services Research and Policy, 1998, 3, 203-206.	0.8	1
114	Public views of coronavirus science and scientists: findings from a cross-sectional survey. Wellcome Open Research, 0, 6, 166.	0.9	1
115	General practice based intervention to prevent repeat episodes of deliberate self harm: cluster randomised controlled trial. , 0, .		1
116	Running a Randomised Controlled Trial in a Service Setting: A Progress Report. International Journal of Language and Communication Disorders, 1998, 33, 31-36.	0.7	0
117	Reconciling different measures of risk in the treatment of hypertension: a community-based study. Journal of Human Hypertension, 1998, 12, 391-395.	1.0	Ο
118	The Significance of Terminal Dribbling in Men With Lower Urinary Tract Symptoms. Journal of Urology, 1998, 160, 959-959.	0.2	0
119	Relationships Between Lower Urinary Tract Symptoms and Bladder Outlet Obstruction: Results From the ICS-"BPH―Study. Journal of Urology, 1999, 161, 1041-1042.	0.2	0