## Patrick Meimoun

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/857784/publications.pdf Version: 2024-02-01



DATRICK MEIMOUN

#	Article	IF	CITATIONS
1	International Expert Consensus Document on Takotsubo Syndrome (Part I): Clinical Characteristics, Diagnostic Criteria, and Pathophysiology. European Heart Journal, 2018, 39, 2032-2046.	2.2	972
2	International Expert Consensus Document on Takotsubo Syndrome (Part II): Diagnostic Workup, Outcome, and Management. European Heart Journal, 2018, 39, 2047-2062.	2.2	521
3	Usefulness of exercise-stress echocardiography for risk stratification of true asymptomatic patients with aortic valve stenosis. European Heart Journal, 2010, 31, 1390-1397.	2.2	231
4	Non-invasive assessment of coronary flow and coronary flow reserve by transthoracic Doppler echocardiography: a magic tool for the real world. European Journal of Echocardiography, 2008, 9, 449-457.	2.3	104
5	Standard and Advanced Echocardiography in Takotsubo (Stress) Cardiomyopathy: Clinical and Prognostic Implications. Journal of the American Society of Echocardiography, 2015, 28, 57-74.	2.8	97
6	The Coronary Flow Reserve Is Transiently Impaired in Tako-Tsubo Cardiomyopathy: A Prospective Study Using Serial Doppler Transthoracic Echocardiography. Journal of the American Society of Echocardiography, 2008, 21, 72-77.	2.8	94
7	Takotsubo cardiomyopathy: an integrated multi-imaging approach. European Heart Journal Cardiovascular Imaging, 2014, 15, 366-377.	1.2	69
8	Multimodality imaging in takotsubo syndrome: a joint consensus document of the European Association of Cardiovascular Imaging (EACVI) and the Japanese Society of Echocardiography (JSE). European Heart Journal Cardiovascular Imaging, 2020, 21, 1184-1207.	1.2	45
9	Assessment of left ventricular twist mechanics in Tako-tsubo cardiomyopathy by two-dimensional speckle-tracking echocardiography. European Journal of Echocardiography, 2011, 12, 931-939.	2.3	41
10	Comparison Between Non-Invasive Coronary Flow Reserve and Fractional Flow Reserve to Assess the Functional Significance of Left Anterior Descending Artery Stenosis of Intermediate Severity. Journal of the American Society of Echocardiography, 2011, 24, 374-381.	2.8	40
11	Transient impairment of coronary flow reserve in tako-tsubo cardiomyopathy is related to left ventricular systolic parameters. European Journal of Echocardiography, 2008, 10, 265-270.	2.3	35
12	Multimodality imaging in takotsubo syndrome: a joint consensus document of the European Association of Cardiovascular Imaging (EACVI) and the Japanese Society of Echocardiography (JSE). Journal of Echocardiography, 2020, 18, 199-224.	0.8	35
13	Prognostic value of transthoracic coronary flow reserve in medically treated patients with proximal left anterior descending artery stenosis of intermediate severity. European Journal of Echocardiography, 2009, 10, 127-132.	2.3	34
14	Evaluation of Left Anterior Descending Coronary Artery Stenosis of Intermediate Severity Using Transthoracic Coronary Flow Reserve and Dobutamine Stress Echocardiography. Journal of the American Society of Echocardiography, 2005, 18, 1233-1240.	2.8	32
15	Usefulness of Noninvasive Myocardial Work to Predict Left Ventricular Recovery and Acute Complications after Acute Anterior Myocardial Infarction Treated by Percutaneous Coronary Intervention. Journal of the American Society of Echocardiography, 2020, 33, 1180-1190.	2.8	32
16	Significance of systolic anterior motion of the mitral valve during dobutamine stress echocardiography. Journal of the American Society of Echocardiography, 2005, 18, 49-56.	2.8	30
17	Non-invasive detection of tako-tsubo cardiomyopathy vs. acute anterior myocardial infarction by transthoracic Doppler echocardiography. European Heart Journal Cardiovascular Imaging, 2013, 14, 464-470.	1.2	24
18	Factors Associated with Noninvasive Coronary Flow Reserve in Severe Aortic Stenosis. Journal of the American Society of Echocardiography, 2012, 25, 835-841.	2.8	22

PATRICK MEIMOUN

#	Article	IF	CITATIONS
19	Left atrial strain and distensibility in relation to left ventricular dysfunction and prognosis in aortic stenosis. Echocardiography, 2019, 36, 469-477.	0.9	21
20	Non-Invasive Coronary Flow Reserve After Successful Primary Angioplasty for Acute Anterior Myocardial Infarction Is an Independent Predictor of Left Ventricular Recovery and In-Hospital Cardiac Events. Journal of the American Society of Echocardiography, 2009, 22, 1071-1079.	2.8	18
21	Non-invasive coronary flow reserve after successful primary angioplasty for acute anterior myocardial infarction is an independent predictor of left ventricular adverse remodelling. European Journal of Echocardiography, 2010, 11, 711-718.	2.3	15
22	Relationship between acute strain pattern and recovery in tako-tsubo cardiomyopathy and acute anterior myocardial infarction: a comparative study using two-dimensional longitudinal strain. International Journal of Cardiovascular Imaging, 2014, 30, 1491-1500.	1.5	15
23	Usefulness of Two-Dimensional Longitudinal StrainÂPattern to Predict Left Ventricular Recovery andÂln-Hospital Complications after Acute Anterior Myocardial Infarction Treated Successfully byÂPrimaryÂAngioplasty. Journal of the American Society of Echocardiography, 2015, 28, 1366-1375.	2.8	15
24	The left atrial function is transiently impaired in Tako-tsubo cardiomyopathy and associated to in-hospital complications: a prospective study using two-dimensional strain. International Journal of Cardiovascular Imaging, 2020, 36, 299-307.	1.5	8
25	Noninvasive Coronary Flow Reserve Predicts Response to Exercise in Asymptomatic Severe Aortic Stenosis. Journal of the American Society of Echocardiography, 2017, 30, 736-744.	2.8	7
26	Systolic anterior motion of the mitral valve in tako-tsubo cardiomyopathy: Still a matter of debate?. Annales De Cardiologie Et D'Angeiologie, 2015, 64, 385-389.	0.6	5
27	Assessment of left anterior descending artery stenosis of intermediate severity by fractional flow reserve, instantaneous wave-free ratio, and non-invasive coronary flow reserve. International Journal of Cardiovascular Imaging, 2017, 33, 999-1007.	1.5	5
28	Assessment of left anterior descending artery stenosis of intermediate severity by fractional flow reserve, instantaneous wave-free ratio and non-invasive coronary flow reserve. Annales De Cardiologie Et D'Angeiologie, 2016, 65, 380-381.	0.6	0
29	Apical rotation, a simplified index of left ventricular twist is independently linked to recovery after acute anterior myocardial infarction. Annales De Cardiologie Et D'Angeiologie, 2016, 65, 381.	0.6	0