Ernesto Igartua

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8588004/publications.pdf Version: 2024-02-01

Version, 2024-02-01



#	Article	lF	CITATIONS
1	BARLEYMAP: physical and genetic mapping of nucleotide sequences and annotation of surrounding loci in barley. Molecular Breeding, 2015, 35, 1.	2.1	91
2	Genetic diversity of Prunus rootstocks analyzed by RAPD markers. Euphytica, 1999, 110, 139-149.	1.2	66
3	Adaptation of barley to mild winters: A role for PPDH2. BMC Plant Biology, 2011, 11, 164.	3.6	66
4	Population structure and marker–trait associations for pomological traits in peach and nectarine cultivars. Tree Genetics and Genomes, 2013, 9, 331-349.	1.6	65
5	Analysis of Plant Pan-Genomes and Transcriptomes with GET_HOMOLOGUES-EST, a Clustering Solution for Sequences of the Same Species. Frontiers in Plant Science, 2017, 8, 184.	3.6	63
6	Patterns of genetic and eco-geographical diversity in Spanish barleys. Theoretical and Applied Genetics, 2008, 116, 271-282.	3.6	62
7	Yield QTL affected by heading date in Mediterranean grown barley. Plant Breeding, 2009, 128, 46-53.	1.9	62
8	The Spanish barley core collection. Genetic Resources and Crop Evolution, 1998, 45, 475-481.	1.6	61
9	Heading date QTL in a springÂ×Âwinter barley cross evaluated in Mediterranean environments. Molecular Breeding, 2008, 21, 455-471.	2.1	58
10	Foliar fertilization of peach (Prunus persica (L.) Batsch) with different iron formulations: Effects on re-greening, iron concentration and mineral composition in treated and untreated leaf surfaces. Scientia Horticulturae, 2008, 117, 241-248.	3.6	57
11	Expression analysis of vernalization and day-length response genes in barley (Hordeum vulgare L.) indicates that VRNH2 is a repressor of PPDH2 (HvFT3) under long days. Journal of Experimental Botany, 2011, 62, 1939-1949.	4.8	57
12	Large Differences in Gene Expression Responses to Drought and Heat Stress between Elite Barley Cultivar Scarlett and a Spanish Landrace. Frontiers in Plant Science, 2017, 8, 647.	3.6	54
13	Changes over time in the adaptation of barley releases in north-eastern Spain. Plant Breeding, 1998, 117, 531-535.	1.9	53
14	Quantitative trait loci for agronomic traits in an elite barley population for Mediterranean conditions. Molecular Breeding, 2014, 33, 249-265.	2.1	52
15	Screening the Spanish Barley Core Collection for disease resistance. Plant Breeding, 2010, 129, 45-52.	1.9	51
16	Further evidence supporting Morocco as a centre of origin of barley. Theoretical and Applied Genetics, 1999, 98, 913-918.	3.6	49
17	Integrating statistical and ecophysiological analyses of genotype by environment interaction for grain filling of barley I Field Crops Research, 1999, 62, 63-74.	5.1	49
18	HvFT1 polymorphism and effectââ,¬â€survey of barley germplasm and expression analysis. Frontiers in Plant Science, 2014, 5, 251.	3.6	49

Ernesto Igartua

#	Article	IF	CITATIONS
19	Field responses of grain sorghum to a salinity gradient. Field Crops Research, 1995, 42, 15-25.	5.1	46
20	Spanish barley landraces outperform modern cultivars at lowâ€productivity sites. Plant Breeding, 2014, 133, 218-226.	1.9	44
21	HvFT1 (VrnH3) drives latitudinal adaptation in Spanish barleys. Theoretical and Applied Genetics, 2011, 122, 1293-1304.	3.6	43
22	Markerâ€Based Selection of QTL Affecting Grain and Malt Quality in Twoâ€Row Barley. Crop Science, 2000, 40, 1426-1433.	1.8	42
23	Major flowering time genes of barley: allelic diversity, effects, and comparison with wheat. Theoretical and Applied Genetics, 2021, 134, 1867-1897.	3.6	41
24	Prognosis of iron chlorosis from the mineral composition of Â⁻owers in peach. Journal of Horticultural Science and Biotechnology, 2000, 75, 111-118.	1.9	33
25	Assessing genetic and phenotypic diversity in pepper (Capsicum annuum L.) landraces from North-West Spain. Scientia Horticulturae, 2016, 203, 1-11.	3.6	33
26	Characterization and genetic control of germination-emergence responses of grain sorghum to salinity. Euphytica, 1994, 76, 185-193.	1.2	30
27	Fine mapping of the Rrs1 resistance locus against scald in two large populations derived from Spanish barley landraces. Theoretical and Applied Genetics, 2013, 126, 3091-3102.	3.6	30
28	Assessing different barley growth habits under Egyptian conditions for enhancing resilience to climate change. Field Crops Research, 2018, 224, 67-75.	5.1	30
29	Joint analysis for heading date QTL in small interconnected barley populations. Molecular Breeding, 2008, 21, 383-399.	2.1	29
30	Harnessing Novel Diversity From Landraces to Improve an Elite Barley Variety. Frontiers in Plant Science, 2019, 10, 434.	3.6	28
31	Fine mapping and comparative genomics integration of two quantitative trait loci controlling resistance to powdery mildew in a Spanish barley landrace. Theoretical and Applied Genetics, 2012, 124, 49-62.	3.6	25
32	Quantitative Trait Loci and Candidate Loci for Heading Date in a Large Population of a Wide Barley Cross. Crop Science, 2012, 52, 2469-2480.	1.8	24
33	Grain yield stability of high-yielding barley genotypes under Egyptian conditions for enhancing resilience to climate change. Crop and Pasture Science, 2018, 69, 681.	1.5	24
34	Choice of selection environment for improving crop yields in saline areas. Theoretical and Applied Genetics, 1995, 91-91, 1016-1021.	3.6	22
35	RFLP markers associated with major genes controlling heading date evaluated in a barley germ plasm pool. Heredity, 1999, 83, 551-559.	2.6	22
36	Development of a costâ€effective pyrosequencing approach for SNP genotyping in barley. Plant Breeding, 2011, 130, 394-397.	1.9	22

Ernesto Igartua

#	Article	IF	CITATIONS
37	Genetic association with highâ€resolution climate data reveals selection footprints in the genomes of barley landraces across the Iberian Peninsula. Molecular Ecology, 2019, 28, 1994-2012.	3.9	22
38	Mechanisms of Malt Extract Development in Barleys from Different European Regions: I. Effect of Environment and Grain Protein Content on Malt Extract Yield. Journal of the Institute of Brewing, 2000, 106, 111-116.	2.3	21
39	Rapid On-Site Phenotyping via Field Fluorimeter Detects Differences in Photosynthetic Performance in a Hybrid—Parent Barley Germplasm Set. Sensors, 2020, 20, 1486.	3.8	21
40	Genetic diversity of barley cultivars grown in Spain, estimated by RFLP, similarity and coancestry coefficients. Plant Breeding, 1998, 117, 429-435.	1.9	20
41	Identification of quantitative trait loci for resistance to powdery mildew in a Spanish barley landrace. Molecular Breeding, 2010, 25, 581-592.	2.1	20
42	Introgression of an intermediate VRNH1 allele in barley (Hordeum vulgare L) leads to reduced vernalization requirement without affecting freezing tolerance. Molecular Breeding, 2011, 28, 475-484.	2.1	20
43	Resistance to powdery mildew in Spanish barley landraces is controlled by different sets of quantitative trait loci. Theoretical and Applied Genetics, 2011, 123, 1019-1028.	3.6	19
44	Perspectives on Low Temperature Tolerance and Vernalization Sensitivity in Barley: Prospects for Facultative Growth Habit. Frontiers in Plant Science, 2020, 11, 585927.	3.6	19
45	Effects of Low Water Availability on Root Placement and Shoot Development in Landraces and Modern Barley Cultivars. Agronomy, 2020, 10, 134.	3.0	19
46	Responses to S1 Selection in Flint and Dent Synthetic Maize Populations. Crop Science, 1996, 36, 1129-1134.	1.8	18
47	Field responses of barley genotypes across a salinity gradient in an arid Mediterranean environment. Agricultural Water Management, 2021, 258, 107206.	5.6	18
48	Whole-genome analysis with SNPs from BOPA1 shows clearly defined groupings of Western Mediterranean, Ethiopian, and Fertile Crescent barleys. Genetic Resources and Crop Evolution, 2013, 60, 251-264.	1.6	15
49	Analysis of powdery mildew resistance in the Spanish barley core collection. Plant Breeding, 2011, 130, 195-202.	1.9	14
50	Developmental patterns of a large set of barley (<i>Hordeum vulgare</i>) cultivars in response to ambient temperature. Annals of Applied Biology, 2013, 162, 309-323.	2.5	14
51	Fine-tuning of the flowering time control in winter barley: the importance of HvOS2 and HvVRN2 in non-inductive conditions. BMC Plant Biology, 2019, 19, 113.	3.6	14
52	Towards Positional Isolation of Three Quantitative Trait Loci Conferring Resistance to Powdery Mildew in Two Spanish Barley Landraces. PLoS ONE, 2013, 8, e67336.	2.5	14
53	Prognosis of iron chlorosis in pear (Pyrus communis L.) and peach (Prunus persica L. Batsch) trees using bud, flower and leaf mineral concentrations. Plant and Soil, 2012, 354, 121-139.	3.7	13
54	A Cluster of Nucleotideâ€Binding Site–Leucineâ€Rich Repeat Genes Resides in a Barley Powdery Mildew Resistance Quantitative Trait Loci on 7HL. Plant Genome, 2016, 9, plantgenome2015.10.0101.	2.8	13

ERNESTO IGARTUA

#	Article	IF	CITATIONS
55	Resistance to powdery mildew in one Spanish barley landrace hardly resembles other previously identified wild barley resistances. European Journal of Plant Pathology, 2013, 136, 459-468.	1.7	12

 $_{56}$ Identification of quantitative trait loci for agronomic traits contributed by a barley (Hordeum) Tj ETQq0 0 0 rgBT /Oyerlock 10 Tf 50 702

57	Resequencing theVrs1 gene in Spanish barley landraces revealed reversion of six-rowed to two-rowed spike. Molecular Breeding, 2018, 38, 1.	2.1	10
58	Responses of Barley to High Ambient Temperature Are Modulated by Vernalization. Frontiers in Plant Science, 2021, 12, 776982.	3.6	10
59	TB1: from domestication gene to tool for many trades. Journal of Experimental Botany, 2020, 71, 4621-4624.	4.8	9
60	Durum Wheat Seminal Root Traits within Modern and Landrace Germplasm in Algeria. Agronomy, 2020, 10, 713.	3.0	9
61	Selection footprints in barley breeding lines detected by combining genotyping-by-sequencing with reference genome information. Molecular Breeding, 2015, 35, 1.	2.1	7
62	Evaluation of glycyrrhizin contents in licorice (Glycyrrhiza glabra L.) under drought and soil salinity conditions using nutrient concentrations and biochemical traits as biomarkers. Acta Physiologiae Plantarum, 2020, 42, 1.	2.1	7
63	Candidate genes underlying QTL for flowering time and their interactions in a wide spring barley (Hordeum vulgare L.) cross. Crop Journal, 2021, 9, 862-872.	5.2	6
64	Root Trait Diversity in Field Grown Durum Wheat and Comparison with Seedlings. Agronomy, 2021, 11, 2545.	3.0	6
65	Genetic diversity in developmental responses to light spectral quality in barley (Hordeum vulgare L.). BMC Plant Biology, 2020, 20, 207.	3.6	5
66	Genomic Prediction of Grain Yield in a Barley MAGIC Population Modeling Genotype per Environment Interaction. Frontiers in Plant Science, 2021, 12, 664148.	3.6	5
67	Rachis brittleness in a hybrid–parent barley (Hordeum vulgare) breeding germplasm with different combinations at the nonâ€brittle rachis genes. Plant Breeding, 2020, 139, 317-327.	1.9	3
68	Barley Types and Varieties in Spain: A Historical Overview. Ciencia E Investigacion Agraria, 2017, 44, 1-12.	0.2	3
69	Genome-wide association studies (GWAS) in barley. Burleigh Dodds Series in Agricultural Science, 2019, , 503-536.	0.2	2
70	Algerian durum wheat assessment for early drought tolerance shows landraces superiority. Egyptian Journal of Agronomy, 2019, .	0.3	2
71	Hybrids Provide More Options for Fine-Tuning Flowering Time Responses of Winter Barley. Frontiers in Plant Science, 2022, 13, 827701.	3.6	1
72	Registration of Four Sorghum Germplasm Randomâ€Mating Populations. Crop Science, 1997, 37, 1036-1037.	1.8	0

#	Article	IF	CITATIONS
73	Barley Adaptation: Teachings from Landraces Will Help to Respond to Climate Change. , 2013, , 327-337.		0