

# Renato Toffanin

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8603356/publications.pdf>

Version: 2024-02-01

65  
papers

1,987  
citations

257450

24  
h-index

243625

44  
g-index

65  
all docs

65  
docs citations

65  
times ranked

2218  
citing authors

#	ARTICLE	IF	CITATIONS
1	Implementing Systemic Innovation Strategies for a More Sustainable Future: The Case of Three Overseas Countries and Territories. <i>Frontiers in Research Metrics and Analytics</i> , 2021, 6, 801789.	1.9	0
2	Innovation system foresight and systemic innovation for the overseas countries and territories. <i>Foresight</i> , 2018, 20, 105-122.	2.1	2
3	Exploitation of Old Wheat Properties for Prevention of Human Disease. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.5	1
4	Fast MRI Methods for the Clinical Evaluation of Skeletal Disorders. , 2011, , .		0
5	3 T magnetic resonance imaging of the musculoskeletal system. <i>Radiologia Medica</i> , 2010, 115, 571-584.	7.7	9
6	Fast T2 mapping of the patellar articular cartilage with gradient and spin-echo magnetic resonance imaging at 1.5 T: validation and initial clinical experience in patients with osteoarthritis. <i>Skeletal Radiology</i> , 2008, 37, 511-517.	2.0	30
7	Mechanical and diffusive properties of homogeneous alginate gels in form of particles and cylinders. <i>Journal of Biomedical Materials Research - Part A</i> , 2008, 87A, 808-818.	4.0	22
8	Chondrocyte-alginate bioconstructs: An nuclear magnetic resonance relaxation study. <i>Journal of Biomedical Materials Research - Part A</i> , 2007, 83A, 345-353.	4.0	6
9	MEDICAL IMAGING ANALYSIS OF THE THREE DIMENSIONAL (3D) ARCHITECTURE OF TRABECULAR BONE: TECHNIQUES AND THEIR APPLICATIONS. , 2005, , 1-41.		3
10	Degenerative Changes of Porcine Intervertebral Disc Induced by Vertebral Endplate Injuries. <i>Spine</i> , 2005, 30, 174-180.	2.0	84
11	Structure of the exopolysaccharide produced by <i>Enterobacter amnigenus</i> . <i>Carbohydrate Research</i> , 2005, 340, 439-447.	2.3	40
12	Transverse relaxation mechanisms in articular cartilage. <i>Journal of Magnetic Resonance</i> , 2004, 169, 300-307.	2.1	127
13	Gel beads from novel ionic polysaccharides. <i>Carbohydrate Polymers</i> , 2004, 55, 163-169.	10.2	6
14	Numerical simulation of trabecular bone magnetic resonance imaging. , 2004, 2004, 1088-91.		0
15	Ex vivo assessment of trabecular bone structure from three-dimensional projection reconstruction mr micro-images. <i>IEEE Transactions on Biomedical Engineering</i> , 2003, 50, 967-977.	4.2	9
16	NOVEL BIOMATERIALS BASED ON CROSS-LINKED HYALURONAN: STRUCTURAL INVESTIGATIONS. , 2002, , 269-276.		0
17	Structural investigations of cross-linked hyaluronan. <i>Biomaterials</i> , 2002, 23, 1161-1167.	11.4	24
18	MR microscopy of hyaline cartilage: current status. <i>European Radiology</i> , 2002, 12, 814-823.	4.5	38

#	ARTICLE	IF	CITATIONS
19	Magnetic resonance microscopy for the quantitative analysis of trabecular bone architecture. <i>Journal of Gravitational Physiology: A Journal of the International Society for Gravitational Physiology</i> , 2002, 9, P173-4.	0.0	0
20	Proteoglycan Depletion and Magnetic Resonance Parameters of Articular Cartilage. <i>Archives of Biochemistry and Biophysics</i> , 2001, 390, 235-242.	3.0	57
21	Short-TE projection reconstruction NMR microscopy of trabecular bone. <i>Magnetic Resonance Imaging</i> , 2001, 19, 485-486.	1.8	3
22	Magnetic resonance microscopy of osteoporotic bone. <i>AIP Conference Proceedings</i> , 2000, , .	0.4	2
23	Inhomogeneous alginate gel spheres: An assessment of the polymer gradients by synchrotron radiation-induced x-ray emission, magnetic resonance microimaging, and mathematical modeling. <i>Biopolymers</i> , 2000, 53, 60-71.	2.4	126
24	Collagen fibrils are differently organized in weight-bearing and not-weight-bearing regions of pig articular cartilage. <i>The Journal of Experimental Zoology</i> , 2000, 287, 346-352.	1.4	24
25	Articular cartilage repair in rabbits by using suspensions of allogenic chondrocytes in alginate. <i>Biomaterials</i> , 2000, 21, 795-801.	11.4	176
26	Short-TE projection reconstruction MR microscopy in the evaluation of articular cartilage thickness. <i>European Radiology</i> , 2000, 10, 1222-1226.	4.5	10
27	Collagen fibrils are differently organized in weight-bearing and not-weight-bearing regions of pig articular cartilage. <i>The Journal of Experimental Zoology</i> , 2000, 287, 346-352.	1.4	0
28	Polysaccharides from hot water extracts of roasted <i>Coffea arabica</i> beans: isolation and characterization. <i>Carbohydrate Polymers</i> , 1999, 40, 71-81.	10.2	177
29	Structural determination of the acidic exopolysaccharide produced by a <i>Pseudomonas</i> sp. strain 1.15. <i>Carbohydrate Research</i> , 1999, 315, 159-168.	2.3	49
30	High-resolution $^1\text{H}$ NMR investigation of coffee. <i>Journal of the Science of Food and Agriculture</i> , 1999, 79, 869-878.	3.5	59
31	A better understanding of the properties of alginate solutions and gels by quantitative magnetic resonance imaging (MRI). <i>Carbohydrate Research</i> , 1998, 306, 19-26.	2.3	20
32	Cell wall polysaccharides from <i>Gelidium</i> species: physico-chemical studies using MRI techniques. <i>Journal of Applied Phycology</i> , 1998, 10, 315-322.	2.8	12
33	Magnetic resonance imaging of articular cartilage: ex vivo study on normal cartilage correlated with magnetic resonance microscopy. <i>European Radiology</i> , 1998, 8, 1130-1136.	4.5	28
34	Correlation between biochemical composition and magnetic resonance appearance of articular cartilage. <i>Osteoarthritis and Cartilage</i> , 1998, 6, 24-32.	1.3	113
35	Structural investigation of the exopolysaccharide produced by <i>Pseudomonas flavescens</i> strain B62 . Degradation by a fungal cellulase and isolation of the oligosaccharide repeating unit. <i>FEBS Journal</i> , 1998, 251, 971-979.	0.2	20
36	$^31\text{P}$ NMR analysis of phospholipids in crude extracts from different sources: improved efficiency of the solvent system. , 1998, 36, 907-912.		27

#	ARTICLE	IF	CITATIONS
37	Sensitivity of chondrocytes of growing cartilage to reactive oxygen species. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 1998, 1425, 103-111.	2.4	21
38	<sup>31</sup> P NMR analysis of phospholipids in crude extracts from different sources: improved efficiency of the solvent system. <i>Magnetic Resonance in Chemistry</i> , 1998, 36, 907-912.	1.9	2
39	Conformational analysis on segments of charged polysaccharides. The case of hyaluronic acid dimer and chondrosine. <i>Computational and Theoretical Chemistry</i> , 1997, 395-396, 437-449.	1.5	9
40	Investigation of the carrageenans extracted from <i>Solieria filiformis</i> and <i>Agardhiella subulata</i> from Mar Piccolo, Taranto. <i>Marine Chemistry</i> , 1997, 58, 319-325.	2.3	42
41	Organic Solvent Systems for <sup>31</sup> P Nuclear Magnetic Resonance Analysis of Lecithin Phospholipids: Applications to Two-Dimensional Gradient-Enhanced <sup>1</sup> H-Detected Heteronuclear Multiple Quantum Coherence Experiments. <i>Analytical Biochemistry</i> , 1997, 245, 38-47.	2.4	64
42	Hydrogen-bonding and conformation of agarose in methyl sulfoxide and aqueous solutions investigated by <sup>1</sup> H and <sup>13</sup> C NMR spectroscopy. <i>Carbohydrate Research</i> , 1997, 304, 293-302.	2.3	27
43	Analysis of Mixtures of Phospholipids in Dimethylformamide. , 1997, , 333-334.		0
44	A method for generating magnetic resonance microimaging $T_2$ maps with low sensitivity to diffusion. <i>Magnetic Resonance in Medicine</i> , 1996, 35, 423-425.	3.0	29
45	Investigation of laminar appearance of articular cartilage by means of magnetic resonance microscopy. <i>Magnetic Resonance Imaging</i> , 1996, 14, 435-442.	1.8	98
46	Structure and properties of agar from two unexploited agarophytes from Venezuela. <i>Hydrobiologia</i> , 1996, 326-327, 497-500.	2.0	12
47	Structure and properties of agar from two unexploited agarophytes from Venezuela. , 1996, , 497-500.		2
48	Detection and quantitation of phosphorus metabolites in crude tissue extracts by <sup>1</sup> H and <sup>31</sup> P NMR: use of gradient assisted <sup>1</sup> H- <sup>31</sup> P HMQC experiments, with selective pulses, for the assignment of less abundant metabolites. <i>NMR in Biomedicine</i> , 1995, 8, 190-196.	2.8	12
49	Modified procedures for extraction and analysis of carrageenan applied to the red alga <i>Hypnea musciformis</i> . <i>Journal of Applied Phycology</i> , 1995, 7, 565-576.	2.8	50
50	Sensitivity of hypertrophic chondrocytes to oxygen derived free radicals.. <i>Bone</i> , 1995, 17, 595.	2.9	0
51	NMR analysis of succinoglycans from different microbial sources: partial assignment of their <sup>1</sup> H and <sup>13</sup> C NMR spectra and location of the succinate and the acetate groups. <i>Carbohydrate Research</i> , 1994, 265, 167-179.	2.3	36
52	A conformational study of the Smith degradation product of the <i>Klebsiella</i> K40 capsular polysaccharide by 1D NOESY and molecular mechanics calculations. <i>Carbohydrate Research</i> , 1994, 265, 151-159.	2.3	3
53	Detection of cellulose in the cell wall of some red algae by <sup>13</sup> C NMR spectroscopy. <i>Carbohydrate Research</i> , 1994, 262, 167-171.	2.3	12
54	Analysis of Lipids in Crude Extracts by <sup>13</sup> C Nuclear Magnetic Resonance. <i>Analytical Biochemistry</i> , 1993, 214, 238-244.	2.4	12

#	ARTICLE	IF	CITATIONS
55	Structural determination of the capsular polysaccharide produced by <i>Klebsiella pneumoniae</i> serotype K40. NMR studies of the oligosaccharide obtained upon depolymerisation of the polysaccharide with a bacteriophage-associated endoglycanase. <i>FEBS Journal</i> , 1993, 213, 445-453.	0.2	20
56	Evaluation of steam explosion as pretreatment in agar extraction from <i>Gracilaria dura</i> (C. Agardh) J. Agardh (Gracilariaceae, Rhodophyta). <i>Journal of Applied Phycology</i> , 1993, 5, 417-424.	2.8	22
57	NMR studies of oligosaccharides derived from hyaluronate: complete assignment of <sup>1</sup> H and <sup>13</sup> C NMR spectra of aqueous di- and tetra-saccharides, and comparison of chemical shifts for oligosaccharides of increasing degree of polymerisation. <i>Carbohydrate Research</i> , 1993, 245, 113-120.	2.3	45
58	<sup>1</sup> H-NMR Spectroscopic Studies of Lipid Extracts from Human Fatty Liver. <i>Biochemical and Biophysical Research Communications</i> , 1993, 192, 1217-1222.	2.1	19
59	Pyruvate-Rich Agarose from the Red Alga <i>Gracilaria dura</i> . <i>Planta Medica</i> , 1992, 58, 588-589.	1.3	2
60	<sup>1</sup> H- and <sup>13</sup> C-NMR studies of solutions of hyaluronic acid esters and salts in methyl sulfoxide: comparison of hydrogen-bond patterns and conformational behaviour. <i>Carbohydrate Research</i> , 1992, 230, 1-13.	2.3	38
61	Chemical and macromolecular characterisation of agar polymers from <i>Gracilaria dura</i> (C. Agardh) J. Agardh (Gracilariaceae, Rhodophyta). <i>Carbohydrate Polymers</i> , 1992, 18, 171-178.	10.2	69
62	Lipid extracts from different algal species: <sup>1</sup> H and <sup>13</sup> C-NMR spectroscopic studies as a new tool to screen differences in the composition of fatty acids, sterols and carotenoids. <i>Journal of Applied Phycology</i> , 1992, 4, 315-322.	2.8	18
63	<sup>1</sup> H- and <sup>13</sup> C-NMR spectroscopic studies of lipid extracts of the red alga <i>Gracilaria longa</i> . <i>Journal of Applied Phycology</i> , 1992, 4, 149-155.	2.8	6
64	Evidence for a boat-chair equilibrium in the glucuronate residue of chondrosine. <i>Carbohydrate Research</i> , 1991, 209, C13-C15.	2.3	13
65	Quantification of trabecular bone structure from three-dimensional <sup>1</sup> H MR images. , 0, , .		0