

Saswata Talukdar

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8637225/publications.pdf>

Version: 2024-02-01

27
papers

6,594
citations

257450

24
h-index

526287

27
g-index

27
all docs

27
docs citations

27
times ranked

13303
citing authors

#	ARTICLE	IF	CITATIONS
1	Positive Reinforcing Mechanisms between GPR120 and PPAR β Modulate Insulin Sensitivity. <i>Cell Metabolism</i> , 2020, 31, 1173-1188.e5.	16.2	43
2	Free fatty acid receptor 4 activation protects against choroidal neovascularization in mice. <i>Angiogenesis</i> , 2020, 23, 385-394.	7.2	17
3	Insulin resistance drives hepatic de novo lipogenesis in nonalcoholic fatty liver disease. <i>Journal of Clinical Investigation</i> , 2020, 130, 1453-1460.	8.2	362
4	Once-a-weekly administration of a long-acting fibroblast growth factor 21 analogue modulates lipids, bone turnover markers, blood pressure and body weight differently in obese people with hypertriglyceridaemia and in non-human primates. <i>Diabetes, Obesity and Metabolism</i> , 2017, 19, 1762-1772.	4.4	106
5	A Long-Acting FGF21 Molecule, PF-05231023, Decreases Body Weight and Improves Lipid Profile in Non-human Primates and Type 2 Diabetic Subjects. <i>Cell Metabolism</i> , 2016, 23, 427-440.	16.2	377
6	FGF21 Regulates Sweet and Alcohol Preference. <i>Cell Metabolism</i> , 2016, 23, 344-349.	16.2	259
7	LTB4 promotes insulin resistance in obese mice by acting on macrophages, hepatocytes and myocytes. <i>Nature Medicine</i> , 2015, 21, 239-247.	30.7	252
8	FGF21 does not require interscapular brown adipose tissue and improves liver metabolic profile in animal models of obesity and insulin-resistance. <i>Scientific Reports</i> , 2015, 5, 11382.	3.3	45
9	Pharmacokinetics (PK), Pharmacodynamics (PD) and Integrated PK/PD Modeling of a Novel Long Acting FGF21 Clinical Candidate PF-05231023 in Diet-Induced Obese and Leptin-Deficient Obese Mice. <i>PLoS ONE</i> , 2015, 10, e0119104.	2.5	55
10	Knock-Down of IL-1Ra in Obese Mice Decreases Liver Inflammation and Improves Insulin Sensitivity. <i>PLoS ONE</i> , 2014, 9, e107487.	2.5	20
11	Fibroblast Growth Factor 21 Improves Insulin Sensitivity and Synergizes with Insulin in Human Adipose Stem Cell-Derived (hASC) Adipocytes. <i>PLoS ONE</i> , 2014, 9, e111767.	2.5	28
12	NCoR Repression of LXRs Restricts Macrophage Biosynthesis of Insulin-Sensitizing Omega 3 Fatty Acids. <i>Cell</i> , 2013, 155, 200-214.	28.9	149
13	Neuronal Sirt1 Deficiency Increases Insulin Sensitivity in Both Brain and Peripheral Tissues. <i>Journal of Biological Chemistry</i> , 2013, 288, 10722-10735.	3.4	50
14	Development of a Novel Long-Acting Antidiabetic FGF21 Mimetic by Targeted Conjugation to a Scaffold Antibody. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2013, 346, 270-280.	2.5	105
15	GPR105 Ablation Prevents Inflammation and Improves Insulin Sensitivity in Mice with Diet-Induced Obesity. <i>Journal of Immunology</i> , 2012, 189, 1992-1999.	0.8	65
16	G protein-coupled receptor 21 deletion improves insulin sensitivity in diet-induced obese mice. <i>Journal of Clinical Investigation</i> , 2012, 122, 2444-2453.	8.2	49
17	Maintenance of Metabolic Homeostasis by Sestrin2 and Sestrin3. <i>Cell Metabolism</i> , 2012, 16, 311-321.	16.2	242
18	Increased Macrophage Migration Into Adipose Tissue in Obese Mice. <i>Diabetes</i> , 2012, 61, 346-354.	0.6	304

#	ARTICLE	IF	CITATIONS
19	Neutrophils mediate insulin resistance in mice fed a high-fat diet through secreted elastase. <i>Nature Medicine</i> , 2012, 18, 1407-1412.	30.7	751
20	Inflammation Is Necessary for Long-Term but Not Short-Term High-Fat Diet-Induced Insulin Resistance. <i>Diabetes</i> , 2011, 60, 2474-2483.	0.6	452
21	Adipocyte NCoR Knockout Decreases PPAR β Phosphorylation and Enhances PPAR β Activity and Insulin Sensitivity. <i>Cell</i> , 2011, 147, 815-826.	28.9	246
22	Targeting GPR120 and other fatty acid-sensing GPCRs ameliorates insulin resistance and inflammatory diseases. <i>Trends in Pharmacological Sciences</i> , 2011, 32, 543-550.	8.7	218
23	Brain PPAR β promotes obesity and is required for the insulin-sensitizing effect of thiazolidinediones. <i>Nature Medicine</i> , 2011, 17, 618-622.	30.7	214
24	GPR120 Is an Omega-3 Fatty Acid Receptor Mediating Potent Anti-inflammatory and Insulin-Sensitizing Effects. <i>Cell</i> , 2010, 142, 687-698.	28.9	2,013
25	Osteopontin Is Required for the Early Onset of High Fat Diet-Induced Insulin Resistance in Mice. <i>PLoS ONE</i> , 2010, 5, e13959.	2.5	71
26	Chenodeoxycholic acid suppresses the activation of acetyl-coenzyme A carboxylase- β gene transcription by the liver X receptor agonist T0-901317. <i>Journal of Lipid Research</i> , 2007, 48, 2647-2663.	4.2	13
27	The mechanism mediating the activation of acetyl-coenzyme A carboxylase- β gene transcription by the liver X receptor agonist T0-901317. <i>Journal of Lipid Research</i> , 2006, 47, 2451-2461.	4.2	88