

Roberto A Perego

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8642707/publications.pdf>

Version: 2024-02-01

23
papers

891
citations

623734

14
h-index

677142

22
g-index

25
all docs

25
docs citations

25
times ranked

1328
citing authors

#	ARTICLE	IF	CITATIONS
1	DNA Damage in Circulating Hematopoietic Progenitor Stem Cells as Promising Biological Sensor of Frailty. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2022, 77, 1279-1286.	3.6	5
2	PKH ^{high} /CD133 ⁺ /CD24 ^{low} Renal Stem-Like Cells Isolated from Human Nephrospheres Exhibit In Vitro Multipotency. <i>Cells</i> , 2020, 9, 1805.	4.1	4
3	36-kDa Annexin A3 Isoform Negatively Modulates Lipid Storage in Clear Cell Renal Cell Carcinoma Cells. <i>American Journal of Pathology</i> , 2020, 190, 2317-2326.	3.8	53
4	Integration of Lipidomics and Transcriptomics Reveals Reprogramming of the Lipid Metabolism and Composition in Clear Cell Renal Cell Carcinoma. <i>Metabolites</i> , 2020, 10, 509.	2.9	51
5	<i>De novo</i> UBE2A mutations are recurrently acquired during chronic myeloid leukemia progression and interfere with myeloid differentiation pathways. <i>Haematologica</i> , 2019, 104, 1789-1797.	3.5	21
6	The 1ALCTL and 1BLCTL isoforms of Arg/Abl2 induce fibroblast activation and extra cellular matrix remodelling differently. <i>Biology Open</i> , 2019, 8, .	1.2	7
7	CORRELATION BETWEEN FRAILTY AND DNA DAMAGE IN HEMATOPOIETIC STEM CELLS: A PILOT STUDY. <i>Innovation in Aging</i> , 2019, 3, S87-S87.	0.1	1
8	Hepcidin regulation in a mouse model of acute hypoxia. <i>European Journal of Haematology</i> , 2018, 100, 636-643.	2.2	17
9	Nephrosphere-Derived Cells Are Induced to Multilineage Differentiation when Cultured on Human Decellularized Kidney Scaffolds. <i>American Journal of Pathology</i> , 2018, 188, 184-195.	3.8	25
10	Integrated multi-omics characterization reveals a distinctive metabolic signature and the role of NDUFA4L2 in promoting angiogenesis, chemoresistance, and mitochondrial dysfunction in clear cell renal cell carcinoma. <i>Aging</i> , 2018, 10, 3957-3985.	3.1	133
11	The glucose and lipid metabolism reprogramming is grade-dependent in clear cell renal cell carcinoma primary cultures and is targetable to modulate cell viability and proliferation. <i>Oncotarget</i> , 2017, 8, 113502-113515.	1.8	95
12	Major Action of Endogenous Lysyl Oxidase in Clear Cell Renal Cell Carcinoma Progression and Collagen Stiffness Revealed by Primary Cell Cultures. <i>American Journal of Pathology</i> , 2016, 186, 2473-2485.	3.8	36
13	Arg tyrosine kinase modulates TGF- β 1 production in human renal tubular cells under high-glucose conditions. <i>Journal of Cell Science</i> , 2016, 129, 2925-2936.	2.0	11
14	Grade-Dependent Metabolic Reprogramming in Kidney Cancer Revealed by Combined Proteomics and Metabolomics Analysis. <i>Cancer Research</i> , 2015, 75, 2541-2552.	0.9	236
15	Response to communication of Paola Romagnani and Giuseppe Remuzzi. <i>Stem Cell Research</i> , 2014, 12, 830-831.	0.7	0
16	PKH ^{high} cells within clonal human nephrospheres provide a purified adult renal stem cell population. <i>Stem Cell Research</i> , 2013, 11, 1163-1177.	0.7	29
17	Renal cell carcinoma primary cultures maintain genomic and phenotypic profile of parental tumor tissues. <i>BMC Cancer</i> , 2011, 11, 244.	2.6	24
18	Primary Cell Cultures from Human Renal Cortex and Renal-Cell Carcinoma Evidence a Differential Expression of Two Spliced Isoforms of Annexin A3. <i>American Journal of Pathology</i> , 2010, 176, 1660-1670.	3.8	44

#	ARTICLE	IF	CITATIONS
19	Eight full-length abelson related gene (Arg) isoforms are constitutively expressed in caki cell line and cell distribution of two isoforms has been analyzed after transfection. <i>Journal of Cellular Biochemistry</i> , 2008, 105, 1219-1227.	2.6	10
20	Concentration and microsatellite status of plasma DNA for monitoring patients with renal carcinoma. <i>European Journal of Cancer</i> , 2008, 44, 1039-1047.	2.8	32
21	Primary Cell Cultures Arising from Normal Kidney and Renal Cell Carcinoma Retain the Proteomic Profile of Corresponding Tissues. <i>Journal of Proteome Research</i> , 2005, 4, 1503-1510.	3.7	38
22	The role of quantitative polymerase chain reaction in the management of follicular lymphoma patients. <i>Tumori</i> , 2005, 91, 59-66.	1.1	1
23	The expression of the non-receptor tyrosine kinases Arg and c-abl is differently modulated in B lymphoid cells at different stages of differentiation. <i>FEBS Letters</i> , 2002, 527, 216-222.	2.8	10