

Bao-Kai Cui

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8645800/publications.pdf>

Version: 2024-02-01

146
papers

5,449
citations

117625
34
h-index

106344
65
g-index

148
all docs

148
docs citations

148
times ranked

3582
citing authors

#	ARTICLE	IF	CITATIONS
1	Diversity of Microbial Communities of <i>Pinus sylvestris</i> var. <i>mongolica</i> at Spatial Scale. <i>Microorganisms</i> , 2022, 10, 371.	3.6	8
2	Phylogenomics and Comparative Genomics Highlight Specific Genetic Features in <i>Ganoderma</i> Species. <i>Journal of Fungi</i> (Basel, Switzerland), 2022, 8, 311.	3.5	10
3	Species Diversity, Molecular Phylogeny, and Ecological Habits of <i>Fomitopsis</i> (Polyporales). <i>Tj ETQq1 1 0.784314 rgBT/Overlock 10 Tf</i> 50	3.5	7
4	A Phylogenetic and Taxonomic Study on <i>Phellodon</i> (Bankeraceae, Thelephorales) from China. <i>Journal of Fungi</i> (Basel, Switzerland), 2022, 8, 429.	3.5	6
5	Species Diversity and Molecular Phylogeny of <i>Cyanosporus</i> (Polyporales, Basidiomycota). <i>Frontiers in Microbiology</i> , 2021, 12, 631166.	3.5	16
6	Taxonomy and Phylogeny of the <i>Fomitopsis pinicola</i> Complex With Descriptions of Six New Species From East Asia. <i>Frontiers in Microbiology</i> , 2021, 12, 644979.	3.5	24
7	Introducing a Thermo-Alkali-Stable, Metallic Ion-Tolerant Laccase Purified From White Rot Fungus <i>Trametes hirsuta</i> . <i>Frontiers in Microbiology</i> , 2021, 12, 670163.	3.5	16
8	Taxonomy and Molecular Phylogeny of <i>Phellodon</i> (Thelephorales) with Descriptions of Four New Species from Southwest China. <i>Forests</i> , 2021, 12, 932.	2.1	4
9	Selection of a pH- and temperature-stable laccase from <i>Ganoderma australe</i> and its application for bioremediation of textile dyes. <i>Journal of Environmental Management</i> , 2021, 299, 113619.	7.8	18
10	Fungal diversity notes 1277–1386: taxonomic and phylogenetic contributions to fungal taxa. <i>Fungal Diversity</i> , 2020, 104, 1-266.	12.3	60
11	Two new species of <i>Neofavolus</i> (Polyporales, Basidiomycota) based on morphological characters and molecular evidence. <i>Mycological Progress</i> , 2020, 19, 471-480.	1.4	1
12	FungalTraits: a user-friendly traits database of fungi and fungus-like stramenopiles. <i>Fungal Diversity</i> , 2020, 105, 1-16.	12.3	387
13	Morphological characters and molecular data reveal three new species of <i>Fomitopsis</i> (Basidiomycota). <i>Mycological Progress</i> , 2019, 18, 1317-1327.	1.4	8
14	Species diversity, taxonomy and phylogeny of Polyporaceae (Basidiomycota) in China. <i>Fungal Diversity</i> , 2019, 97, 137-392.	12.3	111
15	Two new <i>Neofomitella</i> species (Polyporaceae, Basidiomycota) based on morphological and molecular evidence. <i>Mycological Progress</i> , 2019, 18, 593-602.	1.4	6
16	Species Diversity, Phylogeny, Divergence Time, and Biogeography of the Genus <i>Sanghuangporus</i> (Basidiomycota). <i>Frontiers in Microbiology</i> , 2019, 10, 812.	3.5	52
17	Notes, outline and divergence times of Basidiomycota. <i>Fungal Diversity</i> , 2019, 99, 105-367.	12.3	256
18	<p>Morphological and phylogenetic analyses reveal a new species of Fistulina (Fistulinaceae, Agaricales) from Australia</p>. <i>Phytotaxa</i> , 2019, 420, 233-240.	0.3	3

#	ARTICLE	IF	CITATIONS
19	Medium composition optimization, structural characterization, and antioxidant activity of exopolysaccharides from the medicinal mushroom <i>Ganoderma lingzhi</i> . International Journal of Biological Macromolecules, 2019, 124, 1186-1196.	7.5	35
20	<i>Sanghuangporus toxicodendri</i> sp. nov. (Hymenochaetales, Basidiomycota) from China. MycoKeys, 2019, 57, 101-111.	1.9	11
21	Chitosan crosslinked with genipin as supporting matrix for biodegradation of synthetic dyes: Laccase immobilization and characterization. Chemical Engineering Research and Design, 2018, 132, 664-676.	5.6	94
22	<i>Rigidotubus tephroleucus</i> gen. et sp. nov. (Cystostereaceae, Agaricales) evidenced by morphological characters and phylogenetic analyses. Phytotaxa, 2018, 333, 259.	0.3	4
23	Morphological characters and phylogenetic analysis reveal a new species of <i>Phellinus</i> with hooked hymenial setae from Vietnam. Phytotaxa, 2018, 356, 91.	0.3	6
24	Morphological and molecular identification of two new <i>Ganoderma</i> species on <i>Casuarina equisetifolia</i> from China. MycoKeys, 2018, 34, 93-108.	1.9	43
25	Phylogeny and taxonomy of <i>Laetiporus</i> (Basidiomycota, Polyporales) with descriptions of two new species from western China. MycoKeys, 2018, 37, 57-71.	1.9	23
26	Species diversity and molecular systematics of <i>Fibroporia</i> (Polyporales, Basidiomycota) and its related genera. Mycological Progress, 2017, 16, 521-533.	1.4	19
27	Multi-locus phylogeny and morphology reveal five new species of <i>Fomitiporia</i> (Hymenochaetaceae) from China. Mycological Progress, 2017, 16, 687-701.	1.4	14
28	A novel laccase from white rot fungus <i>Trametes orientalis</i> : Purification, characterization, and application. International Journal of Biological Macromolecules, 2017, 102, 758-770.	7.5	63
29	A six-gene phylogenetic overview of Basidiomycota and allied phyla with estimated divergence times of higher taxa and a phyloproteomics perspective. Fungal Diversity, 2017, 84, 43-74.	12.3	124
30	Phylogeny and taxonomy of <i>Favolus</i> (Basidiomycota). Mycologia, 2017, 109, 1-14.	1.9	21
31	Three new species of <i>Phylloporia</i> (Hymenochaetales) with dimitic hyphal systems from tropical China. Mycologia, 2017, 109, 951-964.	1.9	8
32	Phylogeny, divergence time and historical biogeography of <i>Laetiporus</i> (Basidiomycota, Polyporales). BMC Evolutionary Biology, 2017, 17, 102.	3.2	53
33	Morphological characters and phylogenetic analysis reveal a new species of <i>Sanghuangporus</i> from China. Phytotaxa, 2017, 311, 270.	0.3	9
34	<i>Neoalbatrellus odorus</i> sp. nov. (Albatrellaceae, Russulales) from Southwest China. Phytotaxa, 2017, 309, 217.	0.3	5
35	Laccase Production Among Medicinal Mushrooms from the Genus <i>Flammulina</i> (Agaricomycetes) Under Different Treatments in Submerged Fermentation. International Journal of Medicinal Mushrooms, 2016, 18, 1049-1059.	1.5	5
36	Sequential Solid-State and Submerged Cultivation of the White Rot Fungus <i>Pleurotus ostreatus</i> on Biomass and the Activity of Lignocellulolytic Enzymes. BioResources, 2016, 11, .	1.0	18

#	ARTICLE	IF	CITATIONS
37	Global diversity and molecular systematics of <i>Wrightoporia</i> s.l. (<i>Russulales</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 59	4.4	50
38	Taxonomy and multi-gene phylogeny of Haploporus (Polyporales, Basidiomycota). Mycological Progress, 2016, 15, 731-742.	1.4	14
39	<p align="left">Molecular phylogeny and morphology reveal a new species of Amauroderma (Basidiomycota) from China. Phytotaxa, 2016, 260, 47.	0.3	16
40	Phylogeny and biogeography of the remarkable genus Bondarzewia (Basidiomycota, Russulales). Scientific Reports, 2016, 6, 34568.	3.3	31
41	Fungal diversity notes 253â€“366: taxonomic and phylogenetic contributions to fungal taxa. Fungal Diversity, 2016, 78, 1-237.	12.3	239
42	Taxonomy and phylogeny of the brown-rot fungi: Fomitopsis and its related genera. Fungal Diversity, 2016, 80, 343-373.	12.3	101
43	Molecular phylogeny and global diversity of the remarkable genus <i>Bondarzewia</i> (Basidiomycota,) Tj ETQq1 1 0.784314 rgBT /Over	1.9	15
44	Fungal diversity notes 367â€“490: taxonomic and phylogenetic contributions to fungal taxa. Fungal Diversity, 2016, 80, 1-270.	12.3	314
45	Two new species of <i>Fomitiporia</i> (Hymenochaetales, Basidiomycota) from Tibet, southwest China. Mycologia, 2016, 108, 1010-1017.	1.9	11
46	Morphological characters and phylogenetic analysis reveal a new species within the <i>Ganoderma lucidum</i> complex from South Africa. Phytotaxa, 2016, 266, 115.	0.3	31
47	Podoserpula ailaoshanensis sp. nov. (Amylocorticiales, Basidiomycota) from China based on morphological and sequence analyses. Mycoscience, 2016, 57, 295-301.	0.8	3
48	Polypore diversity in North America with an annotated checklist. Mycological Progress, 2016, 15, 771-790.	1.4	22
49	Phylogenetic analysis and taxonomy of the <i>Antrodia heteromorpha</i> complex in China. Mycoscience, 2016, 57, 1-10.	0.8	23
50	Immobilization of laccase onto chitosan beads to enhance its capability to degrade synthetic dyes. International Biodeterioration and Biodegradation, 2016, 110, 69-78.	3.9	166
51	Phylogeny and taxonomy of the genus <i>Anomoloma</i> (Amylocorticiales, Basidiomycota). Mycological Progress, 2016, 15, 1.	1.4	9
52	Biosorption performances of raw and chemically modified biomasses from <i>Perenniporia subacida</i> for heterocycle dye Neutral Red. Desalination and Water Treatment, 2016, 57, 8454-8469.	1.0	5
53	Taxonomy and Phylogeny of Polyporus Group Melanopus (Polyporales, Basidiomycota) from China. PLoS ONE, 2016, 11, e0159495.	2.5	32
54	Genetic Diversity and Relationships of 24 Strains of Genus <i>Auricularia</i> (Agaricomycetes) Assessed Using SRAP Markers. International Journal of Medicinal Mushrooms, 2016, 18, 945-954.	1.5	1

#	ARTICLE	IF	CITATIONS
55	Molecular phylogeny and taxonomy of Fibroporia (Basidiomycota) in China. <i>Phytotaxa</i> , 2015, 203, 47.	0.3	15
56	Daedalea americana sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analysis. <i>Phytotaxa</i> , 2015, 204, 277.	0.3	18
57	Morphological characters and molecular data reveal a new species of Hydnocristella (Gomphales.) Tj ETQq1 1 0.784314 rgBT ₅ /Overlock	0.4	
58	Phylogeny, divergence time estimation, and biogeography of the genus Heterobasidion (Basidiomycota,) Tj ETQq0 0.0 rgBT _{12.3} /Overlock 10	0.3	
59	Morphological characters and molecular data reveal two new species of Postia (Basidiomycota) from China. <i>Mycological Progress</i> , 2015, 14, 1.	1.4	10
60	A Novel Phellinidium sp. Causes Laminated Root Rot on Qilian Juniper (<i>Sabina przewalskii</i>) in Northwest China. <i>Plant Disease</i> , 2015, 99, 39-43.	1.4	12
61	Dynamics of the worldwide number of fungi with emphasis on fungal diversity in China. <i>Mycological Progress</i> , 2015, 14, 1.	1.4	47
62	Phylogeny and taxonomy of the genus Abundisporus (Polyporales, Basidiomycota). <i>Mycological Progress</i> , 2015, 14, 1.	1.4	16
63	Two new triterpenoids from fruiting bodies of fungus< i>Ganoderma lucidum</i>. <i>Journal of Asian Natural Products Research</i> , 2015, 17, 750-755.	1.4	17
64	Fungal diversity notes 111–252 taxonomic and phylogenetic contributions to fungal taxa. <i>Fungal Diversity</i> , 2015, 75, 27-274.	12.3	375
65	< i>Fistulina subhepatica</i> sp. nov. from China inferred from morphological and sequence analyses. <i>Mycotaxon</i> , 2015, 130, 47-56.	0.3	6
66	Fragiliporiaceae, a new family of Polyporales (Basidiomycota). <i>Fungal Diversity</i> , 2015, 70, 115-126.	12.3	53
67	Exploring strategies for adsorption of azo dye Congo Red using free and immobilized biomasses of <i>Trametes pubescens</i> . <i>Annals of Microbiology</i> , 2015, 65, 411-421.	2.6	18
68	Morphological characters and molecular data reveal a new species of Fomitopsis (Polyporales) from southern China. <i>Mycoscience</i> , 2015, 56, 168-176.	0.8	21
69	Taxonomy and multi-gene phylogeny of < i>Datronia</i> (< i>Polyporales</i>, < i>Basidiomycota</i>). <i>Persoonia: Molecular Phylogeny and Evolution of Fungi</i> , 2014, 32, 170-182.	4.4	48
70	Three new species of Berkleasmium (Hyphomycetes) from China. <i>Nova Hedwigia</i> , 2014, 98, 151-161.	0.4	6
71	Morphological and molecular evidence for two new species of < i>Laetiporus</i> (Basidiomycota,) Tj ETQq1 1 0.784314 rgBT _{1.9} /Overlock 32	0.3	
72	A preliminary report on decay and canker of < i>Acacia richii</i> caused by < i>Inonotus rickii</i> in China. <i>Forest Pathology</i> , 2014, 44, 82-84.	1.1	6

#	ARTICLE	IF	CITATIONS
73	Decolorization of heterocycle dye Neutral Red by white-rot fungus <i>< i>Perenniporia subacida</i></i> . Desalination and Water Treatment, 2014, 52, 5594-5604.	1.0	14
74	Taxonomy and phylogeny of Ceriporia (Polyporales, Basidiomycota) with an emphasis of Chinese collections. Mycological Progress, 2014, 13, 81-93.	1.4	39
75	Flammeopellis bambusicola gen. et. sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analysis. Mycological Progress, 2014, 13, 771-780.	1.4	17
76	Phlebioporia bubalina gen. et. sp. nov. (Meruliaceae, Polyporales) from southwest China with a preliminary phylogeny based on rDNA sequences. Mycological Progress, 2014, 13, 563-573.	1.4	30
77	Phylogeny and Taxonomy of Climacocystis (Polyporales) in China. Cryptogamie, Mycologie, 2014, 35, 221-231.	1.0	3
78	Morphological and Molecular Evidence for a New Species of Postia (Basidiomycota) from China. Cryptogamie, Mycologie, 2014, 35, 199-207.	1.0	6
79	Morphology and molecular phylogeny for two new species of Fomitopsis (Basidiomycota) from South China. Mycological Progress, 2014, 13, 905-914.	1.4	21
80	Perenniporia cinereofusca sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analysis. Mycoscience, 2014, 55, 417-422.	0.8	7
81	Morphological and Molecular Identification of Two New Species of <i>< i>Hypodontia</i></i> (Schizophoraceae, Hymenochaetales) from Southern China. Cryptogamie, Mycologie, 2014, 35, 87-97.	1.0	24
82	<p align="left">A new species of Postia (Polyporales, Basidiomycota) from China based on morphological and molecular evidence. Phytotaxa, 2014, 162, 147.	0.3	13
83	Helicascus gallicus sp. nov., a new freshwater pleosporalean ascomycete from France. Phytotaxa, 2014, 183, 183.	0.3	12
84	Phylogeny and taxonomy of Ceriporiopsis (Polyporales) with descriptions of two new species from southern China. Phytotaxa, 2014, 164, 17.	0.3	22
85	Studies on Wrightoporia from China 3. Wrightoporia subavellanea sp. nov. based on morphological characters and rDNA sequence data. Phytotaxa, 2014, 175, 225.	0.3	11
86	Biological Pretreatment with White Rot Fungi and Their Co-Culture to Overcome Lignocellulosic Recalcitrance for Improved Enzymatic Digestion. BioResources, 2014, 9, .	1.0	10
87	<i>< i>Dichomitus hubeiensis</i></i> sp. nov. and a new record of <i>< i>Dichomitus</i></i> (Basidiomycota) from China. Nordic Journal of Botany, 2013, 31, 118-121.	0.5	3
88	Decolorization of chemically different dyes by white-rot fungi in submerged cultures. Annals of Microbiology, 2013, 63, 1099-1108.	2.6	36
89	Molecular phylogeny and morphology reveal a new species of Amyloporia (Basidiomycota) from China. Antonie Van Leeuwenhoek, 2013, 104, 817-827.	1.7	24
90	Morphological and molecular identification of four new resupinate species of <i>< i>Perenniporia</i></i> (Polyporales) from southern China. Mycologia, 2013, 105, 945-958.	1.9	32

#	ARTICLE	IF	CITATIONS
91	Two new <i>Fomitopsis</i> species from southern China based on morphological and molecular characters. <i>Mycological Progress</i> , 2013, 12, 709-718.	1.4	33
92	Fungal treatment followed by FeCl ₃ treatment to enhance enzymatic hydrolysis of poplar wood for high sugar yields. <i>Biotechnology Letters</i> , 2013, 35, 2061-2067.	2.2	21
93	< i>Yuchengia</i>, a new polypore genus segregated from < i>Perenniporia</i> (Polyporales) based on morphological and molecular evidence. <i>Nordic Journal of Botany</i> , 2013, 31, 331-338.	0.5	14
94	< i>Antrodia kmetii</i>, a New European Polypore Similar to< i>Antrodia variiformis</i>. <i>Cryptogamie, Mycologie</i> , 2013, 34, 203-209.	1.0	4
95	Purification, biochemical characterization and dye decolorization capacity of an alkali-resistant and metal-tolerant laccase from <i>Trametes pubescens</i> . <i>Bioresource Technology</i> , 2013, 128, 49-57.	9.6	114
96	Dye Congo Red adsorptive decolorization by adsorbents obtained from <i>Trametes pubescens</i> pellets. <i>Desalination and Water Treatment</i> , 2013, 51, 7088-7100.	1.0	5
97	Three new <i>Perenniporia</i> (Polyporales, Basidiomycota) species from China based on morphological and molecular data. <i>Mycoscience</i> , 2013, 54, 231-240.	0.8	10
98	Investigating lignin and hemicellulose in white rot fungus-pretreated wood that affect enzymatic hydrolysis. <i>Bioresource Technology</i> , 2013, 134, 381-385.	9.6	46
99	Genetic diversity of wild <i>Auricularia auricula-judae</i> revealed by ISSR analysis. <i>Biochemical Systematics and Ecology</i> , 2013, 48, 199-205.	1.3	15
100	Two new <i>Daedalea</i> species (Polyporales, Basidiomycota) from South China. <i>Mycoscience</i> , 2013, 54, 62-68.	0.8	21
101	A new fungal peroxidase with alkaline-tolerant, chloride-enhancing activity and dye decolorization capacity. <i>Journal of Molecular Catalysis B: Enzymatic</i> , 2013, 89, 6-14.	1.8	13
102	Taxonomy and phylogeny of the genus < i>Megasporoporia</i> and its related genera. <i>Mycologia</i> , 2013, 105, 368-383.	1.9	58
103	Novel sesquiterpenoids from cultures of the basidiomycete <i>Irpea lacteus</i> . <i>Tetrahedron Letters</i> , 2013, 54, 2651-2654.	1.4	25
104	New species and phylogeny of <i>Perenniporia</i> based on morphological and molecular characters. <i>Fungal Diversity</i> , 2013, 58, 47-60.	12.3	76
105	<i>Antrodia tropica</i> sp. nov. from southern China inferred from morphological characters and molecular data. <i>Mycological Progress</i> , 2013, 12, 223-230.	1.4	30
106	<i>Phellinus castanopsisidis</i> sp. nov. (Hymenochaetaceae) from southern China, with preliminary phylogeny based on rDNA sequences. <i>Mycological Progress</i> , 2013, 12, 341-351.	1.4	27
107	Two new species of < i>Ceriporia</i> (< i>Basidiomycota,</i> < i>Polyporales</i>) with a key to the accepted species in China. <i>Mycotaxon</i> , 2013, 121, 305-312.	0.3	4
108	A new species of < i>Grammothelopsis (Polyporales, Basidiomycota)</i> from southern China. <i>Mycotaxon</i> , 2013, 121, 291-296.	0.3	6

#	ARTICLE	IF	CITATIONS
109	< i>Truncospora macrospora</i> sp. nov. (Polyporales) from Southwest China based on morphological and molecular data. Phytotaxa, 2013, 87, 30.	0.3	10
110	Studies on < i>Wrightoporia</i> from China 2. A new species and three new records from South China. Mycotaxon, 2013, 121, 333-343.	0.3	3
111	Chemical Characterization and Structure of Exopolysaccharides from Submerged Culture of New Medicinal Mushroom from China, <i>Phellinus mori</i> (Higher Basidiomycetes). International Journal of Medicinal Mushrooms, 2013, 15, 57-69.	1.5	12
112	Two new polypores (< i>Ceriporiopsis lavendula</i> and < i>Skeletocutis inflata</i> spp. nov.) from Guangdong Province, China. Nordic Journal of Botany, 2013, 31, 326-330.	0.5	6
113	Study of the Physiological Characteristics of the Medicinal Mushroom <i>Trametes pubescens</i> (Higher) Tj ETQq1 1 0.784314 rgBT /Overlooked Mushrooms, 2013, 15, 199-210.	1.5	8
114	A new species of < i>Postia</i> (< i>Basidiomycota</i>) from Northeast China. Mycotaxon, 2012, 120, 231-237.	0.3	13
115	Morphological and molecular evidence for a new species of <i>Perenniporia</i> (Basidiomycota) from Tibet, southwestern China. Mycoscience, 2012, 53, 365-372.	0.8	24
116	Pretreatment of <i>Populus tomentosa</i> with <i>Trametes velutina</i> supplemented with inorganic salts enhances enzymatic hydrolysis for ethanol production. Biotechnology Letters, 2012, 34, 2241-2246.	2.2	14
117	Statistical optimization of cellulase production by the brown rot fungi, <i>Fomitopsis palustris</i> , and its application in the enzymatic hydrolysis of LHW-pretreated woody biomass. Process Biochemistry, 2012, 47, 2552-2556.	3.7	15
118	A new species of <i>Perenniporia</i> (Polyporales, Basidiomycota) described from southern China based on morphological and molecular characters. Mycological Progress, 2012, 11, 555-560.	1.4	35
119	Combination of biological pretreatment with liquid hot water pretreatment to enhance enzymatic hydrolysis of <i>Populus tomentosa</i> . Bioresource Technology, 2012, 107, 282-286.	9.6	96
120	An Antitumor Component from <i>Fomitiporia ellipsoidea</i> . Journal of Microbiology and Biotechnology, 2012, 22, 1482-1485.	2.1	4
121	Wood-Inhabiting Fungi in Southern China. 4. Polypores from Hainan Province. Annales Botanici Fennici, 2011, 48, 219-231.	0.1	56
122	Wood-Rotting Fungi in Eastern China. 5. Polypore Diversity in Jiangxi Province. Annales Botanici Fennici, 2011, 48, 237-246.	0.1	22
123	<i>Fomitiporia ellipsoidea</i> has the largest fruiting body among the fungi. Fungal Biology, 2011, 115, 813-814.	2.5	42
124	Notes on < i>Ceriporia</i> (< i>Basidiomycota</i>, < i>Polyporales</i>) in China. Mycotaxon, 2011, 116, 457-468.	0.3	16
125	A new species of Pyrofomes (Basidiomycota, Polyporaceae) from China. Nova Hedwigia, 2011, 93, 437-441.	0.4	6
126	Wood-rotting fungi in eastern China 6. Two new species of < i>Antrodia</i> (< i>Basidiomycota</i>) from Mt. Huangshan, Anhui Province. Mycotaxon, 2011, 116, 13-20.	0.3	17

#	ARTICLE	IF	CITATIONS
127	Three new species of <i>Inonotus</i> (Basidiomycota, Hymenochaetaceae) from China. <i>Mycological Progress</i> , 2011, 10, 107-114.	1.4	38
128	Morphological and molecular evidences for a new species of <i>Lignosus</i> (Polyporales, Basidiomycota) from tropical China. <i>Mycological Progress</i> , 2011, 10, 267-271.	1.4	29
129	< i> <i>Melanoderma microcarpum</i> </i> gen. et sp. nov. (< i>Basidiomycota</i>) from China. <i>Mycotaxon</i> , 2011, 116, 295-302.	0.3	25
130	High Genetic Diversity in Wild Culinary-Medicinal Wood Ear Mushroom, <i>Auricularia polytricha</i> (Mont.) Sacc., in Tropical China Revealed by ISSR Analysis. <i>International Journal of Medicinal Mushrooms</i> , 2011, 13, 289-298.	1.5	15
131	Two new species of < i>Phylloporia </i> (< i>Basidiomycota</i>,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 0.3 37 10 50 58 2 1		
132	A new < i>Trametes </i> species from Southwest China. <i>Mycotaxon</i> , 2010, 113, 263-267.	0.3	23
133	Current advances in <i>Phellinus sensu lato</i> : medicinal species, functions, metabolites and mechanisms. <i>Applied Microbiology and Biotechnology</i> , 2010, 87, 1587-1593.	3.6	86
134	< i> <i>Bondarzewia podocarpi</i> </i>, a new and remarkable polypore from tropical China. <i>Mycologia</i> , 2010, 102, 881-886.	1.9	37
135	Notes on the genus <i>Rigidoporus</i> (Basidiomycota, Polyporaceae) in China. <i>Nova Hedwigia</i> , 2009, 88, 189-197.	0.4	10
136	Trichaptum (Basidiomycota, Hymenochaetales) from China with a description of three new species. <i>Mycological Progress</i> , 2009, 8, 281-287.	1.4	36
137	< i> <i>Oxyporus piceicola</i> </i> sp. nov. with a key to species of the genus in China. <i>Mycotaxon</i> , 2009, 109, 307-313.	0.3	4
138	Wood-inhabiting fungi in southern China 3. A new species of < i>Phellinus </i> (< i>Hymenochaetales</i>) from tropical China. <i>Mycotaxon</i> , 2009, 110, 125-130.	0.3	30
139	Two new species of < i>Megasporoporia </i> (< i>Polyporales, Basidiomycota</i>) from tropical China. <i>Mycotaxon</i> , 2009, 110, 131-138.	0.3	24
140	Species Diversity and Utilization of Medicinal Mushrooms and Fungi in China (Review). <i>International Journal of Medicinal Mushrooms</i> , 2009, 11, 287-302.	1.5	221
141	A new species of <i>Fomitiporia</i> (Hymenochaetaceae, Basidiomycota) from China based on morphological and molecular characters. <i>Mycological Research</i> , 2008, 112, 375-380.	2.5	29
142	Polypores from eastern Inner Mongolia, northeastern China. <i>Nova Hedwigia</i> , 2007, 84, 513-520.	0.4	3
143	Pathogenic wood-decaying fungi in China. <i>Forest Pathology</i> , 2007, 37, 105-120.	1.1	108
144	<i>Wrightoporia</i> (Basidiomycota, Aphyllophorales) in China. <i>Nova Hedwigia</i> , 2006, 83, 159-166.	0.4	12

ARTICLE

IF

CITATIONS

145	Species diversity, molecular phylogeny and ecological habits of <i>Cyanosporus</i> (Polyporales.) Tj ETQql 1 0.784314 rgBT /Overlock 10	1.9	1
146	Taxonomy and molecular phylogeny of <i>Trametopsis</i> (Polyporales, Basidiomycota) with descriptions of two new species. MycoKeys, 0, 90, 31-51.	1.9	3