

Lambros Kordelas

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8672942/publications.pdf>

Version: 2024-02-01

34
papers

1,894
citations

567281

15
h-index

501196

28
g-index

35
all docs

35
docs citations

35
times ranked

4027
citing authors

#	ARTICLE	IF	CITATIONS
1	Pre-Transplant Serum Leptin Levels and Relapse of Acute Myeloid Leukemia after Allogeneic Transplantation. <i>International Journal of Molecular Sciences</i> , 2022, 23, 2337.	4.1	1
2	Treatment of myeloid malignancies relapsing after allogeneic hematopoietic stem cell transplantation with venetoclax and hypomethylating agents—a retrospective multicenter analysis on behalf of the German Cooperative Transplant Study Group. <i>Annals of Hematology</i> , 2021, 100, 959-968.	1.8	34
3	EASIX and mortality after allogeneic stem cell transplantation. <i>Bone Marrow Transplantation</i> , 2020, 55, 553-561.	2.4	70
4	Allogeneic transplantation of peripheral blood stem cell grafts results in a massive decrease of primitive hematopoietic progenitor frequencies in reconstituted bone marrows. <i>Bone Marrow Transplantation</i> , 2020, 55, 100-109.	2.4	1
5	Optimizing anti-T-lymphocyte globulin dosing to improve long-term outcome after unrelated hematopoietic cell transplantation for hematologic malignancies. <i>American Journal of Transplantation</i> , 2020, 20, 677-688.	4.7	9
6	Interleukin-18 and Hematopoietic Recovery after Allogeneic Stem Cell Transplantation. <i>Cancers</i> , 2020, 12, 2789.	3.7	7
7	Pre-transplant testosterone and outcome of men after allogeneic stem cell transplantation. <i>Haematologica</i> , 2020, 105, 1454-1464.	3.5	2
8	Rapid Rebound of a Preexisting CXCR4-tropic Human Immunodeficiency Virus Variant After Allogeneic Transplantation With CCR5 Δ32 Homozygous Stem Cells. <i>Clinical Infectious Diseases</i> , 2019, 68, 684-687.	5.8	42
9	Human multipotent hematopoietic progenitor cell expansion is neither supported in endothelial and endothelial/mesenchymal co-cultures nor in NSG mice. <i>Scientific Reports</i> , 2019, 9, 12914.	3.3	4
10	Individual Immune-Modulatory Capabilities of MSC-Derived Extracellular Vesicle (EV) Preparations and Recipient-Dependent Responsiveness. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1642.	4.1	36
11	Interleukin-18 and outcome after allogeneic stem cell transplantation: A retrospective cohort study. <i>EBioMedicine</i> , 2019, 49, 202-212.	6.1	11
12	Asymmetric dimethylarginine serum levels are associated with early mortality after allogeneic stem cell transplantation. <i>Haematologica</i> , 2019, 104, 827-834.	3.5	4
13	Decreased Soluble Human Leukocyte Antigen E Levels in Patients After Allogeneic Hematopoietic Stem Cell Transplantation Are Associated With Severe Acute and Extended Chronic Graft-versus-Host Disease and Inferior Overall Survival. <i>Frontiers in Immunology</i> , 2019, 10, 3027.	4.8	7
14	Treatment of Relapse after Allogeneic Hematopoietic Stem Cell Transplantation with Venetoclax, Hypomethylating Agents and DLI - a Retrospective Multi Center Study. <i>Blood</i> , 2019, 134, 4563-4563.	1.4	3
15	High Pre-Transplant Free Interleukin-18 Is Associated with Poor Hematopoietic Recovery after Allogeneic Stem Cell Transplantation. <i>Blood</i> , 2019, 134, 4507-4507.	1.4	0
16	High-dose melphalan-based sequential conditioning chemotherapy followed by allogeneic haematopoietic stem cell transplantation in adult patients with relapsed or refractory acute myeloid leukaemia. <i>British Journal of Haematology</i> , 2018, 180, 840-853.	2.5	15
17	Elevated soluble human leukocyte antigen G levels in patients after allogeneic stem cell transplantation are associated with less severe acute and chronic graft-versus-host disease. <i>Bone Marrow Transplantation</i> , 2018, 53, 1149-1156.	2.4	9
18	Pre-Conditioning Serum Leptin Levels Predict Early Relapse Rates of High Risk AML after Allogeneic Stem Cell Transplantation. <i>Blood</i> , 2018, 132, 3419-3419.	1.4	0

#	ARTICLE	IF	CITATIONS
19	Intestinal T lymphocyte homing is associated with gastric emptying and epithelial barrier function in critically ill: a prospective observational study. <i>Critical Care</i> , 2017, 21, 70.	5.8	20
20	Pretransplant Vitamin D Deficiency Is Associated With Higher Relapse Rates in Patients Allografted for Myeloid Malignancies. <i>Journal of Clinical Oncology</i> , 2017, 35, 3143-3152.	1.6	27
21	Clinical potential of mesenchymal stem/stromal cell-derived extracellular vesicles. <i>Stem Cell Investigation</i> , 2017, 4, 84-84.	3.0	131
22	The Activating NKG2C Receptor Is Significantly Reduced in NK Cells after Allogeneic Stem Cell Transplantation in Patients with Severe Graft-versus-Host Disease. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1797.	4.1	30
23	CD34+ highly enriched allogeneic stem cell transplantation in a patient with mixed phenotype acute leukemia and <i>Fusarium solani</i> sepsis. <i>Annals of Hematology</i> , 2016, 95, 155-156.	1.8	3
24	Adoptive Transfer of Purified Donor-B-Lymphocytes after Allogeneic Stem Cell Transplantation: Results from a Phase I/IIa Clinical Trial. <i>Blood</i> , 2016, 128, 502-502.	1.4	11
25	Applying extracellular vesicles based therapeutics in clinical trials – an ISEV position paper. <i>Journal of Extracellular Vesicles</i> , 2015, 4, 30087.	12.2	1,020
26	<sc>CD</sc>133 allows elaborated discrimination and quantification of haematopoietic progenitor subsets in human haematopoietic stem cell transplants. <i>British Journal of Haematology</i> , 2015, 169, 868-878.	2.5	31
27	More on Shift of HIV Tropism in Stem-Cell Transplantation with <i>CCR5</i> Delta32/Delta32 Mutation. <i>New England Journal of Medicine</i> , 2014, 371, 2437-2438.	27.0	69
28	Shift of HIV Tropism in Stem-Cell Transplantation with <i>CCR5</i> Delta32 Mutation. <i>New England Journal of Medicine</i> , 2014, 371, 880-882.	27.0	144
29	Haploidentical allogeneic hematopoietic cell transplantation in adults using CD3/CD19 depletion and reduced intensity conditioning: a phase II study. <i>Haematologica</i> , 2012, 97, 1523-1531.	3.5	116
30	Small interfering RNA against BCR-ABL transcripts sensitize mutated T315I cells to nilotinib. <i>Haematologica</i> , 2010, 95, 388-397.	3.5	20
31	Results of a Phase II Study of Haploidentical Hematopoietic Cell Transplantation (HHCT) in Adults Using Reduced Intensity Conditioning and CD3/CD19-Depleted Grafts: Clinical Outcome and Immune Reconstitution. <i>Blood</i> , 2009, 114, 1203-1203.	1.4	0
32	Successful Treatment of EBV PTLD with CNS Lymphomas with the Monoclonal Anti-CD20 Antibody Rituximab. <i>Oncology Research and Treatment</i> , 2008, 31, 691-693.	1.2	17
33	Improved Overall Survival in Patients Recovering with High Gamma/Delta T Cells after Allogeneic Hematopoietic Stem Cell Transplantation. <i>Blood</i> , 2008, 112, 2223-2223.	1.4	0
34	Haploidentical Allogeneic Hematopoietic Cell Transplantation in Adults Using Reduced Intensity Conditioning and CD3/CD19-Depleted Grafts: Interim Analysis of a Phase I/II Study. <i>Blood</i> , 2008, 112, 1962-1962.	1.4	0