

Olivier Outteryck

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8711455/publications.pdf>

Version: 2024-02-01

31
papers

1,599
citations

394421

19
h-index

454955

30
g-index

31
all docs

31
docs citations

31
times ranked

2302
citing authors

#	ARTICLE	IF	CITATIONS
1	Optic Nerve Lesion Length at the Acute Phase of Optic Neuritis Is Predictive of Retinal Neuronal Loss. <i>Neurology: Neuroimmunology and NeuroInflammation</i> , 2022, 9, .	6.0	16
2	Long-term effect of natalizumab in patients with RRMS: TYSTEN cohort. <i>Multiple Sclerosis Journal</i> , 2021, 27, 729-741.	3.0	9
3	Artificial intelligence extension of the OSCAR criteria. <i>Annals of Clinical and Translational Neurology</i> , 2021, 8, 1528-1542.	3.7	33
4	Optical coherence tomography for detection of asymptomatic optic nerve lesions in clinically isolated syndrome. <i>Neurology</i> , 2020, 95, e733-e744.	1.1	29
5	COVID-19 Neurologic Complication with CNS Vasculitis-Like Pattern. <i>American Journal of Neuroradiology</i> , 2020, 41, 1384-1387.	2.4	128
6	Follow-Up MRI for Small Brain AVMs Treated by Radiosurgery: Is Gadolinium Really Necessary?. <i>American Journal of Neuroradiology</i> , 2020, 41, 437-445.	2.4	9
7	Asymptomatic optic nerve lesions. <i>Neurology</i> , 2020, 94, e2468-e2478.	1.1	37
8	Optical coherence tomography in myelin-oligodendrocyte-glycoprotein antibody-seropositive patients: a longitudinal study. <i>Journal of Neuroinflammation</i> , 2019, 16, 154.	7.2	61
9	Optical coherence tomography: a window to the optic nerve in clinically isolated syndrome. <i>Brain</i> , 2019, 142, 903-915.	7.6	33
10	Optic nerve double inversion recovery hypersignal in patients with clinically isolated syndrome is associated with asymptomatic gadolinium-enhanced lesion. <i>Multiple Sclerosis Journal</i> , 2019, 25, 1888-1895.	3.0	12
11	Magnetic resonance imaging changes following natalizumab discontinuation in multiple sclerosis patients with progressive multifocal leukoencephalopathy. <i>Multiple Sclerosis Journal</i> , 2018, 24, 1902-1908.	3.0	3
12	MD1003 (High-Dose Pharmaceutical-Grade Biotin) for the Treatment of Chronic Visual Loss Related to Optic Neuritis in Multiple Sclerosis: A Randomized, Double-Blind, Placebo-Controlled Study. <i>CNS Drugs</i> , 2018, 32, 661-672.	5.9	26
13	Continuous hemifacial myokymia as the revealing symptom of demyelinating disease of the CNS. <i>Multiple Sclerosis and Related Disorders</i> , 2017, 11, 10-11.	2.0	4
14	Neuralgic amyotrophy triggered by hepatitis E virus: a particular phenotype. <i>Journal of Neurology</i> , 2017, 264, 770-780.	3.6	25
15	Autoimmune cerebellar ataxia with glutamic acid decarboxylase 65 antibodies associated with central vestibular symptoms. <i>Acta Neurologica Belgica</i> , 2017, 117, 775-776.	1.1	1
16	Retinal layer segmentation in multiple sclerosis: a systematic review and meta-analysis. <i>Lancet Neurology</i> , The, 2017, 16, 797-812.	10.2	397
17	Late-onset of Alpers-Huttenlocher syndrome: an unusual cause of refractory epilepsy and liver failure. <i>Acta Neurologica Belgica</i> , 2017, 117, 399-401.	1.1	3
18	Retinal thickness measured with optical coherence tomography and risk of disability worsening in multiple sclerosis: a cohort study. <i>Lancet Neurology</i> , The, 2016, 15, 574-584.	10.2	266

#	ARTICLE	IF	CITATIONS
19	Punctate pattern. <i>Neurology</i> , 2016, 86, 1516-1523.	1.1	65
20	Natalizumab in relapsing-remitting multiple sclerosis. <i>Expert Review of Neurotherapeutics</i> , 2016, 16, 471-481.	2.8	8
21	OCT Findings in Neuromyelitis Optica Spectrum Disorders. , 2016, , 85-96.		2
22	Asymptomatic Progressive Multifocal Leukoencephalopathy Associated with Natalizumab: Diagnostic Precision with MR Imaging. <i>Radiology</i> , 2016, 278, 863-872.	7.3	38
23	Rituximab as first-line therapy in neuromyelitis optica: efficiency and tolerability. <i>Journal of Neurology</i> , 2015, 262, 2329-2335.	3.6	86
24	A comparative optical coherence tomography study in neuromyelitis optica spectrum disorder and multiple sclerosis. <i>Multiple Sclerosis Journal</i> , 2015, 21, 1781-1793.	3.0	64
25	JC-virus seroconversion in multiple sclerosis patients receiving natalizumab. <i>Multiple Sclerosis Journal</i> , 2014, 20, 822-829.	3.0	62
26	Comparison of 3D double inversion recovery and 2D STIR FLAIR MR sequences for the imaging of optic neuritis: pilot study. <i>European Radiology</i> , 2014, 24, 3069-3075.	4.5	36
27	Extensive myelitis associated with anti-NMDA receptor antibodies. <i>BMC Neurology</i> , 2013, 13, 211.	1.8	21
28	Methionine synthase deficiency: A rare cause of adult-onset leukoencephalopathy. <i>Neurology</i> , 2012, 79, 386-388.	1.1	13
29	Anti-JCV antibody prevalence in a French cohort of MS patients under natalizumab therapy. <i>Journal of Neurology</i> , 2012, 259, 2293-2298.	3.6	34
30	Demographic and clinic characteristics of French patients treated with natalizumab in clinical practice. <i>Journal of Neurology</i> , 2010, 257, 207-211.	3.6	41
31	Optical Coherence Tomography in Clinically Isolated Syndrome. <i>Archives of Neurology</i> , 2009, 66, 1373-7.	4.5	37