

Dawn M Guthrie

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8732536/publications.pdf>

Version: 2024-02-01

47
papers

764
citations

687363

13
h-index

580821

25
g-index

47
all docs

47
docs citations

47
times ranked

960
citing authors

#	ARTICLE	IF	CITATIONS
1	Combined impairments in vision, hearing and cognition are associated with greater levels of functional and communication difficulties than cognitive impairment alone: Analysis of interRAI data for home care and long-term care recipients in Ontario. PLoS ONE, 2018, 13, e0192971.	2.5	102
2	The Health and Well-Being of Older Adults with Dual Sensory Impairment (DSI) in Four Countries. PLoS ONE, 2016, 11, e0155073.	2.5	68
3	Risk Factors for Restriction in Activity Associated With Fear of Falling Among Seniors Within the Community. Journal of Patient Safety, 2010, 6, 187-191.	1.7	63
4	Older Adults With a Combination of Vision and Hearing Impairment Experience Higher Rates of Cognitive Impairment, Functional Dependence, and Worse Outcomes Across a Set of Quality Indicators. Journal of Aging and Health, 2019, 31, 85-108.	1.7	55
5	Associations between sensory loss and social networks, participation, support, and loneliness: Analysis of the Canadian Longitudinal Study on Aging. Canadian Family Physician, 2018, 64, e33-e41.	0.4	51
6	The Prevalence of Hearing, Vision, and Dual Sensory Loss in Older Canadians: An Analysis of Data from the Canadian Longitudinal Study on Aging. Canadian Journal on Aging, 2021, 40, 1-22.	1.1	35
7	Effect of Early Palliative Care on End-of-Life Health Care Costs: A Population-Based, Propensity Score-Matched Cohort Study. JCO Oncology Practice, 2022, 18, e183-e192.	2.9	25
8	Patient Characteristics Associated With Prognostic Awareness: A Study of a Canadian Palliative Care Population Using the InterRAI Palliative Care Instrument. Journal of Pain and Symptom Management, 2015, 49, 716-725.	1.2	24
9	Hearing and vision screening tools for long-term care residents with dementia: protocol for a scoping review. BMJ Open, 2016, 6, e011945.	1.9	24
10	Prevalence and risk factors of depressive symptoms in a Canadian palliative home care population: a cross-sectional study. BMC Palliative Care, 2014, 13, 10.	1.8	21
11	A comparison of home care quality indicator rates in two Canadian provinces. BMC Health Services Research, 2014, 14, 37.	2.2	20
12	Reliability of standardized assessment for adults who are deafblind. Journal of Rehabilitation Research and Development, 2011, 48, 545.	1.6	19
13	Development and Validation of a Prognostic Survival Model With Patient-Reported Outcomes for Patients With Cancer. JAMA Network Open, 2020, 3, e201768.	5.9	16
14	End-of-life outcomes with or without early palliative care: a propensity score matched, population-based cancer cohort study. BMJ Open, 2021, 11, e041432.	1.9	15
15	The CaregiverVoice Survey: A Pilot Study Surveying Bereaved Caregivers To Measure the Caregiver and Patient Experience at End of Life. Journal of Palliative Medicine, 2016, 19, 712-719.	1.1	14
16	Trajectory of psychosocial symptoms among home care patients with cancer at end-of-life. Psycho-Oncology, 2021, 30, 103-110.	2.3	14
17	Detection of vision and /or hearing loss using the interRAI Community Health Assessment aligns well with common behavioral vision/hearing measurements. PLoS ONE, 2019, 14, e0223123.	2.5	13
18	A Comparison Between End-of-Life Home Care Clients With Cancer and Heart Failure in Ontario. Home Health Care Services Quarterly, 2015, 34, 14-29.	0.7	12

#	ARTICLE	IF	CITATIONS
19	Quality Indicator Rates for Seriously Ill Home Care Clients: Analysis of Resident Assessment Instrument for Home Care Data in Six Canadian Provinces. <i>Journal of Palliative Medicine</i> , 2019, 22, 1346-1356.	1.1	12
20	Self-Rated Health, Cognition, and Dual Sensory Impairment Are Important Predictors of Depression Among Home Care Clients in Ontario. <i>Home Health Care Management and Practice</i> , 2016, 28, 35-43.	1.0	11
21	Changes in social engagement and depression predict incident loneliness among seriously ill home care clients. <i>Palliative and Supportive Care</i> , 2018, 16, 170-179.	1.0	11
22	Trajectory of End-of-Life Pain and Other Physical Symptoms among Cancer Patients Receiving Home Care. <i>Current Oncology</i> , 2021, 28, 1641-1651.	2.2	11
23	The relationship between caregiver burden and depressive symptoms in Ontario home care clients. <i>Home Health Care Services Quarterly</i> , 2018, 37, 60-76.	0.7	10
24	Development and Validation of Caregiver Risk Evaluation (CaRE): A New Algorithm to Screen for Caregiver Burden. <i>Journal of Applied Gerontology</i> , 2021, 40, 731-741.	2.0	10
25	The Role of Medications in Predicting Activity Restriction Due to a Fear of Falling. <i>Journal of Aging and Health</i> , 2012, 24, 269-286.	1.7	9
26	Potential quality indicators for seriously ill home care clients: a cross-sectional analysis using Resident Assessment Instrument for Home Care (RAI-HC) data for Ontario. <i>BMC Palliative Care</i> , 2019, 18, 3.	1.8	9
27	A Deterioration in Hearing Is Associated With Functional and Cognitive Impairments, Difficulty With Communication, and Greater Health Instability. <i>Journal of Applied Gerontology</i> , 2020, 39, 159-171.	2.0	9
28	Data sharing between home care professionals: a feasibility study using the RAI Home Care instrument. <i>BMC Geriatrics</i> , 2014, 14, 81.	2.7	8
29	Dimensionality, Reliability and Validity of the InterRAI Depression Rating Scale in a Canadian Palliative Care Population. <i>Social Indicators Research</i> , 2015, 122, 297-314.	2.7	7
30	Hearing and Cognitive Impairments Increase the Risk of Long-term Care Admissions. <i>Innovation in Aging</i> , 2020, 4, igz053.	0.1	7
31	Development and validation of a prediction model of poor performance status and severe symptoms over time in cancer patients (PROVIEW+). <i>Palliative Medicine</i> , 2021, 35, 1713-1723.	3.1	7
32	Older adults with dual sensory loss in rehabilitation show high functioning and may fare better than those with single sensory loss. <i>PLoS ONE</i> , 2020, 15, e0237152.	2.5	6
33	Development of a case-mix funding system for adults with combined vision and hearing loss. <i>BMC Health Services Research</i> , 2013, 13, 137.	2.2	5
34	Factors Associated With Terminally Ill People Who Want to Die. <i>Journal of Pain and Symptom Management</i> , 2020, 60, 539-548.e1.	1.2	5
35	What should be measured to assess the quality of community-based palliative care? Results from a collaborative expert workshop. <i>Palliative and Supportive Care</i> , 2022, 20, 226-232.	1.0	5
36	Prevalence and Risk Factors of Breathlessness Across Canada: A National Retrospective Cohort Study in Home Care and Nursing Home Populations. <i>Journal of Pain and Symptom Management</i> , 2021, 62, 346-354.e1.	1.2	5

#	ARTICLE	IF	CITATIONS
37	Cross-sectional Nutrition Profile of Palliative Home Care Clients in Ontario and Performance of the interRAI Palliative Care Nutrition Clinical Assessment Protocol. <i>Journal of Parenteral and Enteral Nutrition</i> , 2021, 45, 183-192.	2.6	4
38	“The system is well intentioned, but complicated and fallible”-interviews with caregivers and decision makers about palliative care in Canada. <i>BMC Palliative Care</i> , 2021, 20, 149.	1.8	4
39	A Newly Identified Impairment in Both Vision and Hearing Increases the Risk of Deterioration in Both Communication and Cognitive Performance. <i>Canadian Journal on Aging</i> , 0, , 1-14.	1.1	4
40	Prognosis does not change the landscape: palliative home care clients experience high rates of pain and nausea, regardless of prognosis. <i>BMC Palliative Care</i> , 2021, 20, 165.	1.8	4
41	A multi-stage process to develop quality indicators for community-based palliative care using interRAI data. <i>PLoS ONE</i> , 2022, 17, e0266569.	2.5	4
42	Symptom trajectories of non-cancer patients in the last six months of life: Identifying needs in a population-based home care cohort. <i>PLoS ONE</i> , 2021, 16, e0252814.	2.5	3
43	The Influence of Physical and Psychosocial Factors on Disruptive Pain Among Seriously Ill Home Care Patients. <i>Journal of Palliative Care</i> , 2017, 32, 61-68.	1.0	1
44	Pan-Canadian Quality Indicators for Patients at End of Life Derived from interRAI Data. <i>Journal of Pain and Symptom Management</i> , 2018, 56, e59-e60.	1.2	1
45	A Profile of Older Adults With a History of Stroke Receiving Home Care in Ontario, Canada. <i>Home Health Care Management and Practice</i> , 2019, 31, 162-171.	1.0	1
46	Benefits of Early Palliative Care in the Community: A Propensity Score Matched Cancer Cohort. <i>Journal of Pain and Symptom Management</i> , 2018, 56, e42.	1.2	0
47	Optimizing Evaluation of Older Adults With Vision and/or Hearing Loss Using the interRAI Community Health Assessment and Deafblind Supplement. <i>Frontiers in Rehabilitation Sciences</i> , 2021, 2, .	1.2	0