

Dawn M Guthrie

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8732536/publications.pdf>

Version: 2024-02-01

47
papers

764
citations

687363

13
h-index

580821

25
g-index

47
all docs

47
docs citations

47
times ranked

960
citing authors

#	ARTICLE	IF	CITATIONS
1	What should be measured to assess the quality of community-based palliative care? Results from a collaborative expert workshop. <i>Palliative and Supportive Care</i> , 2022, 20, 226-232.	1.0	5
2	Effect of Early Palliative Care on End-of-Life Health Care Costs: A Population-Based, Propensity Score-Matched Cohort Study. <i>JCO Oncology Practice</i> , 2022, 18, e183-e192.	2.9	25
3	A multi-stage process to develop quality indicators for community-based palliative care using interRAI data. <i>PLoS ONE</i> , 2022, 17, e0266569.	2.5	4
4	Cross-sectional Nutrition Profile of Palliative Home Care Clients in Ontario and Performance of the interRAI Palliative Care Nutrition Clinical Assessment Protocol. <i>Journal of Parenteral and Enteral Nutrition</i> , 2021, 45, 183-192.	2.6	4
5	Development and Validation of Caregiver Risk Evaluation (CaRE): A New Algorithm to Screen for Caregiver Burden. <i>Journal of Applied Gerontology</i> , 2021, 40, 731-741.	2.0	10
6	The Prevalence of Hearing, Vision, and Dual Sensory Loss in Older Canadians: An Analysis of Data from the Canadian Longitudinal Study on Aging. <i>Canadian Journal on Aging</i> , 2021, 40, 1-22.	1.1	35
7	Trajectory of psychosocial symptoms among home care patients with cancer at end-of-life. <i>Psycho-Oncology</i> , 2021, 30, 103-110.	2.3	14
8	End-of-life outcomes with or without early palliative care: a propensity score matched, population-based cancer cohort study. <i>BMJ Open</i> , 2021, 11, e041432.	1.9	15
9	Trajectory of End-of-Life Pain and Other Physical Symptoms among Cancer Patients Receiving Home Care. <i>Current Oncology</i> , 2021, 28, 1641-1651.	2.2	11
10	Symptom trajectories of non-cancer patients in the last six months of life: Identifying needs in a population-based home care cohort. <i>PLoS ONE</i> , 2021, 16, e0252814.	2.5	3
11	Development and validation of a prediction model of poor performance status and severe symptoms over time in cancer patients (PROVIEW+). <i>Palliative Medicine</i> , 2021, 35, 1713-1723.	3.1	7
12	Prevalence and Risk Factors of Breathlessness Across Canada: A National Retrospective Cohort Study in Home Care and Nursing Home Populations. <i>Journal of Pain and Symptom Management</i> , 2021, 62, 346-354.e1.	1.2	5
13	"The system is well intentioned, but complicated and fallible" - interviews with caregivers and decision makers about palliative care in Canada. <i>BMC Palliative Care</i> , 2021, 20, 149.	1.8	4
14	Prognosis does not change the landscape: palliative home care clients experience high rates of pain and nausea, regardless of prognosis. <i>BMC Palliative Care</i> , 2021, 20, 165.	1.8	4
15	Optimizing Evaluation of Older Adults With Vision and/or Hearing Loss Using the interRAI Community Health Assessment and Deafblind Supplement. <i>Frontiers in Rehabilitation Sciences</i> , 2021, 2, .	1.2	0
16	A Deterioration in Hearing Is Associated With Functional and Cognitive Impairments, Difficulty With Communication, and Greater Health Instability. <i>Journal of Applied Gerontology</i> , 2020, 39, 159-171.	2.0	9
17	Older adults with dual sensory loss in rehabilitation show high functioning and may fare better than those with single sensory loss. <i>PLoS ONE</i> , 2020, 15, e0237152.	2.5	6
18	Hearing and Cognitive Impairments Increase the Risk of Long-term Care Admissions. <i>Innovation in Aging</i> , 2020, 4, igz053.	0.1	7

#	ARTICLE	IF	CITATIONS
19	Factors Associated With Terminally Ill People Who Want to Die. <i>Journal of Pain and Symptom Management</i> , 2020, 60, 539-548.e1.	1.2	5
20	Development and Validation of a Prognostic Survival Model With Patient-Reported Outcomes for Patients With Cancer. <i>JAMA Network Open</i> , 2020, 3, e201768.	5.9	16
21	Detection of vision and /or hearing loss using the interRAI Community Health Assessment aligns well with common behavioral vision/hearing measurements. <i>PLoS ONE</i> , 2019, 14, e0223123.	2.5	13
22	Quality Indicator Rates for Seriously Ill Home Care Clients: Analysis of Resident Assessment Instrument for Home Care Data in Six Canadian Provinces. <i>Journal of Palliative Medicine</i> , 2019, 22, 1346-1356.	1.1	12
23	A Profile of Older Adults With a History of Stroke Receiving Home Care in Ontario, Canada. <i>Home Health Care Management and Practice</i> , 2019, 31, 162-171.	1.0	1
24	Potential quality indicators for seriously ill home care clients: a cross-sectional analysis using Resident Assessment Instrument for Home Care (RAI-HC) data for Ontario. <i>BMC Palliative Care</i> , 2019, 18, 3.	1.8	9
25	Older Adults With a Combination of Vision and Hearing Impairment Experience Higher Rates of Cognitive Impairment, Functional Dependence, and Worse Outcomes Across a Set of Quality Indicators. <i>Journal of Aging and Health</i> , 2019, 31, 85-108.	1.7	55
26	The relationship between caregiver burden and depressive symptoms in Ontario home care clients. <i>Home Health Care Services Quarterly</i> , 2018, 37, 60-76.	0.7	10
27	Changes in social engagement and depression predict incident loneliness among seriously ill home care clients. <i>Palliative and Supportive Care</i> , 2018, 16, 170-179.	1.0	11
28	Benefits of Early Palliative Care in the Community: A Propensity Score Matched Cancer Cohort. <i>Journal of Pain and Symptom Management</i> , 2018, 56, e42.	1.2	0
29	Pan-Canadian Quality Indicators for Patients at End of Life Derived from interRAI Data. <i>Journal of Pain and Symptom Management</i> , 2018, 56, e59-e60.	1.2	1
30	Combined impairments in vision, hearing and cognition are associated with greater levels of functional and communication difficulties than cognitive impairment alone: Analysis of interRAI data for home care and long-term care recipients in Ontario. <i>PLoS ONE</i> , 2018, 13, e0192971.	2.5	102
31	Associations between sensory loss and social networks, participation, support, and loneliness: Analysis of the Canadian Longitudinal Study on Aging. <i>Canadian Family Physician</i> , 2018, 64, e33-e41.	0.4	51
32	The Influence of Physical and Psychosocial Factors on Disruptive Pain Among Seriously Ill Home Care Patients. <i>Journal of Palliative Care</i> , 2017, 32, 61-68.	1.0	1
33	Hearing and vision screening tools for long-term care residents with dementia: protocol for a scoping review. <i>BMJ Open</i> , 2016, 6, e011945.	1.9	24
34	The CaregiverVoice Survey: A Pilot Study Surveying Bereaved Caregivers To Measure the Caregiver and Patient Experience at End of Life. <i>Journal of Palliative Medicine</i> , 2016, 19, 712-719.	1.1	14
35	Self-Rated Health, Cognition, and Dual Sensory Impairment Are Important Predictors of Depression Among Home Care Clients in Ontario. <i>Home Health Care Management and Practice</i> , 2016, 28, 35-43.	1.0	11
36	The Health and Well-Being of Older Adults with Dual Sensory Impairment (DSI) in Four Countries. <i>PLoS ONE</i> , 2016, 11, e0155073.	2.5	68

#	ARTICLE	IF	CITATIONS
37	Dimensionality, Reliability and Validity of the InterRAI Depression Rating Scale in a Canadian Palliative Care Population. <i>Social Indicators Research</i> , 2015, 122, 297-314.	2.7	7
38	A Comparison Between End-of-Life Home Care Clients With Cancer and Heart Failure in Ontario. <i>Home Health Care Services Quarterly</i> , 2015, 34, 14-29.	0.7	12
39	Patient Characteristics Associated With Prognostic Awareness: A Study of a Canadian Palliative Care Population Using the InterRAI Palliative Care Instrument. <i>Journal of Pain and Symptom Management</i> , 2015, 49, 716-725.	1.2	24
40	Prevalence and risk factors of depressive symptoms in a Canadian palliative home care population: a cross-sectional study. <i>BMC Palliative Care</i> , 2014, 13, 10.	1.8	21
41	A comparison of home care quality indicator rates in two Canadian provinces. <i>BMC Health Services Research</i> , 2014, 14, 37.	2.2	20
42	Data sharing between home care professionals: a feasibility study using the RAI Home Care instrument. <i>BMC Geriatrics</i> , 2014, 14, 81.	2.7	8
43	Development of a case-mix funding system for adults with combined vision and hearing loss. <i>BMC Health Services Research</i> , 2013, 13, 137.	2.2	5
44	The Role of Medications in Predicting Activity Restriction Due to a Fear of Falling. <i>Journal of Aging and Health</i> , 2012, 24, 269-286.	1.7	9
45	Reliability of standardized assessment for adults who are deafblind. <i>Journal of Rehabilitation Research and Development</i> , 2011, 48, 545.	1.6	19
46	Risk Factors for Restriction in Activity Associated With Fear of Falling Among Seniors Within the Community. <i>Journal of Patient Safety</i> , 2010, 6, 187-191.	1.7	63
47	A Newly Identified Impairment in Both Vision and Hearing Increases the Risk of Deterioration in Both Communication and Cognitive Performance. <i>Canadian Journal on Aging</i> , 0, , 1-14.	1.1	4