

# Jianguo Zhuang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8734178/publications.pdf>

Version: 2024-02-01

13  
papers

1,285  
citations

1040056

9  
h-index

1125743

13  
g-index

13  
all docs

13  
docs citations

13  
times ranked

1582  
citing authors

#	ARTICLE	IF	CITATIONS
1	Distinct Caspase Cascades Are Initiated in Receptor-mediated and Chemical-induced Apoptosis. <i>Journal of Biological Chemistry</i> , 1999, 274, 5053-5060.	3.4	729
2	Mcl-1 Interacts with Truncated Bid and Inhibits Its Induction of Cytochrome c Release and Its Role in Receptor-mediated Apoptosis. <i>Journal of Biological Chemistry</i> , 2006, 281, 5750-5759.	3.4	155
3	Apoptosis, in human monocytic THP.1 cells, results in the release of cytochrome c from mitochondria prior to their ultracondensation, formation of outer membrane discontinuities and reduction in inner membrane potential. <i>Cell Death and Differentiation</i> , 1998, 5, 953-962.	11.2	88
4	Dissociation of Phagocyte Recognition of Cells Undergoing Apoptosis from Other Features of the Apoptotic Program. <i>Journal of Biological Chemistry</i> , 1998, 273, 15628-15632.	3.4	70
5	Characterisation of Mcl-1 cleavage during apoptosis of haematopoietic cells. <i>British Journal of Haematology</i> , 2004, 125, 655-665.	2.5	69
6	Akt is activated in chronic lymphocytic leukemia cells and delivers a pro-survival signal: the therapeutic potential of Akt inhibition. <i>Haematologica</i> , 2010, 95, 110-118.	3.5	69
7	Redistribution of Cytochrome c Precedes the Caspase-Dependent Formation of Ultracondensed Mitochondria, with a Reduced Inner Membrane Potential, in Apoptotic Monocytes. <i>American Journal of Pathology</i> , 1999, 155, 607-618.	3.8	33
8	Total Proteome Analysis Identifies Migration Defects as a Major Pathogenetic Factor in Immunoglobulin Heavy Chain Variable Region (IGHV)-unmutated Chronic Lymphocytic Leukemia. <i>Molecular and Cellular Proteomics</i> , 2015, 14, 933-945.	3.8	31
9	Glucocorticoid resistance in chronic lymphocytic leukaemia is associated with a failure of upregulated Bim/Bcl-2 complexes to activate Bax and Bak. <i>Cell Death and Disease</i> , 2012, 3, e372-e372.	6.3	16
10	Selective IAP inhibition results in sensitization of unstimulated but not CD40-stimulated chronic lymphocytic leukaemia cells to TRAIL-induced apoptosis. <i>Pharmacology Research and Perspectives</i> , 2014, 2, e00081.	2.4	9
11	Delineating the distinct role of AKT in mediating cell survival and proliferation induced by CD154 and IL-4/IL-21 in chronic lymphocytic leukemia. <i>Oncotarget</i> , 2017, 8, 102948-102964.	1.8	9
12	Assessing technical and biological variation in SWATH-MS-based proteomic analysis of chronic lymphocytic leukaemia cells. <i>Scientific Reports</i> , 2021, 11, 2932.	3.3	5
13	Development of a cell-line model to mimic the pro-survival effect of nurse-like cells in chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2021, 62, 45-57.	1.3	2