

Margaret Maxwell

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8769434/publications.pdf>

Version: 2024-02-01

34
papers

2,481
citations

361045

20
h-index

377514

34
g-index

34
all docs

34
docs citations

34
times ranked

3600
citing authors

#	ARTICLE	IF	CITATIONS
1	The consultation and relational empathy (CARE) measure: development and preliminary validation and reliability of an empathy-based consultation process measure. <i>Family Practice</i> , 2004, 21, 699-705.	0.8	601
2	Effectiveness of Motivational Interviewing on adult behaviour change in health and social care settings: A systematic review of reviews. <i>PLoS ONE</i> , 2018, 13, e0204890.	1.1	236
3	Improving reporting of meta-ethnography: the eMERGe reporting guidance. <i>BMC Medical Research Methodology</i> , 2019, 19, 25.	1.4	224
4	Best Practice Elements of Multilevel Suicide Prevention Strategies. <i>Crisis</i> , 2011, 32, 319-333.	0.9	214
5	What Are Reasons for the Large Gender Differences in the Lethality of Suicidal Acts? An Epidemiological Analysis in Four European Countries. <i>PLoS ONE</i> , 2015, 10, e0129062.	1.1	151
6	A methodological systematic review of what's wrong with meta-ethnography reporting. <i>BMC Medical Research Methodology</i> , 2014, 14, 119.	1.4	132
7	Barriers and facilitators to the implementation of the advanced nurse practitioner role in primary care settings: A scoping review. <i>International Journal of Nursing Studies</i> , 2020, 104, 103443.	2.5	106
8	Optimizing Suicide Prevention Programs and Their Implementation in Europe (OSPI Europe): an evidence-based multi-level approach. <i>BMC Public Health</i> , 2009, 9, 428.	1.2	104
9	The influence of contextual factors on healthcare quality improvement initiatives: a realist review. <i>Systematic Reviews</i> , 2020, 9, 94.	2.5	83
10	The influence of contextual factors on healthcare quality improvement initiatives: what works, for whom and in what setting? Protocol for a realist review. <i>Systematic Reviews</i> , 2017, 6, 168.	2.5	76
11	Improving reporting of meta-ethnography: The eMERGe reporting guidance. <i>Journal of Advanced Nursing</i> , 2019, 75, 1126-1139.	1.5	58
12	"Doing the right thing": factors influencing GP prescribing of antidepressants and prescribed doses. <i>BMC Family Practice</i> , 2017, 18, 72.	2.9	55
13	Women's experiences of receiving care for pelvic organ prolapse: a qualitative study. <i>BMC Women's Health</i> , 2019, 19, 45.	0.8	47
14	Exploring synergistic interactions and catalysts in complex interventions: longitudinal, mixed methods case studies of an optimised multi-level suicide prevention intervention in four European countries (Ospi-Europe). <i>BMC Public Health</i> , 2016, 16, 268.	1.2	44
15	The Patient Centered Assessment Method (PCAM): Integrating the Social Dimensions of Health into Primary Care. <i>Journal of Comorbidity</i> , 2015, 5, 110-119.	3.9	39
16	Epidemiology of emergency ambulance service calls related to mental health problems and self harm: a national record linkage study. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2019, 27, 34.	1.1	39
17	A qualitative study of primary care professionals' views of case finding for depression in patients with diabetes or coronary heart disease in the UK. <i>BMC Family Practice</i> , 2013, 14, 46.	2.9	37
18	Developing social capital in implementing a complex intervention: a process evaluation of the early implementation of a suicide prevention intervention in four European countries. <i>BMC Public Health</i> , 2013, 13, 158.	1.2	35

#	ARTICLE	IF	CITATIONS
19	Improving reporting of meta-ethnography: The eMERGe reporting guidance. <i>Psycho-Oncology</i> , 2019, 28, 447-458.	1.0	34
20	Prevention of suicidal behaviour: Results of a controlled community-based intervention study in four European countries. <i>PLoS ONE</i> , 2019, 14, e0224602.	1.1	27
21	Effectiveness of Depression- Suicidal Behaviour Gatekeeper Training among police officers in three European regions: Outcomes of the Optimising Suicide Prevention Programmes and Their Implementation in Europe (OSPI-Europe) study. <i>International Journal of Social Psychiatry</i> , 2016, 62, 651-660.	1.6	23
22	Depression Awareness and Self-Management Through the Internet: Protocol for an Internationally Standardized Approach. <i>JMIR Research Protocols</i> , 2015, 4, e99.	0.5	22
23	Effectiveness of General Practitioner training to improve suicide awareness and knowledge and skills towards depression. <i>Journal of Affective Disorders</i> , 2018, 227, 17-23.	2.0	21
24	Developing a reporting guideline to improve meta-ethnography in health research: the eMERGe mixed-methods study. <i>Health Services and Delivery Research</i> , 2019, 7, 1-116.	1.4	16
25	Improving reporting of meta-ethnography: The eMERGe reporting guidance. <i>Review of Education</i> , 2019, 7, 430-451.	1.1	14
26	PROPEL: implementation of an evidence based pelvic floor muscle training intervention for women with pelvic organ prolapse: a realist evaluation and outcomes study protocol. <i>BMC Health Services Research</i> , 2017, 17, 843.	0.9	10
27	An analysis of suicide trends in Scotland 1950-2014: comparison with England & Wales. <i>BMC Public Health</i> , 2017, 17, 970.	1.2	9
28	The Patient Centred Assessment Method for improving nurse-led biopsychosocial assessment of patients with long-term conditions: a feasibility RCT. <i>Health Services and Delivery Research</i> , 2018, 6, 1-120.	1.4	7
29	Pelvic floor muscle training for women with pelvic organ prolapse: the PROPEL realist evaluation. <i>Health Services and Delivery Research</i> , 2020, 8, 1-104.	1.4	6
30	Prevention and management of depression in primary care in Europe: a holistic model of care and interventions-position paper of the European Forum for Primary Care. <i>Quality in Primary Care</i> , 2008, 16, 187-96.	0.8	3
31	Survey to identify research priorities for primary care in Scotland during and following the COVID-19 pandemic. <i>BMJ Open</i> , 2022, 12, e056817.	0.8	3
32	Targeted education for general practitioners reduces risk of depression or suicide ideation or attempts in older primary care patients. <i>Evidence-Based Mental Health</i> , 2013, 16, 8-8.	2.2	2
33	Implementing pelvic floor muscle training for women with pelvic organ prolapse: a realist evaluation of different delivery models. <i>BMC Health Services Research</i> , 2020, 20, 910.	0.9	2
34	Evaluation of the impact and implementation of a national clinical programme for the management of self-harm in hospital emergency departments: study protocol for a natural experiment. <i>BMJ Open</i> , 2021, 11, e055962.	0.8	1