
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8780894/publications.pdf Version: 2024-02-01



FELLY ANCST

#	Article	IF	CITATIONS
1	Smallest detectable and minimal clinically important differences of rehabilitation intervention with their implications for required sample sizes using WOMAC and SF-36 quality of life measurement instruments in patients with osteoarthritis of the lower extremities. Arthritis and Rheumatism, 2001, MeasureSof.adult shoulder function: Disabilities of the Arm, Shoulder, and Hand Questionnaire	6.7	632
2	(DASH) and Its Short Version ( <i>Quick</i> DASH), Shoulder Pain and Disability Index (SPADI), American Shoulder and Elbow Surgeons (ASES) Society Standardized Shoulder Assessment Form, Constant (Murley) Score (CS), Simple Shoulder Test (SST), Oxford Shoulder Score (OSS), Shoulder Disability Questionnaire (SDQ), and Western Ontario Shoulder Instability Index (WOSI). Arthritis Care and	3.4	447
3	Research, 2011, 63, S174-88. Minimal clinically important rehabilitation effects in patients with osteoarthritis of the lower extremities. Journal of Rheumatology, 2002, 29, 131-8.	2.0	188
4	The minimal clinically important difference raised the significance of outcome effects above the statistical level, with methodological implications for future studies. Journal of Clinical Epidemiology, 2017, 82, 128-136.	5.0	183
5	Prediction of grip and key pinch strength in 978 healthy subjects. BMC Musculoskeletal Disorders, 2010, 11, 94.	1.9	147
6	Motor and Functional Recovery After Stroke. Stroke, 2007, 38, 2101-2107.	2.0	115
7	Responsiveness of six outcome assessment instruments in total shoulder arthroplasty. Arthritis and Rheumatism, 2008, 59, 391-398.	6.7	110
8	Anxiety and depression in the first six months after stroke. A longitudinal multicentre study. Disability and Rehabilitation, 2008, 30, 1858-1866.	1.8	100
9	Comprehensive assessment of clinical outcome and quality of life after total shoulder arthroplasty: Usefulness and validity of subjective outcome measures. Arthritis and Rheumatism, 2004, 51, 819-828.	6.7	99
10	Responsiveness of five condition-specific and generic outcome assessment instruments for chronic pain. BMC Medical Research Methodology, 2008, 8, 26.	3.1	72
11	Translation, Cross-Cultural Adaptation, Reliability, and Validity of the German Version of the Coping Strategies Questionnaire (CSQ-D). Journal of Pain, 2006, 7, 327-336.	1.4	69
12	Comprehensive assessment of clinical outcome and quality of life after total elbow arthroplasty. Arthritis and Rheumatism, 2005, 53, 73-82.	6.7	62
13	How sharp is the short QuickDASH? A refined content and validity analysis of the short form of the disabilities of the shoulder, arm and hand questionnaire in the strata of symptoms and function and specific joint conditions. Quality of Life Research, 2009, 18, 1043-1051.	3.1	62
14	Determinants of Patient Satisfaction After Orthopedic Interventions to the Hand: A Review of the Literature. Journal of Hand Therapy, 2011, 24, 303-312.e10.	1.5	59
15	Interdisciplinary Rehabilitation in Fibromyalgia and Chronic Back Pain: A Prospective Outcome Study. Journal of Pain, 2006, 7, 807-815.	1.4	54
16	Short-term results after reversed shoulder arthroplasty (Delta III) in patients with rheumatoid arthritis and irreparable rotator cuff tear. International Orthopaedics, 2010, 34, 71-77.	1.9	54
17	What Counts: Outcome Assessment After Distal Radius Fractures in Aged Patients. Journal of Orthopaedic Trauma, 2008, 22, S126-S130.	1.4	53
18	Reliability and Validity of the German Version of "The Patient-rated Wrist Evaluation (PRWE)―as an Outcome Measure of Wrist Pain and Disability in Patients with Acute Distal Radius Fractures. Journal of Hand Therapy, 2008, 21, 366-376.	1.5	50

#	Article	IF	CITATIONS
19	Comprehensive assessment of clinical outcome and quality of life after resection interposition arthroplasty of the thumb saddle joint. Arthritis and Rheumatism, 2005, 53, 205-213.	6.7	46
20	Lessons learned during the cross-cultural adaptation of the American Shoulder and Elbow Surgeons shoulder form into German. Journal of Shoulder and Elbow Surgery, 2008, 17, 248-254.	2.6	42
21	Multidimensional minimal clinically important differences in knee osteoarthritis after comprehensive rehabilitation: a prospective evaluation from the Bad Zurzach Osteoarthritis Study. RMD Open, 2018, 4, e000685.	3.8	42
22	Extended overview of the longitudinal pain-depression association: A comparison of six cohorts treated for specific chronic pain conditions. Journal of Affective Disorders, 2020, 273, 508-516.	4.1	39
23	A concept for comprehensively measuring health, function and quality of life following orthopaedic interventions of the upper extremity. Archives of Orthopaedic and Trauma Surgery, 2009, 129, 113-118.	2.4	38
24	The factor subdimensions of the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) help to specify hip and knee osteoarthritis. a prospective evaluation and validation study. Journal of Rheumatology, 2005, 32, 1324-30.	2.0	38
25	Does Classification of Persons with Fibromyalgia into Multidimensional Pain Inventory Subgroups Detect Differences in Outcome after a Standard Chronic Pain Management Program?. Pain Research and Management, 2009, 14, 445-453.	1.8	33
26	ICF Core Sets for diabetes mellitus. Journal of Rehabilitation Medicine, 2004, 36, 100-106.	1.1	31
27	Clinical effectiveness of an interdisciplinary pain management programme compared with standard inpatient rehabilitation in chronic pain: A naturalistic, prospective controlled cohort study. Journal of Rehabilitation Medicine, 2009, 41, 569-575.	1.1	28
28	Reliability of the Multidimensional Pain Inventory and stability of the MPI classification system in chronic back pain. BMC Musculoskeletal Disorders, 2012, 13, 155.	1.9	27
29	Responsiveness of five outcome measurement instruments in total elbow arthroplasty. Arthritis Care and Research, 2012, 64, 1749-1755.	3.4	26
30	Multidimensional associative factors for improvement in pain, function, and working capacity after rehabilitation of whiplash associated disorder: a prognostic, prospective outcome study. BMC Musculoskeletal Disorders, 2014, 15, 130.	1.9	25
31	High molecular weight Intraarticular hyaluronic acid for the treatment of knee osteoarthritis: a network meta-analysis. BMC Musculoskeletal Disorders, 2020, 21, 702.	1.9	25
32	The American Shoulder and Elbow Surgeons Elbow Questionnaire: Cross-cultural Adaptation into German and Evaluation of Its Psychometric Properties. Journal of Hand Therapy, 2010, 23, 301-314.	1.5	22
33	Association of the sense of coherence with physical and psychosocial health in the rehabilitation of osteoarthritis of the hip and knee: a prospective cohort study. BMC Musculoskeletal Disorders, 2013, 14, 159.	1.9	22
34	Differences in pain, function and coping in Multidimensional Pain Inventory subgroups of chronic back pain: a one-group pretest-posttest study. BMC Musculoskeletal Disorders, 2011, 12, 145.	1.9	21
35	Epidemiology of Back Pain in Young and Middle-Aged Adults: A Longitudinal Population Cohort Survey From Age 27–50 Years. Psychosomatics, 2017, 58, 604-613.	2.5	21
36	Evaluation of wrist and hand handicap and postoperative outcome in rheumatoid arthritis. Hand Clinics, 2003, 19, 471-481.	1.0	20

#	Article	IF	CITATIONS
37	Quality of life assessment in osteoarthritis. Expert Review of Pharmacoeconomics and Outcomes Research, 2003, 3, 623-636.	1.4	20
38	Refined Insights Into the Pain-Depression Association in Chronic Pain Patients. Clinical Journal of Pain, 2008, 24, 808-816.	1.9	19
39	Health and quality of life in patients with primary and secondary lymphedema of the lower extremity. Vasa - European Journal of Vascular Medicine, 2015, 44, 129-137.	1.4	19
40	Prognostic Factors for Pain Relief and Functional Improvement in Chronic Pain After Inpatient Rehabilitation. Clinical Journal of Pain, 2014, 30, 279-285.	1.9	16
41	Responsiveness of the cervical Northern American Spine Society questionnaire (NASS) and the Short Form 36 (SF-36) in chronic whiplash. Clinical Rehabilitation, 2012, 26, 142-151.	2.2	15
42	Effectiveness of subgroup-specific pain rehabilitation: a randomized controlled trial in patients with chronic back pain. European Journal of Physical and Rehabilitation Medicine, 2018, 54, 358-370.	2.2	15
43	Multidisciplinary biopsychosocial rehabilitation in chronic neck pain: a naturalistic prospective cohort study with intraindividual control of effects and 12-month follow-up. European Journal of Physical and Rehabilitation Medicine, 2019, 55, 665-675.	2.2	15
44	Interdisciplinary rehabilitation after whiplash injury: An observational prospective outcome study. Journal of Rehabilitation Medicine, 2010, 42, 350-356.	1,1	14
45	Effects of Inpatient Rehabilitation in Hip and Knee Osteoarthritis: A Naturalistic Prospective Cohort Study With Intraindividual Control of Effects. Archives of Physical Medicine and Rehabilitation, 2013, 94, 2139-2145.	0.9	14
46	Health and quality of life in patients with medication overuse headache syndrome after standardized inpatient rehabilitation. Medicine (United States), 2017, 96, e8493.	1.0	13
47	Cross-sectional validity and specificity of comprehensive measurement in lymphedema and lipedema of the lower extremity: a comparison of five outcome instruments. Health and Quality of Life Outcomes, 2020, 18, 245.	2.4	13
48	Long-term survival of GSB III elbow prostheses and risk factors for revisions. Archives of Orthopaedic and Trauma Surgery, 2013, 133, 1415-1424.	2.4	12
49	MCID — The Minimal Clinically Important Difference Assigns Significance to Outcome Effects. Journal of Rheumatology, 2016, 43, 258-259.	2.0	11
50	Translation, transcultural adaptation, reliability and validity of the Patient Satisfaction Questionnaire in German. Physiotherapy, 2008, 94, 43-55.	0.4	10
51	Common and Contrasting Characteristics of the Chronic Soft-Tissue Pain Conditions Fibromyalgia and Lipedema. Journal of Pain Research, 2021, Volume 14, 2931-2941.	2.0	9
52	Prognostic factors for the improvement of pain and disability following multidisciplinary rehabilitation in patients with chronic neck pain. BMC Musculoskeletal Disorders, 2021, 22, 330.	1.9	8
53	Catastrophizing as a prognostic factor for pain and physical function in the multidisciplinary rehabilitation of fibromyalgia and low back pain. European Journal of Pain, 2022, 26, 1569-1580.	2.8	8
54	Acupuncture Improves Peripheral Perfusion in Patients with Reflex Sympathetic Dystrophy. Journal of Clinical Rheumatology, 2002, 8, 6-12.	0.9	7

#	Article	IF	CITATIONS
55	Interdisciplinary rehabilitation after whiplash injury. Medicine (United States), 2017, 96, e6113.	1.0	7
56	Comparison of short- and mid-term outcomes of Italian- and German-speaking patients after an interdisciplinary pain management programme in Switzerland: A prospective cohort study. Journal of Rehabilitation Medicine, 2019, 51, 127-135.	1.1	7
57	Comprehensiveness and validity of a multidimensional assessment in patients with chronic low back pain: a prospective cohort study. BMC Musculoskeletal Disorders, 2021, 22, 291.	1.9	7
58	Mental and psychosocial health and health related quality of life before and after cardiac rehabilitation: a prospective cohort study with comparison to specific population norms. Health and Quality of Life Outcomes, 2022, 20, .	2.4	2
59	Quadriceps performance under activation of foot dorsal extension in healthy volunteers: an interventional cohort study. BMC Musculoskeletal Disorders, 2015, 16, 340.	1.9	1
60	Disabilities of the Arm, Shoulder and Hand Questionnaire (DASH). , 2021, , 1-22.		0
61	Begleitende Messungen zur QualitÃ <b>t</b> ssicherung bei Ellenbogenprothetik. , 2013, , 217-227.		0