

Susan M O'brien

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8801980/publications.pdf>

Version: 2024-02-01

81
papers

14,673
citations

31902

53
h-index

60497

81
g-index

81
all docs

81
docs citations

81
times ranked

10847
citing authors

#	ARTICLE	IF	CITATIONS
1	Idelalisib and Rituximab in Relapsed Chronic Lymphocytic Leukemia. <i>New England Journal of Medicine</i> , 2014, 370, 997-1007.	13.9	1,535
2	Inotuzumab Ozogamicin versus Standard Therapy for Acute Lymphoblastic Leukemia. <i>New England Journal of Medicine</i> , 2016, 375, 740-753.	13.9	1,047
3	Safety and activity of blinatumomab for adult patients with relapsed or refractory B-precursor acute lymphoblastic leukaemia: a multicentre, single-arm, phase 2 study. <i>Lancet Oncology</i> , The, 2015, 16, 57-66.	5.1	1,031
4	Long-term follow-up results of hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone (Hyper-CVAD), a dose-intensive regimen, in adult acute lymphocytic leukemia. <i>Cancer</i> , 2004, 101, 2788-2801.	2.0	550
5	Results of intensive chemotherapy in 998 patients age 65 years or older with acute myeloid leukemia or high-risk myelodysplastic syndrome. <i>Cancer</i> , 2006, 106, 1090-1098.	2.0	550
6	Treatment of Philadelphia chromosome-positive acute lymphocytic leukemia with hyper-CVAD and imatinib mesylate. <i>Blood</i> , 2004, 103, 4396-4407.	0.6	522
7	Chemoimmunotherapy with hyper-CVAD plus rituximab for the treatment of adult Burkitt and Burkitt-type lymphoma or acute lymphoblastic leukemia. <i>Cancer</i> , 2006, 106, 1569-1580.	2.0	503
8	Inotuzumab ozogamicin, an anti-CD22 ^{â€} calcicheamicin conjugate, for refractory and relapsed acute lymphocytic leukaemia: a phase 2 study. <i>Lancet Oncology</i> , The, 2012, 13, 403-411.	5.1	401
9	Chemoimmunotherapy With a Modified Hyper-CVAD and Rituximab Regimen Improves Outcome in De Novo Philadelphia Chromosome ^{â€} Negative Precursor B-Lineage Acute Lymphoblastic Leukemia. <i>Journal of Clinical Oncology</i> , 2010, 28, 3880-3889.	0.8	361
10	Phase I/II Study of Combination Therapy With Sorafenib, Idarubicin, and Cytarabine in Younger Patients With Acute Myeloid Leukemia. <i>Journal of Clinical Oncology</i> , 2010, 28, 1856-1862.	0.8	347
11	First report of phase 2 study of dasatinib with hyper-CVAD for the frontline treatment of patients with Philadelphia chromosome ^{â€} positive (Ph+) acute lymphoblastic leukemia. <i>Blood</i> , 2010, 116, 2070-2077.	0.6	319
12	Safety and activity of ibrutinib plus rituximab for patients with high-risk chronic lymphocytic leukaemia: a single-arm, phase 2 study. <i>Lancet Oncology</i> , The, 2014, 15, 1090-1099.	5.1	315
13	Outcomes of patients with chronic lymphocytic leukemia after discontinuing ibrutinib. <i>Blood</i> , 2015, 125, 2062-2067.	0.6	303
14	Improved survival in chronic myeloid leukemia since the introduction of imatinib therapy: a single-institution historical experience. <i>Blood</i> , 2012, 119, 1981-1987.	0.6	298
15	Clonal evolution in patients with chronic lymphocytic leukaemia developing resistance to BTK inhibition. <i>Nature Communications</i> , 2016, 7, 11589.	5.8	285
16	Results of inotuzumab ozogamicin, a CD22 monoclonal antibody, in refractory and relapsed acute lymphocytic leukemia. <i>Cancer</i> , 2013, 119, 2728-2736.	2.0	265
17	Early T-cell precursor acute lymphoblastic leukemia/lymphoma (ETP-ALL/LBL) in adolescents and adults: a high-risk subtype. <i>Blood</i> , 2016, 127, 1863-1869.	0.6	253
18	Combination of hyper-CVAD with ponatinib as first-line therapy for patients with Philadelphia chromosome-positive acute lymphoblastic leukaemia: a single-centre, phase 2 study. <i>Lancet Oncology</i> , The, 2015, 16, 1547-1555.	5.1	245

#	ARTICLE	IF	CITATIONS
19	Nilotinib As Front-Line Treatment for Patients With Chronic Myeloid Leukemia in Early Chronic Phase. <i>Journal of Clinical Oncology</i> , 2010, 28, 392-397.	0.8	231
20	Results of Dasatinib Therapy in Patients With Early Chronic-Phase Chronic Myeloid Leukemia. <i>Journal of Clinical Oncology</i> , 2010, 28, 398-404.	0.8	227
21	Relative survival in patients with chronic-phase chronic myeloid leukaemia in the tyrosine-kinase inhibitor era: analysis of patient data from six prospective clinical trials. <i>Lancet Haematology</i> , 2015, 2, e186-e193.	2.2	227
22	A phase 2 study of idelalisib plus rituximab in treatment-naïve older patients with chronic lymphocytic leukemia. <i>Blood</i> , 2015, 126, 2686-2694.	0.6	224
23	Complete cytogenetic and molecular responses to interferon- γ -based therapy for chronic myelogenous leukemia are associated with excellent long-term prognosis. <i>Cancer</i> , 2003, 97, 1033-1041.	2.0	219
24	Inotuzumab ozogamicin versus standard of care in relapsed or refractory acute lymphoblastic leukemia: Final report and long-term survival follow-up from the randomized, phase 3 INO-VATE study. <i>Cancer</i> , 2019, 125, 2474-2487.	2.0	210
25	Survival benefit with imatinib mesylate versus interferon- γ -based regimens in newly diagnosed chronic-phase chronic myelogenous leukemia. <i>Blood</i> , 2006, 108, 1835-1840.	0.6	204
26	New insights into the pathophysiology and therapy of adult acute lymphoblastic leukemia. <i>Cancer</i> , 2015, 121, 2517-2528.	2.0	200
27	Impact of complete molecular response on survival in patients with Philadelphia chromosome-positive acute lymphoblastic leukemia. <i>Blood</i> , 2016, 128, 504-507.	0.6	194
28	Final report of a phase II study of imatinib mesylate with hyper-CVAD for the front-line treatment of adult patients with Philadelphia chromosome-positive acute lymphoblastic leukemia. <i>Haematologica</i> , 2015, 100, 653-661.	1.7	191
29	Detection of MRD may predict the outcome of patients with Philadelphia chromosome-positive ALL treated with tyrosine kinase inhibitors plus chemotherapy. <i>Blood</i> , 2013, 122, 1214-1221.	0.6	190
30	Combination of hyper-CVAD with ponatinib as first-line therapy for patients with Philadelphia chromosome-positive acute lymphoblastic leukaemia: long-term follow-up of a single-centre, phase 2 study. <i>Lancet Haematology</i> , 2018, 5, e618-e627.	2.2	190
31	Long-term follow-up of a phase 2 study of chemotherapy plus dasatinib for the initial treatment of patients with Philadelphia chromosome-positive acute lymphoblastic leukemia. <i>Cancer</i> , 2015, 121, 4158-4164.	2.0	181
32	Hepatic adverse event profile of inotuzumab ozogamicin in adult patients with relapsed or refractory acute lymphoblastic leukaemia: results from the open-label, randomised, phase 3 INO-VATE study. <i>Lancet Haematology</i> , 2017, 4, e387-e398.	2.2	158
33	Hyper-CVAD plus ponatinib versus hyper-CVAD plus dasatinib as frontline therapy for patients with Philadelphia chromosome-positive acute lymphoblastic leukemia: A propensity score analysis. <i>Cancer</i> , 2016, 122, 3650-3656.	2.0	156
34	Delayed achievement of cytogenetic and molecular response is associated with increased risk of progression among patients with chronic myeloid leukemia in early chronic phase receiving high-dose or standard-dose imatinib therapy. <i>Blood</i> , 2009, 113, 6315-6321.	0.6	153
35	Acalabrutinib monotherapy in patients with relapsed/refractory chronic lymphocytic leukemia: updated phase 2 results. <i>Blood</i> , 2020, 135, 1204-1213.	0.6	130
36	Long-term outcome of patients with chronic myeloid leukemia treated with second-generation tyrosine kinase inhibitors after imatinib failure is predicted by the in vitro sensitivity of BCR-ABL kinase domain mutations. <i>Blood</i> , 2009, 114, 2037-2043.	0.6	119

#	ARTICLE	IF	CITATIONS
37	Prognostic factors and survival outcomes in patients with chronic myeloid leukemia in blast phase in the tyrosine kinase inhibitor era: Cohort study of 477 patients. <i>Cancer</i> , 2017, 123, 4391-4402.	2.0	114
38	Results of the hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone regimen in elderly patients with acute lymphocytic leukemia. <i>Cancer</i> , 2008, 113, 2097-2101.	2.0	109
39	Defining the course and prognosis of adults with acute lymphocytic leukemia in first salvage after induction failure or short first remission duration. <i>Cancer</i> , 2010, 116, 5568-5574.	2.0	104
40	Long-term outcomes for patients with chronic lymphocytic leukemia who discontinue ibrutinib. <i>Cancer</i> , 2017, 123, 2268-2273.	2.0	103
41	Final results of a single institution experience with a pediatric-based regimen, the augmented Berlin-Frankfurt-Münster, in adolescents and young adults with acute lymphoblastic leukemia, and comparison to the hyper-CVAD regimen. <i>American Journal of Hematology</i> , 2016, 91, 819-823.	2.0	102
42	Minimal residual disease assessed by multi-parameter flow cytometry is highly prognostic in adult patients with acute lymphoblastic leukaemia. <i>British Journal of Haematology</i> , 2016, 172, 392-400.	1.2	102
43	Characteristics and outcome of patients with acute myeloid leukemia refractory to 1 cycle of high-dose cytarabine-based induction chemotherapy. <i>Blood</i> , 2010, 116, 5818-5823.	0.6	93
44	Augmented Berlin-Frankfurt-Münster therapy in adolescents and young adults (AYAs) with acute lymphoblastic leukemia (ALL). <i>Cancer</i> , 2014, 120, 3660-3668.	2.0	91
45	Malignancies occurring during therapy with tyrosine kinase inhibitors (TKIs) for chronic myeloid leukemia (CML) and other hematologic malignancies. <i>Blood</i> , 2011, 118, 4353-4358.	0.6	89
46	Ponatinib as first-line treatment for patients with chronic myeloid leukaemia in chronic phase: a phase 2 study. <i>Lancet Haematology</i> , 2015, 2, e376-e383.	2.2	86
47	Survival is poorer in patients with secondary core-binding factor acute myelogenous leukemia compared with de novo core-binding factor leukemia. <i>Cancer</i> , 2009, 115, 3217-3221.	2.0	76
48	Long-term molecular and cytogenetic response and survival outcomes with imatinib 400 mg, imatinib 800 mg, dasatinib, and nilotinib in patients with chronic-phase chronic myeloid leukaemia: retrospective analysis of patient data from five clinical trials. <i>Lancet Haematology</i> , 2015, 2, e118-e128.	2.2	65
49	Predictive factors for outcome and response in patients treated with second-generation tyrosine kinase inhibitors for chronic myeloid leukemia in chronic phase after imatinib failure. <i>Blood</i> , 2011, 117, 1822-1827.	0.6	64
50	Inotuzumab ozogamicin in combination with low-intensity chemotherapy (mini-HCVAD) with or without blinatumomab versus standard intensive chemotherapy (HCVAD) as frontline therapy for older patients with Philadelphia chromosome-negative acute lymphoblastic leukemia: A propensity score analysis. <i>Cancer</i> , 2019, 125, 2579-2586.	2.0	63
51	Outcome of patients with acute myelogenous leukemia after second salvage therapy. <i>Cancer</i> , 2005, 104, 547-554.	2.0	61
52	Significance of Increasing Levels of Minimal Residual Disease in Patients With Philadelphia Chromosome-Positive Chronic Myelogenous Leukemia in Complete Cytogenetic Response. <i>Journal of Clinical Oncology</i> , 2009, 27, 3659-3663.	0.8	61
53	Augmented Hyper-CVAD Based on Dose-Intensified Vincristine, Dexamethasone, and Asparaginase in Adult Acute Lymphoblastic Leukemia Salvage Therapy. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2011, 11, 54-59.	0.2	61
54	HCVAD plus imatinib or dasatinib in lymphoid blastic phase chronic myeloid leukemia. <i>Cancer</i> , 2014, 120, 373-380.	2.0	54

#	ARTICLE	IF	CITATIONS
55	Phase 3 randomized, placebo-controlled, double-blind study of high-dose continuous infusion cytarabine alone or with laromustine (VNP40101M) in patients with acute myeloid leukemia in first relapse. <i>Blood</i> , 2009, 114, 4027-4033.	0.6	52
56	Acute lymphoblastic leukemia in adolescents and young adults. <i>Cancer</i> , 2017, 123, 2398-2403.	2.0	49
57	Prognostic impact of pretreatment cytogenetics in adult Philadelphia chromosome-negative acute lymphoblastic leukemia in the era of minimal residual disease. <i>Cancer</i> , 2017, 123, 459-467.	2.0	49
58	Efficacy and safety analysis by age cohort of inotuzumab ozogamicin in patients with relapsed or refractory acute lymphoblastic leukemia enrolled in INO-VATE. <i>Cancer</i> , 2018, 124, 1722-1732.	2.0	43
59	Poor outcomes associated with +der(22)t(9;22) and $\hat{9}/9p$ in patients with Philadelphia chromosome-positive acute lymphoblastic leukemia receiving chemotherapy plus a tyrosine kinase inhibitor. <i>American Journal of Hematology</i> , 2017, 92, 238-243.	2.0	41
60	Efficacy of Ponatinib Versus Earlier Generation Tyrosine Kinase Inhibitors for Front-line Treatment of Newly Diagnosed Philadelphia-positive Acute Lymphoblastic Leukemia. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2018, 18, 257-265.	0.2	39
61	A randomized phase 2 study of idarubicin and cytarabine with clofarabine or fludarabine in patients with newly diagnosed acute myeloid leukemia. <i>Cancer</i> , 2017, 123, 4430-4439.	2.0	37
62	Outcomes of patients with chronic lymphocytic leukemia treated with first-line idelalisib plus rituximab after cessation of treatment for toxicity. <i>Cancer</i> , 2016, 122, 2505-2511.	2.0	31
63	Conditional survival in patients with chronic myeloid leukemia in chronic phase in the era of tyrosine kinase inhibitors. <i>Cancer</i> , 2016, 122, 238-248.	2.0	30
64	Philadelphia chromosome-positive acute lymphoblastic leukemia at first relapse in the era of tyrosine kinase inhibitors. <i>American Journal of Hematology</i> , 2019, 94, 1388-1395.	2.0	26
65	The early achievement of measurable residual disease negativity in the treatment of adults with Philadelphia-negative B-cell acute lymphoblastic leukemia is a strong predictor for survival. <i>American Journal of Hematology</i> , 2020, 95, 144-150.	2.0	25
66	Long-term follow-up of salvage therapy using a combination of inotuzumab ozogamicin and mini-hyper-CVD with or without blinatumomab in relapsed/refractory Philadelphia chromosome-negative acute lymphoblastic leukemia. <i>Cancer</i> , 2021, 127, 2025-2038.	2.0	24
67	Outcomes of acute lymphoblastic leukemia with <i>KMT2A</i> (<i>MLL</i>) rearrangement: the MD Anderson experience. <i>Blood Advances</i> , 2021, 5, 5415-5419.	2.5	24
68	Results of second salvage therapy in 673 adults with acute myelogenous leukemia treated at a single institution since 2000. <i>Cancer</i> , 2018, 124, 2534-2540.	2.0	23
69	Prediction for sustained deep molecular response of <i>BCR-ABL1</i> levels in patients with chronic myeloid leukemia in chronic phase. <i>Cancer</i> , 2018, 124, 1160-1168.	2.0	23
70	Frontline Inotuzumab Ozogamicin in Combination with Low-Intensity Chemotherapy (mini-hyper-CVD) for Older Patients with Acute Lymphoblastic Leukemia (ALL). <i>Blood</i> , 2015, 126, 83-83.	0.6	19
71	<i>TP53</i> mutation does not confer a poor outcome in adult patients with acute lymphoblastic leukemia who are treated with frontline hyper-CVD-based regimens. <i>Cancer</i> , 2017, 123, 3717-3724.	2.0	18
72	Prognostic significance of day 14 bone marrow evaluation in adults with Philadelphia chromosome-negative acute lymphoblastic leukemia. <i>Cancer</i> , 2016, 122, 3812-3820.	2.0	17

#	ARTICLE	IF	CITATIONS
73	Prognostic implications of cytogenetics in adults with acute lymphoblastic leukemia treated with inotuzumab ozogamicin. <i>American Journal of Hematology</i> , 2019, 94, 408-416.	2.0	11
74	Inotuzumab Ozogamicin in Combination with Low-Intensity Chemotherapy (mini-hyper-CVD) As Frontline Therapy for Older Patients with Acute Lymphoblastic Leukemia (ALL): Interim Result of a Phase II Clinical Trial. <i>Blood</i> , 2016, 128, 588-588.	0.6	11
75	Hyper- Δ CVAD plus ofatumumab versus hyper- Δ CVAD plus rituximab as frontline therapy in adults with Philadelphia chromosome- Δ negative acute lymphoblastic leukemia: A propensity score analysis. <i>Cancer</i> , 2021, 127, 3381-3389.	2.0	10
76	Discontinuation of Maintenance Tyrosine Kinase Inhibitors in Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia outside of Transplant. <i>Acta Haematologica</i> , 2021, 144, 285-292.	0.7	10
77	The safety of Bruton's tyrosine kinase inhibitors for the treatment of chronic lymphocytic leukemia. <i>Expert Opinion on Drug Safety</i> , 2017, 16, 1079-1088.	1.0	9
78	Using ibrutinib in earlier lines of treatment results in better outcomes for patients with chronic lymphocytic leukemia/small lymphocytic lymphoma. <i>Leukemia and Lymphoma</i> , 2021, 62, 3278-3282.	0.6	7
79	Similar Outcome of Patients With Chronic Myeloid Leukemia Treated With Imatinib in or Out of Clinical Trials. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2013, 13, 693-699.	0.2	6
80	Chronic myeloid leukemia among patients with a history of prior malignancies: A tale of dual survivorship. <i>Cancer</i> , 2017, 123, 609-616.	2.0	4
81	Sustained long-lasting responses after lenalidomide discontinuation in patients with chronic lymphocytic leukemia. <i>Leukemia</i> , 2018, 32, 2278-2281.	3.3	3