

Seung Min Jung

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8839693/publications.pdf>

Version: 2024-02-01

123
papers

2,005
citations

257450

24
h-index

345221

36
g-index

125
all docs

125
docs citations

125
times ranked

3527
citing authors

#	ARTICLE	IF	CITATIONS
1	Cytokine-Mediated Bone Destruction in Rheumatoid Arthritis. <i>Journal of Immunology Research</i> , 2014, 2014, 1-15.	2.2	119
2	2018 update of the APLAR recommendations for treatment of rheumatoid arthritis. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 357-375.	1.9	115
3	<scp>APLAR</scp> rheumatoid arthritis treatment recommendations. <i>International Journal of Rheumatic Diseases</i> , 2015, 18, 685-713.	1.9	109
4	Role of intrahepatic tumor control in the prognosis of patients with hepatocellular carcinoma and extrahepatic metastases. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2012, 27, 684-689.	2.8	61
5	Cord blood cell-derived iPSCs as a new candidate for chondrogenic differentiation and cartilage regeneration. <i>Stem Cell Research and Therapy</i> , 2017, 8, 16.	5.5	59
6	Targeted Immunotherapy for Autoimmune Disease. <i>Immune Network</i> , 2022, 22, e9.	3.6	46
7	Generation of disease-specific induced pluripotent stem cells from patients with rheumatoid arthritis and osteoarthritis. <i>Arthritis Research and Therapy</i> , 2014, 16, R41.	3.5	44
8	Risk of tuberculosis in patients treated with anti-tumor necrosis factor therapy: a nationwide study in South Korea, a country with an intermediate tuberculosis burden. <i>International Journal of Rheumatic Diseases</i> , 2015, 18, 323-330.	1.9	43
9	Downregulation of Tryptophan-related Metabolomic Profile in Rheumatoid Arthritis Synovial Fluid. <i>Journal of Rheumatology</i> , 2015, 42, 2003-2011.	2.0	43
10	Application of the 2016 EULAR/ACR/PRINTO Classification Criteria for Macrophage Activation Syndrome in Patients with Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2017, 44, 996-1003.	2.0	43
11	Systemic immune inflammation index could estimate the cross-sectional high activity and the poor outcomes in immunosuppressive drug-naïve patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Nephrology</i> , 2019, 24, 711-717.	1.6	42
12	Recent progress of national banking project on homozygous <scp>HLA</scp> -typed induced pluripotent stem cells in <scp>S</scp>outh <scp>K</scp>orea. <i>Journal of Tissue Engineering and Regenerative Medicine</i> , 2018, 12, e1531-e1536.	2.7	39
13	Arthritic role of <i>Porphyromonas gingivalis</i> in collagen-induced arthritis mice. <i>PLoS ONE</i> , 2017, 12, e0188698.	2.5	39
14	Etanercept-Synthesising Mesenchymal Stem Cells Efficiently Ameliorate Collagen-Induced Arthritis. <i>Scientific Reports</i> , 2017, 7, 39593.	3.3	37
15	Evaluation of Spleen Glucose Metabolism Using ¹⁸ F-FDG PET/CT in Patients with Febrile Autoimmune Disease. <i>Journal of Nuclear Medicine</i> , 2017, 58, 507-513.	5.0	33
16	Recent technological updates and clinical applications of induced pluripotent stem cells. <i>Korean Journal of Internal Medicine</i> , 2014, 29, 547.	1.7	32
17	Neutrophil to lymphocyte ratio at diagnosis can estimate vasculitis activity and poor prognosis in patients with ANCA-associated vasculitis: a retrospective study. <i>BMC Nephrology</i> , 2018, 19, 187.	1.8	32
18	In-hospital mortality in febrile lupus patients based on 2016 EULAR/ACR/PRINTO classification criteria for macrophage activation syndrome. <i>Seminars in Arthritis and Rheumatism</i> , 2017, 47, 216-221.	3.4	28

#	ARTICLE	IF	CITATIONS
19	Platelet to lymphocyte ratio is associated with the current activity of ANCA-associated vasculitis at diagnosis: a retrospective monocentric study. <i>Rheumatology International</i> , 2018, 38, 1865-1871.	3.0	28
20	Interrupting oral infection of <i>Porphyromonas gingivalis</i> with anti-FimA antibody attenuates bacterial dissemination to the arthritic joint and improves experimental arthritis. <i>Experimental and Molecular Medicine</i> , 2018, 50, e460-e460.	7.7	27
21	Safety of Tocilizumab in Rheumatoid Arthritis Patients with Resolved Hepatitis B Virus Infection: Data from Real-World Experience. <i>Yonsei Medical Journal</i> , 2018, 59, 452.	2.2	27
22	A Dual Target-directed Agent against Interleukin-6 Receptor and Tumor Necrosis Factor $\hat{\pm}$ ameliorates experimental arthritis. <i>Scientific Reports</i> , 2016, 6, 20150.	3.3	26
23	Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. <i>Rheumatology International</i> , 2019, 39, 1937-1944.	3.0	26
24	Sodium Chloride Aggravates Arthritis via Th17 Polarization. <i>Yonsei Medical Journal</i> , 2019, 60, 88.	2.2	26
25	1,25-dihydroxy Vitamin D3 and Interleukin-6 Blockade Synergistically Regulate Rheumatoid Arthritis by Suppressing Interleukin-17 Production and Osteoclastogenesis. <i>Journal of Korean Medical Science</i> , 2020, 35, e40.	2.5	26
26	Pulmonary hypertension in systemic lupus erythematosus: an independent predictor of patient survival. <i>Korean Journal of Internal Medicine</i> , 2015, 30, 232.	1.7	26
27	Intraperitoneal infusion of mesenchymal stem cell attenuates severity of collagen antibody induced arthritis. <i>PLoS ONE</i> , 2018, 13, e0198740.	2.5	25
28	The Interleukin 33/ST2 Axis in Patients with Primary Sjögren Syndrome: Expression in Serum and Salivary Glands, and the Clinical Association. <i>Journal of Rheumatology</i> , 2015, 42, 264-271.	2.0	22
29	No overlap between IgG4-related disease and microscopic polyangiitis and granulomatosis with polyangiitis despite elevated serum IgG4 at diagnosis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 1147-1154.	2.2	21
30	Induced production of anti-etanercept antibody in collagen-induced arthritis. <i>Molecular Medicine Reports</i> , 2014, 9, 2301-2308.	2.4	20
31	Low serum complement 3 level is associated with severe ANCA-associated vasculitis at diagnosis. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 223-230.	1.6	20
32	The use of biological disease-modifying antirheumatic drugs for inflammatory arthritis in Korea: results of a Korean Expert Consensus. <i>Korean Journal of Internal Medicine</i> , 2020, 35, 41-59.	1.7	20
33	A New Strategy to Deliver Synthetic Protein Drugs: Self-reproducible Biologics Using Minicircles. <i>Scientific Reports</i> , 2014, 4, 5961.	3.3	19
34	Ethyl pyruvate ameliorates inflammatory arthritis in mice. <i>International Immunopharmacology</i> , 2017, 52, 333-341.	3.8	19
35	Immunogenicity of anti-tumour necrosis factor therapy in Korean patients with rheumatoid arthritis and ankylosing spondylitis. <i>International Immunopharmacology</i> , 2014, 21, 20-25.	3.8	18
36	Eupatilin Ameliorates Collagen Induced Arthritis. <i>Journal of Korean Medical Science</i> , 2015, 30, 233.	2.5	17

#	ARTICLE	IF	CITATIONS
37	Birmingham vasculitis activity score of more than 9.5 at diagnosis is an independent predictor of refractory disease in granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1593-1605.	1.9	17
38	The Use of Biological Disease-modifying Antirheumatic Drugs for Inflammatory Arthritis in Korea: Results of a Korean Expert Consensus. <i>Journal of Rheumatic Diseases</i> , 2020, 27, 4.	1.1	17
39	Role of Fractalkine in the Pathogenesis of Primary Sjögren Syndrome: Increased Serum Levels of Fractalkine, Its Expression in Labial Salivary Glands, and the Association with Clinical Manifestations. <i>Journal of Rheumatology</i> , 2014, 41, 2425-2438.	2.0	16
40	Delta Neutrophil Index Is Associated with Vasculitis Activity and Risk of Relapse in ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018, 59, 397.	2.2	16
41	Prognostic nutritional index is associated with disease severity and relapse in ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 797-804.	1.9	16
42	Integrative analysis of lung molecular signatures reveals key drivers of systemic sclerosis-associated interstitial lung disease. <i>Annals of the Rheumatic Diseases</i> , 2022, 81, 108-116.	0.9	16
43	Clinical and prognostic features of Korean patients with MPO-ANCA, PR3-ANCA and ANCA-negative vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 111-118.	0.8	16
44	The Risk of Osteoporotic Fractures According to the FRAX Model in Korean Patients with Rheumatoid Arthritis. <i>Journal of Korean Medical Science</i> , 2014, 29, 1082.	2.5	15
45	Antinociceptive and anti-inflammatory effects of essential oil extracted from <i>Chamaecyparis obtusa</i> in mice. <i>International Immunopharmacology</i> , 2015, 29, 320-325.	3.8	15
46	Decreased muscle mass is independently associated with knee pain in female patients with radiographically mild osteoarthritis: a nationwide cross-sectional study (KNHANES 2010-2011). <i>Clinical Rheumatology</i> , 2018, 37, 1333-1340.	2.2	15
47	Radiographic Structural Damage Is Worse in the Dominant than the Non-Dominant Hand in Individuals with Early Rheumatoid Arthritis. <i>PLoS ONE</i> , 2015, 10, e0135409.	2.5	14
48	Decreased ex vivo production of interferon-gamma is associated with severity and poor prognosis in patients with lupus. <i>Arthritis Research and Therapy</i> , 2017, 19, 193.	3.5	14
49	Comparison of the Clinical Implications among Five Different Nutritional Indices in Patients with Lupus Nephritis. <i>Nutrients</i> , 2019, 11, 1456.	4.1	14
50	The utility of the ACR/EULAR 2017 provisional classification criteria for granulomatosis with polyangiitis in Korean patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 85-87.	0.8	14
51	Risk factors associated with inadequate control of disease activity in elderly patients with rheumatoid arthritis: Results from a nationwide Korean College of Rheumatology BIOlogics (KOBIO) registry. <i>PLoS ONE</i> , 2018, 13, e0205651.	2.5	13
52	Fibrosis-4 index at diagnosis is associated with all-cause mortality in patients with microscopic polyangiitis and granulomatosis with polyangiitis. <i>BMC Gastroenterology</i> , 2019, 19, 90.	2.0	13
53	Risk of malignancy in patients with rheumatoid arthritis after anti-tumor necrosis factor therapy: results from Korean National Health Insurance claims data. <i>Korean Journal of Internal Medicine</i> , 2019, 34, 669-677.	1.7	13
54	Self in vivo production of a synthetic biological drug CTLA4Ig using a minicircle vector. <i>Scientific Reports</i> , 2014, 4, 6935.	3.3	12

#	ARTICLE	IF	CITATIONS
55	Birmingham vasculitis activity and chest manifestation at diagnosis can predict hospitalised infection in ANCA-associated vasculitis. <i>Clinical Rheumatology</i> , 2018, 37, 2133-2141.	2.2	12
56	Brief Report: Atypical Femoral Fracture in Rheumatoid Arthritis Patients Treated With Bisphosphonates: A Nested Caseâ€“Control Study. <i>Arthritis and Rheumatology</i> , 2016, 68, 77-82.	5.6	11
57	Birmingham vasculitis activity score at diagnosis is a significant predictor of relapse of polyarteritis nodosa. <i>Rheumatology International</i> , 2017, 37, 685-694.	3.0	11
58	Fraxinellone Attenuates Rheumatoid Inflammation in Mice. <i>International Journal of Molecular Sciences</i> , 2018, 19, 829.	4.1	11
59	Clinical role of albumin to globulin ratio in microscopic polyangiitis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 487-494.	2.2	11
60	Clinical characteristics and long-term outcomes of Libmanâ€“Sacks endocarditis in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020, 29, 1115-1120.	1.6	11
61	Preemptive antiviral therapy with entecavir can reduce acute deterioration of hepatic function following transarterial chemoembolization. <i>Clinical and Molecular Hepatology</i> , 2016, 22, 458-465.	8.9	11
62	Recommendations for the management of patients with systemic rheumatic diseases during the coronavirus disease pandemic. <i>Korean Journal of Internal Medicine</i> , 2020, 35, 1317-1332.	1.7	11
63	Integrative analysis of lung molecular signatures reveals key drivers of idiopathic pulmonary fibrosis. <i>BMC Pulmonary Medicine</i> , 2021, 21, 404.	2.0	11
64	Chest and renal involvements, Birmingham vascular activity score more than 13.5 and five factor score (1996) more than 1 at diagnosis are significant predictors of relapse of microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 47-54.	0.8	11
65	Serum leucine-rich Î±2-glycoprotein is elevated in patients with systemic lupus erythematosus and correlates with disease activity. <i>Clinica Chimica Acta</i> , 2018, 486, 253-258.	1.1	10
66	Risk of Cancers in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Results from the Korea National Health Insurance Claims Database 2010â€“2018. <i>Journal of Clinical Medicine</i> , 2019, 8, 1871.	2.4	10
67	Comparison of Radiological and Histological Findings of Lung Parenchyma in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019, 60, 454.	2.2	10
68	Echocardiographic features in patients with ANCA-associated vasculitis within 3Â½months before and after diagnosis. <i>Clinical Rheumatology</i> , 2017, 36, 2751-2759.	2.2	9
69	Mean platelet volume can estimate the current vasculitis activity of microscopic polyangiitis. <i>Rheumatology International</i> , 2018, 38, 1095-1101.	3.0	9
70	ANCA positivity at the time of renal biopsy is associated with chronicity index of lupus nephritis. <i>Rheumatology International</i> , 2019, 39, 879-884.	3.0	9
71	Predictors of mortality in autoimmune disease patients with concurrent cytomegalovirus infections detected by quantitative real-time PCR. <i>PLoS ONE</i> , 2017, 12, e0181590.	2.5	9
72	The initial predictors of death in 153 patients with ANCA-associated vasculitis in a single Korean centre. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 65-72.	0.8	9

#	ARTICLE	IF	CITATIONS
73	Evaluation of macrophage activation syndrome in hospitalised patients with Kikuchi-Fujimoto disease based on the 2016 EULAR/ACR/PRINTO classification criteria. <i>PLoS ONE</i> , 2019, 14, e0219970.	2.5	8
74	Controlling Nutritional Status Score is Associated with All-Cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019, 60, 1164.	2.2	8
75	Clinical significance of delta neutrophil index in the differential diagnosis between septic arthritis and acute gout attack within 24 hours after hospitalization. <i>Medicine (United States)</i> , 2017, 96, e7431.	1.0	7
76	Predictor of depressive disorders in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2019, 38, 3485-3491.	2.2	7
77	Deep phenotyping of synovial molecular signatures by integrative systems analysis in rheumatoid arthritis. <i>Rheumatology</i> , 2021, 60, 3420-3431.	1.9	7
78	Hyperuricemia is associated with decreased renal function and occurrence of end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis: a retrospective study. <i>Rheumatology International</i> , 2020, 40, 1089-1099.	3.0	7
79	Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020, 251, 117-123.	1.2	7
80	Generation of Functional Cardiomyocytes from the Synoviocytes of Patients with Rheumatoid Arthritis via Induced Pluripotent Stem Cells. <i>Scientific Reports</i> , 2016, 6, 32669.	3.3	6
81	The clinical utility of splenic fluorodeoxyglucose uptake for diagnosis and prognosis in patients with macrophage activation syndrome. <i>Medicine (United States)</i> , 2017, 96, e7901.	1.0	6
82	Subsequent Thrombotic Outcomes in Patients with Ischemic Stroke with Antiphospholipid Antibody Positivity. <i>Yonsei Medical Journal</i> , 2017, 58, 1128.	2.2	6
83	Red Blood Cell Distribution Width Can Predict Vasculitis Activity and Poor Prognosis in Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2018, 59, 294.	2.2	6
84	Persistent antiphospholipid antibodies are associated with thrombotic events in ANCA-associated vasculitis: A retrospective monocentric study. <i>Nefrologia</i> , 2019, 39, 395-401.	0.4	6
85	Development of synthetic anti-cyclic citrullinated peptide antibody and its arthritogenic role. <i>Clinical and Translational Immunology</i> , 2015, 4, e51.	3.8	5
86	Fever as an initial manifestation of spondyloarthritis: A retrospective study. <i>PLoS ONE</i> , 2017, 12, e0184323.	2.5	5
87	Non-histologic factors discriminating proliferative lupus nephritis from membranous lupus nephritis. <i>Arthritis Research and Therapy</i> , 2020, 22, 138.	3.5	5
88	Comparison of clinical features and outcomes between patients with early and delayed lupus nephritis. <i>BMC Nephrology</i> , 2020, 21, 258.	1.8	5
89	Human adipose-derived mesenchymal stem cells attenuate collagen antibody-induced autoimmune arthritis by inducing expression of FcγRIIb receptors. <i>BMC Musculoskeletal Disorders</i> , 2015, 16, 170.	1.9	4
90	Clinical characteristics of patients with systemic lupus erythematosus showing a false-positive result of syphilis screening. <i>Rheumatology International</i> , 2019, 39, 1859-1866.	3.0	4

#	ARTICLE	IF	CITATIONS
91	Can antineutrophil cytoplasmic antibody positivity at diagnosis predict the poor outcomes of Sjögren's syndrome?. <i>Rheumatology International</i> , 2020, 40, 1063-1070.	3.0	4
92	Characterization of Early-Onset Finger Osteoarthritis-Like Condition Using Patient-Derived Induced Pluripotent Stem Cells. <i>Cells</i> , 2021, 10, 317.	4.1	4
93	Drug Survival of Biologic Therapy in Elderly Patients With Rheumatoid Arthritis Compared With Nonelderly Patients. <i>Journal of Clinical Rheumatology</i> , 2022, 28, e81-e88.	0.9	4
94	Positron Emission Tomography of Chronic Panniculitis Mimicking Extranodal Lymphoma. <i>Arthritis and Rheumatology</i> , 2014, 66, 3250-3250.	5.6	3
95	Could hypereosinophilia at diagnosis estimate the current activity or predict relapse in systemic immunosuppressive drug-naïve patients with eosinophilic granulomatosis with polyangiitis?. <i>Rheumatology International</i> , 2019, 39, 1899-1905.	3.0	3
96	Should nasal biopsy inevitably be performed for classifying granulomatosis with polyangiitis in patients with rhinosinusitis? A retrospective chart review study. <i>Rheumatology International</i> , 2019, 39, 885-892.	3.0	3
97	Clinical characteristics associated with drug-free sustained remission in patients with rheumatoid arthritis: Data from Korean Intensive Management of Early Rheumatoid Arthritis (KIMERA). <i>Seminars in Arthritis and Rheumatism</i> , 2020, 50, 1414-1420.	3.4	3
98	Reclassification of Korean patients with polymyositis and dermatomyositis based on the Bohan and Peter criteria by the 2017 European League Against Rheumatism/American College of Rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 441-446.	1.7	3
99	Cancer development in Korean patients with ANCA-associated vasculitis: a single centre study. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 73-77.	0.8	3
100	Denosumab-Related Osteonecrosis of the Jaw May Not Be a Subject of Teriparatide Treatment. <i>Journal of Bone and Mineral Research</i> , 2020, 37, 2044-2045.	2.8	3
101	Hemoglobin A1c, Not Glycated Albumin, Can Independently Reflect the Ankylosing Spondylitis Disease Activity Score. <i>Journal of Rheumatic Diseases</i> , 2018, 25, 131.	1.1	2
102	Application of 2012 EULAR/ACR criteria for polymyalgia rheumatica to Korean patients previously classified by Chuang and Hunder criteria or Healey criteria. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 1838-1843.	1.9	2
103	Ex Vivo Interferon Gamma Production by Peripheral Immune Cells Predicts Survival in Lung Adenocarcinoma. <i>Clinical Lung Cancer</i> , 2019, 20, e299-e308.	2.6	2
104	Clinical implication of chronic paranasal sinusitis for the classification of microscopic polyangiitis. <i>International Journal of Clinical Practice</i> , 2020, 74, e13431.	1.7	2
105	Double positivity for antineutrophil cytoplasmic antibody (ANCA) and anti-glomerular basement membrane antibody could predict end-stage renal disease in ANCA-associated vasculitis: a monocentric pilot study. <i>Clinical Rheumatology</i> , 2020, 39, 831-840.	2.2	2
106	Clinical and imaging findings suggestive of histopathological immunoglobulin G4-related disease: a single-center retrospective study. <i>Clinical Rheumatology</i> , 2021, 40, 1423-1430.	2.2	2
107	Prevalence of Osteopenia in Drug-Naïve Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Journal of Clinical Rheumatology</i> , 2021, 27, e330-e335.	0.9	2
108	Reclassification of polyarteritis nodosa based on the 1990 ACR criteria using the 2007 EMA algorithm modified by the 2012 CHCC definitions. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 165-166.	0.8	2

#	ARTICLE	IF	CITATIONS
109	Renal outcome of kidney-transplantation in Korean recipients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 115-120.	0.8	2
110	Serum aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1), a novel disease activity predictive biomarker of systemic lupus erythematosus. <i>Clinical and Experimental Rheumatology</i> , 2018, 36, 533-539.	0.8	2
111	Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 117, 116-121.	0.8	2
112	Use of Disease-modifying Antirheumatic Drugs After Cancer Diagnosis in Rheumatoid Arthritis Patients. <i>Journal of Rheumatic Diseases</i> , 2022, 29, 162-170.	1.1	2
113	Clinical Implications of Shared Epitope and Anti-citrullinated Peptide Antibody in Patients With Rheumatoid Arthritis. <i>Journal of Rheumatic Diseases</i> , 2022, 29, 171-180.	1.1	2
114	Evaluation of body composition using computed tomography in patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 1221-1232.	1.7	1
115	Type 4 dual left anterior descending coronary artery. <i>Korean Journal of Internal Medicine</i> , 2015, 30, 727-729.	1.7	1
116	Recommendations for the Management of Patients With Systemic Rheumatic Diseases During the Coronavirus Disease Pandemic. <i>Journal of Rheumatic Diseases</i> , 2020, 27, 218-232.	1.1	1
117	Clinical phenotype with high risk for initiation of biologic therapy in rheumatoid arthritis: a data-driven cluster analysis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 1282-1290.	0.8	1
118	Comparison of developing tuberculosis following tumor necrosis factor inhibition and interleukin-6 inhibition in patients with rheumatoid arthritis: a nationwide observational study in South Korea, 2013â€“2018. <i>Arthritis Research and Therapy</i> , 2022, 24, .	3.5	1
119	Platelet Distribution Width and Mean Platelet Volume Are Not Correlated with the Disease Activity Indices of Ankylosing Spondylitis. <i>Journal of Rheumatic Diseases</i> , 2017, 24, 143.	1.1	0
120	122â€œ...Persistence of anti-smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. , 2019, , .		0
121	123â€œ...Clinical characteristics of patients with systemic lupus erythematosus showing the false positive result of syphilis test. , 2019, , .		0
122	125â€œ...Treatment outcome in lupus nephritis patients treated with mycophenolate mofetil: from a real-world clinical practice. , 2019, , .		0
123	Clinical characteristics and long-term outcomes in patients with mixed Class III/IV + V and pure proliferative lupus nephritis: A single-center experience. <i>Lupus</i> , 2022, 31, 588-595.	1.6	0