

# Seung Min Jung

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8839693/publications.pdf>

Version: 2024-02-01

123  
papers

2,005  
citations

257450

24  
h-index

345221

36  
g-index

125  
all docs

125  
docs citations

125  
times ranked

3527  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Integrative analysis of lung molecular signatures reveals key drivers of systemic sclerosis-associated interstitial lung disease. <i>Annals of the Rheumatic Diseases</i> , 2022, 81, 108-116.   | 0.9 | 16        |
| 2  | Drug Survival of Biologic Therapy in Elderly Patients With Rheumatoid Arthritis Compared With Nonelderly Patients. <i>Journal of Clinical Rheumatology</i> , 2022, 28, e81-e88.  | 0.9 | 4         |
| 3  | Targeted Immunotherapy for Autoimmune Disease. <i>Immune Network</i> , 2022, 22, e9.   | 3.6 | 46        |
| 4  | Clinical characteristics and long-term outcomes in patients with mixed Class III/IV + V and pure proliferative lupus nephritis: A single-center experience. <i>Lupus</i> , 2022, 31, 588-595.  | 1.6 | 0         |
| 5  | Use of Disease-modifying Antirheumatic Drugs After Cancer Diagnosis in Rheumatoid Arthritis Patients. <i>Journal of Rheumatic Diseases</i> , 2022, 29, 162-170.  | 1.1 | 2         |
| 6  | Clinical Implications of Shared Epitope and Anti-citrullinated Peptide Antibody in Patients With Rheumatoid Arthritis. <i>Journal of Rheumatic Diseases</i> , 2022, 29, 171-180.   | 1.1 | 2         |
| 7  | Comparison of developing tuberculosis following tumor necrosis factor inhibition and interleukin-6 inhibition in patients with rheumatoid arthritis: a nationwide observational study in South Korea, 2013-2018. <i>Arthritis Research and Therapy</i> , 2022, 24, .   | 3.5 | 1         |
| 8  | Deep phenotyping of synovial molecular signatures by integrative systems analysis in rheumatoid arthritis. <i>Rheumatology</i> , 2021, 60, 3420-3431.  | 1.9 | 7         |
| 9  | Clinical and imaging findings suggestive of histopathological immunoglobulin G4-related disease: a single-center retrospective study. <i>Clinical Rheumatology</i> , 2021, 40, 1423-1430.  | 2.2 | 2         |
| 10 | Characterization of Early-Onset Finger Osteoarthritis-Like Condition Using Patient-Derived Induced Pluripotent Stem Cells. <i>Cells</i> , 2021, 10, 317.   | 4.1 | 4         |
| 11 | Reclassification of Korean patients with polymyositis and dermatomyositis based on the Bohan and Peter criteria by the 2017 European League Against Rheumatism/American College of Rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 441-446. | 1.7 | 3         |
| 12 | Evaluation of body composition using computed tomography in patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 1221-1232.  | 1.7 | 1         |
| 13 | Prevalence of Osteopenia in Drug-Naive Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Journal of Clinical Rheumatology</i> , 2021, 27, e330-e335.   | 0.9 | 2         |
| 14 | Integrative analysis of lung molecular signatures reveals key drivers of idiopathic pulmonary fibrosis. <i>BMC Pulmonary Medicine</i> , 2021, 21, 404.   | 2.0 | 11        |
| 15 | Clinical phenotype with high risk for initiation of biologic therapy in rheumatoid arthritis: a data-driven cluster analysis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 1282-1290.   | 0.8 | 1         |
| 16 | Can antineutrophil cytoplasmic antibody positivity at diagnosis predict the poor outcomes of Sjögren's syndrome?. <i>Rheumatology International</i> , 2020, 40, 1063-1070.   | 3.0 | 4         |
| 17 | Clinical implication of chronic paranasal sinusitis for the classification of microscopic polyangiitis. <i>International Journal of Clinical Practice</i> , 2020, 74, e13431.  | 1.7 | 2         |
| 18 | Double positivity for antineutrophil cytoplasmic antibody (ANCA) and anti-glomerular basement membrane antibody could predict end-stage renal disease in ANCA-associated vasculitis: a monocentric pilot study. <i>Clinical Rheumatology</i> , 2020, 39, 831-840.  | 2.2 | 2         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Clinical characteristics and long-term outcomes of Libmanâ€Sacks endocarditis in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020, 29, 1115-1120.  | 1.6 | 11        |
| 20 | Non-histologic factors discriminating proliferative lupus nephritis from membranous lupus nephritis. <i>Arthritis Research and Therapy</i> , 2020, 22, 138.   | 3.5 | 5         |
| 21 | Comparison of clinical features and outcomes between patients with early and delayed lupus nephritis. <i>BMC Nephrology</i> , 2020, 21, 258.  | 1.8 | 5         |
| 22 | Hyperuricemia is associated with decreased renal function and occurrence of end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis: a retrospective study. <i>Rheumatology International</i> , 2020, 40, 1089-1099. | 3.0 | 7         |
| 23 | Clinical characteristics associated with drug-free sustained remission in patients with rheumatoid arthritis: Data from Korean Intensive Management of Early Rheumatoid Arthritis (KIMERA). <i>Seminars in Arthritis and Rheumatism</i> , 2020, 50, 1414-1420.      | 3.4 | 3         |
| 24 | Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020, 251, 117-123.                              | 1.2 | 7         |
| 25 | 1,25-dihydroxy Vitamin D3 and Interleukin-6 Blockade Synergistically Regulate Rheumatoid Arthritis by Suppressing Interleukin-17 Production and Osteoclastogenesis. <i>Journal of Korean Medical Science</i> , 2020, 35, e40.                                       | 2.5 | 26        |
| 26 | The use of biological disease-modifying antirheumatic drugs for inflammatory arthritis in Korea: results of a Korean Expert Consensus. <i>Korean Journal of Internal Medicine</i> , 2020, 35, 41-59.  | 1.7 | 20        |
| 27 | Recommendations for the management of patients with systemic rheumatic diseases during the coronavirus disease pandemic. <i>Korean Journal of Internal Medicine</i> , 2020, 35, 1317-1332.  | 1.7 | 11        |
| 28 | The Use of Biological Disease-modifying Antirheumatic Drugs for Inflammatory Arthritis in Korea: Results of a Korean Expert Consensus. <i>Journal of Rheumatic Diseases</i> , 2020, 27, 4.  | 1.1 | 17        |
| 29 | Recommendations for the Management of Patients With Systemic Rheumatic Diseases During the Coronavirus Disease Pandemic. <i>Journal of Rheumatic Diseases</i> , 2020, 27, 218-232.  | 1.1 | 1         |
| 30 | Denosumab-Related Osteonecrosis of the Jaw May Not Be a Subject of Teriparatide Treatment. <i>Journal of Bone and Mineral Research</i> , 2020, 37, 2044-2045.   | 2.8 | 3         |
| 31 | Predictor of depressive disorders in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2019, 38, 3485-3491.   | 2.2 | 7         |
| 32 | Evaluation of macrophage activation syndrome in hospitalised patients with Kikuchi-Fujimoto disease based on the 2016 EULAR/ACR/PRINTO classification criteria. <i>PLoS ONE</i> , 2019, 14, e0219970.   | 2.5 | 8         |
| 33 | Could hypereosinophilia at diagnosis estimate the current activity or predict relapse in systemic immunosuppressive drug-naïve patients with eosinophilic granulomatosis with polyangiitis?. <i>Rheumatology International</i> , 2019, 39, 1899-1905.               | 3.0 | 3         |
| 34 | Comparison of the Clinical Implications among Five Different Nutritional Indices in Patients with Lupus Nephritis. <i>Nutrients</i> , 2019, 11, 1456.   | 4.1 | 14        |
| 35 | Risk of Cancers in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Results from the Korea National Health Insurance Claims Database 2010â€2018. <i>Journal of Clinical Medicine</i> , 2019, 8, 1871.   | 2.4 | 10        |
| 36 | Clinical characteristics of patients with systemic lupus erythematosus showing a false-positive result of syphilis screening. <i>Rheumatology International</i> , 2019, 39, 1859-1866.  | 3.0 | 4         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. <i>Rheumatology International</i> , 2019, 39, 1937-1944.   | 3.0 | 26        |
| 38 | Fibrosis-4 index at diagnosis is associated with all-cause mortality in patients with microscopic polyangiitis and granulomatosis with polyangiitis. <i>BMC Gastroenterology</i> , 2019, 19, 90.  | 2.0 | 13        |
| 39 | Comparison of Radiological and Histological Findings of Lung Parenchyma in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019, 60, 454.  | 2.2 | 10        |
| 40 | Should nasal biopsy inevitably be performed for classifying granulomatosis with polyangiitis in patients with rhinosinusitis? A retrospective chart review study. <i>Rheumatology International</i> , 2019, 39, 885-892.                                  | 3.0 | 3         |
| 41 | Ex Vivo Interferon Gamma Production by Peripheral Immune Cells Predicts Survival in Lung Adenocarcinoma. <i>Clinical Lung Cancer</i> , 2019, 20, e299-e308.   | 2.6 | 2         |
| 42 | Prognostic nutritional index is associated with disease severity and relapse in ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 797-804.   | 1.9 | 16        |
| 43 | ANCA positivity at the time of renal biopsy is associated with chronicity index of lupus nephritis. <i>Rheumatology International</i> , 2019, 39, 879-884.  | 3.0 | 9         |
| 44 | Persistent antiphospholipid antibodies are associated with thrombotic events in ANCA-associated vasculitis: A retrospective monocentric study. <i>Nefrologia</i> , 2019, 39, 395-401.   | 0.4 | 6         |
| 45 | 2018 update of the APLAR recommendations for treatment of rheumatoid arthritis. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 357-375.   | 1.9 | 115       |
| 46 | 122 Persistence of anti-smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. , 2019, , .   |     | 0         |
| 47 | 123 Clinical characteristics of patients with systemic lupus erythematosus showing the false positive result of syphilis test. , 2019, , .  |     | 0         |
| 48 | 125 Treatment outcome in lupus nephritis patients treated with mycophenolate mofetil: from a real-world clinical practice. , 2019, , .  |     | 0         |
| 49 | Clinical role of albumin to globulin ratio in microscopic polyangiitis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 487-494.  | 2.2 | 11        |
| 50 | Low serum complement 3 level is associated with severe ANCA-associated vasculitis at diagnosis. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 223-230.  | 1.6 | 20        |
| 51 | Systemic immune-inflammatory index could estimate the cross-sectional high activity and the poor outcomes in immunosuppressive drug-naïve patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Nephrology</i> , 2019, 24, 711-717. | 1.6 | 42        |
| 52 | No overlap between IgG4-related disease and microscopic polyangiitis and granulomatosis with polyangiitis despite elevated serum IgG4 at diagnosis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 1147-1154.                | 2.2 | 21        |
| 53 | Sodium Chloride Aggravates Arthritis via Th17 Polarization. <i>Yonsei Medical Journal</i> , 2019, 60, 88.   | 2.2 | 26        |
| 54 | Controlling Nutritional Status Score is Associated with All-Cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019, 60, 1164.   | 2.2 | 8         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Risk of malignancy in patients with rheumatoid arthritis after anti-tumor necrosis factor therapy: results from Korean National Health Insurance claims data. <i>Korean Journal of Internal Medicine</i> , 2019, 34, 669-677.   | 1.7 | 13        |
| 56 | Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 117, 116-121. | 0.8 | 2         |
| 57 | Birmingham vasculitis activity and chest manifestation at diagnosis can predict hospitalised infection in ANCA-associated vasculitis. <i>Clinical Rheumatology</i> , 2018, 37, 2133-2141.   | 2.2 | 12        |
| 58 | Mean platelet volume can estimate the current vasculitis activity of microscopic polyangiitis. <i>Rheumatology International</i> , 2018, 38, 1095-1101.   | 3.0 | 9         |
| 59 | Interrupting oral infection of <i>Porphyromonas gingivalis</i> with anti-FimA antibody attenuates bacterial dissemination to the arthritic joint and improves experimental arthritis. <i>Experimental and Molecular Medicine</i> , 2018, 50, e460-e460.                   | 7.7 | 27        |
| 60 | Recent progress of national banking project on homozygous <sc>HLA</sc> $\alpha$ 1 typed induced pluripotent stem cells in <sc>S</sc> outh <sc>K</sc> orea. <i>Journal of Tissue Engineering and Regenerative Medicine</i> , 2018, 12, e1531-e1536.                        | 2.7 | 39        |
| 61 | Decreased muscle mass is independently associated with knee pain in female patients with radiographically mild osteoarthritis: a nationwide cross-sectional study (KNHANES 2010-2011). <i>Clinical Rheumatology</i> , 2018, 37, 1333-1340.                                | 2.2 | 15        |
| 62 | Hemoglobin A1c, Not Glycated Albumin, Can Independently Reflect the Ankylosing Spondylitis Disease Activity Score. <i>Journal of Rheumatic Diseases</i> , 2018, 25, 131.  | 1.1 | 2         |
| 63 | Risk factors associated with inadequate control of disease activity in elderly patients with rheumatoid arthritis: Results from a nationwide Korean College of Rheumatology BIOlogics (KOBIO) registry. <i>PLoS ONE</i> , 2018, 13, e0205651.                             | 2.5 | 13        |
| 64 | Application of 2012 EULAR/ACR criteria for polymyalgia rheumatica to Korean patients previously classified by Chuang and Hunder criteria or Healey criteria. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 1838-1843.                                    | 1.9 | 2         |
| 65 | Red Blood Cell Distribution Width Can Predict Vasculitis Activity and Poor Prognosis in Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2018, 59, 294.  | 2.2 | 6         |
| 66 | Neutrophil to lymphocyte ratio at diagnosis can estimate vasculitis activity and poor prognosis in patients with ANCA-associated vasculitis: a retrospective study. <i>BMC Nephrology</i> , 2018, 19, 187.  | 1.8 | 32        |
| 67 | Safety of Tocilizumab in Rheumatoid Arthritis Patients with Resolved Hepatitis B Virus Infection: Data from Real-World Experience. <i>Yonsei Medical Journal</i> , 2018, 59, 452.   | 2.2 | 27        |
| 68 | Delta Neutrophil Index Is Associated with Vasculitis Activity and Risk of Relapse in ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018, 59, 397.   | 2.2 | 16        |
| 69 | Fraxinellone Attenuates Rheumatoid Inflammation in Mice. <i>International Journal of Molecular Sciences</i> , 2018, 19, 829.  | 4.1 | 11        |
| 70 | Platelet to lymphocyte ratio is associated with the current activity of ANCA-associated vasculitis at diagnosis: a retrospective monocentric study. <i>Rheumatology International</i> , 2018, 38, 1865-1871.  | 3.0 | 28        |
| 71 | Serum leucine-rich $\alpha$ 2-glycoprotein is elevated in patients with systemic lupus erythematosus and correlates with disease activity. <i>Clinica Chimica Acta</i> , 2018, 486, 253-258.  | 1.1 | 10        |
| 72 | Intraperitoneal infusion of mesenchymal stem cell attenuates severity of collagen antibody induced arthritis. <i>PLoS ONE</i> , 2018, 13, e0198740.   | 2.5 | 25        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | Reclassification of polyarteritis nodosa based on the 1990 ACR criteria using the 2007 EMA algorithm modified by the 2012 CHCC definitions. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 165-166.   | 0.8 | 2         |
| 74 | The utility of the ACR/EULAR 2017 provisional classification criteria for granulomatosis with polyangiitis in Korean patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 85-87. | 0.8 | 14        |
| 75 | Renal outcome of kidney-transplantation in Korean recipients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 115-120.   | 0.8 | 2         |
| 76 | Serum aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1), a novel disease activity predictive biomarker of systemic lupus erythematosus. <i>Clinical and Experimental Rheumatology</i> , 2018, 36, 533-539.   | 0.8 | 2         |
| 77 | Cancer development in Korean patients with ANCA-associated vasculitis: a single centre study. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 73-77.   | 0.8 | 3         |
| 78 | The initial predictors of death in 153 patients with ANCA-associated vasculitis in a single Korean centre. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 65-72.  | 0.8 | 9         |
| 79 | Cord blood cell-derived iPSCs as a new candidate for chondrogenic differentiation and cartilage regeneration. <i>Stem Cell Research and Therapy</i> , 2017, 8, 16.   | 5.5 | 59        |
| 80 | Etanercept-Synthesising Mesenchymal Stem Cells Efficiently Ameliorate Collagen-Induced Arthritis. <i>Scientific Reports</i> , 2017, 7, 39593.  | 3.3 | 37        |
| 81 | In-hospital mortality in febrile lupus patients based on 2016 EULAR/ACR/PRINTO classification criteria for macrophage activation syndrome. <i>Seminars in Arthritis and Rheumatism</i> , 2017, 47, 216-221.  | 3.4 | 28        |
| 82 | Evaluation of Spleen Glucose Metabolism Using <sup>18</sup> F-FDG PET/CT in Patients with Febrile Autoimmune Disease. <i>Journal of Nuclear Medicine</i> , 2017, 58, 507-513.  | 5.0 | 33        |
| 83 | Application of the 2016 EULAR/ACR/PRINTO Classification Criteria for Macrophage Activation Syndrome in Patients with Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2017, 44, 996-1003.   | 2.0 | 43        |
| 84 | Birmingham vasculitis activity score at diagnosis is a significant predictor of relapse of polyarteritis nodosa. <i>Rheumatology International</i> , 2017, 37, 685-694.  | 3.0 | 11        |
| 85 | Echocardiographic features in patients with ANCA-associated vasculitis within 3 months before and after diagnosis. <i>Clinical Rheumatology</i> , 2017, 36, 2751-2759.   | 2.2 | 9         |
| 86 | Ethyl pyruvate ameliorates inflammatory arthritis in mice. <i>International Immunopharmacology</i> , 2017, 52, 333-341.  | 3.8 | 19        |
| 87 | The clinical utility of splenic fluorodeoxyglucose uptake for diagnosis and prognosis in patients with macrophage activation syndrome. <i>Medicine (United States)</i> , 2017, 96, e7901.  | 1.0 | 6         |
| 88 | Birmingham vasculitis activity score of more than 9.5 at diagnosis is an independent predictor of refractory disease in granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1593-1605.                                      | 1.9 | 17        |
| 89 | Clinical significance of delta neutrophil index in the differential diagnosis between septic arthritis and acute gout attack within 24 hours after hospitalization. <i>Medicine (United States)</i> , 2017, 96, e7431.   | 1.0 | 7         |
| 90 | Fever as an initial manifestation of spondyloarthritis: A retrospective study. <i>PLoS ONE</i> , 2017, 12, e0184323.   | 2.5 | 5         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 91  | Decreased ex vivo production of interferon-gamma is associated with severity and poor prognosis in patients with lupus. <i>Arthritis Research and Therapy</i> , 2017, 19, 193.  | 3.5 | 14        |
| 92  | Platelet Distribution Width and Mean Platelet Volume Are Not Correlated with the Disease Activity Indices of Ankylosing Spondylitis. <i>Journal of Rheumatic Diseases</i> , 2017, 24, 143.  | 1.1 | 0         |
| 93  | Subsequent Thrombotic Outcomes in Patients with Ischemic Stroke with Antiphospholipid Antibody Positivity. <i>Yonsei Medical Journal</i> , 2017, 58, 1128.  | 2.2 | 6         |
| 94  | Predictors of mortality in autoimmune disease patients with concurrent cytomegalovirus infections detected by quantitative real-time PCR. <i>PLoS ONE</i> , 2017, 12, e0181590.   | 2.5 | 9         |
| 95  | Arthritic role of <i>Porphyromonas gingivalis</i> in collagen-induced arthritis mice. <i>PLoS ONE</i> , 2017, 12, e0188698.   | 2.5 | 39        |
| 96  | Chest and renal involvements, Birmingham vascular activity score more than 13.5 and five factor score (1996) more than 1 at diagnosis are significant predictors of relapse of microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 47-54. | 0.8 | 11        |
| 97  | Clinical and prognostic features of Korean patients with MPO-ANCA, PR3-ANCA and ANCA-negative vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 111-118.  | 0.8 | 16        |
| 98  | A Dual Target-directed Agent against Interleukin-6 Receptor and Tumor Necrosis Factor $\hat{\pm}$ ameliorates experimental arthritis. <i>Scientific Reports</i> , 2016, 6, 20150.   | 3.3 | 26        |
| 99  | Brief Report: Atypical Femoral Fracture in Rheumatoid Arthritis Patients Treated With Bisphosphonates: A Nested Caseâ€“Control Study. <i>Arthritis and Rheumatology</i> , 2016, 68, 77-82.  | 5.6 | 11        |
| 100 | Generation of Functional Cardiomyocytes from the Synoviocytes of Patients with Rheumatoid Arthritis via Induced Pluripotent Stem Cells. <i>Scientific Reports</i> , 2016, 6, 32669.   | 3.3 | 6         |
| 101 | Preemptive antiviral therapy with entecavir can reduce acute deterioration of hepatic function following transarterial chemoembolization. <i>Clinical and Molecular Hepatology</i> , 2016, 22, 458-465.   | 8.9 | 11        |
| 102 | Eupatilin Ameliorates Collagen Induced Arthritis. <i>Journal of Korean Medical Science</i> , 2015, 30, 233.   | 2.5 | 17        |
| 103 | Radiographic Structural Damage Is Worse in the Dominant than the Non-Dominant Hand in Individuals with Early Rheumatoid Arthritis. <i>PLoS ONE</i> , 2015, 10, e0135409.  | 2.5 | 14        |
| 104 | Antinociceptive and anti-inflammatory effects of essential oil extracted from <i>Chamaecyparis obtusa</i> in mice. <i>International Immunopharmacology</i> , 2015, 29, 320-325.   | 3.8 | 15        |
| 105 | Development of synthetic anti-cyclic citrullinated peptide antibody and its arthritogenic role. <i>Clinical and Translational Immunology</i> , 2015, 4, e51.  | 3.8 | 5         |
| 106 | Risk of tuberculosis in patients treated with anti-tumor necrosis factor therapy: a nationwide study in South Korea, a country with an intermediate tuberculosis burden. <i>International Journal of Rheumatic Diseases</i> , 2015, 18, 323-330.                                    | 1.9 | 43        |
| 107 | <sc>APLAR</sc> rheumatoid arthritis treatment recommendations. <i>International Journal of Rheumatic Diseases</i> , 2015, 18, 685-713.  | 1.9 | 109       |
| 108 | Downregulation of Tryptophan-related Metabolomic Profile in Rheumatoid Arthritis Synovial Fluid. <i>Journal of Rheumatology</i> , 2015, 42, 2003-2011.  | 2.0 | 43        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | Human adipose-derived mesenchymal stem cells attenuate collagen antibody-induced autoimmune arthritis by inducing expression of FCγRIIB receptors. <i>BMC Musculoskeletal Disorders</i> , 2015, 16, 170.  | 1.9 | 4         |
| 110 | The Interleukin 33/ST2 Axis in Patients with Primary Sjögren Syndrome: Expression in Serum and Salivary Glands, and the Clinical Association. <i>Journal of Rheumatology</i> , 2015, 42, 264-271.   | 2.0 | 22        |
| 111 | Pulmonary hypertension in systemic lupus erythematosus: an independent predictor of patient survival. <i>Korean Journal of Internal Medicine</i> , 2015, 30, 232.   | 1.7 | 26        |
| 112 | Type 4 dual left anterior descending coronary artery. <i>Korean Journal of Internal Medicine</i> , 2015, 30, 727-729.   | 1.7 | 1         |
| 113 | Recent technological updates and clinical applications of induced pluripotent stem cells. <i>Korean Journal of Internal Medicine</i> , 2014, 29, 547.   | 1.7 | 32        |
| 114 | The Risk of Osteoporotic Fractures According to the FRAX Model in Korean Patients with Rheumatoid Arthritis. <i>Journal of Korean Medical Science</i> , 2014, 29, 1082.   | 2.5 | 15        |
| 115 | Role of Fractalkine in the Pathogenesis of Primary Sjögren Syndrome: Increased Serum Levels of Fractalkine, Its Expression in Labial Salivary Glands, and the Association with Clinical Manifestations. <i>Journal of Rheumatology</i> , 2014, 41, 2425-2438. | 2.0 | 16        |
| 116 | Positron Emission Tomography of Chronic Panniculitis Mimicking Extranodal Lymphoma. <i>Arthritis and Rheumatology</i> , 2014, 66, 3250-3250.  | 5.6 | 3         |
| 117 | Cytokine-Mediated Bone Destruction in Rheumatoid Arthritis. <i>Journal of Immunology Research</i> , 2014, 2014, 1-15.   | 2.2 | 119       |
| 118 | Generation of disease-specific induced pluripotent stem cells from patients with rheumatoid arthritis and osteoarthritis. <i>Arthritis Research and Therapy</i> , 2014, 16, R41.  | 3.5 | 44        |
| 119 | Immunogenicity of anti-tumour necrosis factor therapy in Korean patients with rheumatoid arthritis and ankylosing spondylitis. <i>International Immunopharmacology</i> , 2014, 21, 20-25.   | 3.8 | 18        |
| 120 | Induced production of anti-etanercept antibody in collagen-induced arthritis. <i>Molecular Medicine Reports</i> , 2014, 9, 2301-2308.   | 2.4 | 20        |
| 121 | A New Strategy to Deliver Synthetic Protein Drugs: Self-reproducible Biologics Using Minicircles. <i>Scientific Reports</i> , 2014, 4, 5961.  | 3.3 | 19        |
| 122 | Self in vivo production of a synthetic biological drug CTLA4Ig using a minicircle vector. <i>Scientific Reports</i> , 2014, 4, 6935.  | 3.3 | 12        |
| 123 | Role of intrahepatic tumor control in the prognosis of patients with hepatocellular carcinoma and extrahepatic metastases. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2012, 27, 684-689.   | 2.8 | 61        |