

Yiangos Yiangou

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8865172/publications.pdf>

Version: 2024-02-01

32
papers

3,106
citations

304743

22
h-index

414414

32
g-index

32
all docs

32
docs citations

32
times ranked

3533
citing authors

#	ARTICLE	IF	CITATIONS
1	Disruption of the P2X7 purinoceptor gene abolishes chronic inflammatory and neuropathic pain. <i>Pain</i> , 2005, 114, 386-396.	4.2	687
2	COX-2, CB2 and P2X7-immunoreactivities are increased in activated microglial cells/macrophages of multiple sclerosis and amyotrophic lateral sclerosis spinal cord. <i>BMC Neurology</i> , 2006, 6, 12.	1.8	420
3	Capsaicin receptor TRPV1 in urothelium of neurogenic human bladders and effect of intravesical resiniferatoxin. <i>Urology</i> , 2005, 65, 400-405.	1.0	186
4	Expression of the TRPV1 receptor differs in quiescent inflammatory bowel disease with or without abdominal pain. <i>Gut</i> , 2010, 59, 767-774.	12.1	184
5	Localization of M ₂ and M ₃ Muscarinic Receptors in Human Bladder Disorders and Their Clinical Correlations. <i>Journal of Urology</i> , 2006, 176, 367-373.	0.4	163
6	Cool and menthol receptor TRPM8 in human urinary bladder disorders and clinical correlations. <i>BMC Urology</i> , 2006, 6, 6.	1.4	160
7	Increased vanilloid receptor VR1 innervation in vulvodynia. <i>European Journal of Pain</i> , 2004, 8, 129-133.	2.8	159
8	P2X3-Immunoreactive Nerve Fibres in Neurogenic Detrusor Overactivity and the Effect of Intravesical Resiniferatoxin*1. <i>European Urology</i> , 2004, 46, 247-253.	1.9	158
9	Cannabinoid receptor CB2 localisation and agonist-mediated inhibition of capsaicin responses in human sensory neurons. <i>Pain</i> , 2008, 138, 667-680.	4.2	140
10	Increased acid-sensing ion channel ASIC-3 in inflamed human intestine. <i>European Journal of Gastroenterology and Hepatology</i> , 2001, 13, 891-896.	1.6	106
11	Prostanoid receptor EP1 and Cox-2 in injured human nerves and a rat model of nerve injury: a time-course study. <i>BMC Neurology</i> , 2006, 6, 1.	1.8	74
12	Transient Receptor Potential Vanilloid Receptor Subtype 1 in Painful Bladder Syndrome and its Correlation With Pain. <i>Journal of Urology</i> , 2006, 176, 797-801.	0.4	60
13	<p>Rational treatment of chemotherapy-induced peripheral neuropathy with capsaicin 8% patch: from pain relief towards disease modification</p>. <i>Journal of Pain Research</i> , 2019, Volume 12, 2039-2052.	2.0	58
14	Expression and Function of Junctional Adhesion Molecule-C in Myelinated Peripheral Nerves. <i>Science</i> , 2007, 318, 1472-1475.	12.6	55
15	Increased Cannabinoid Receptor 1-Immunoreactive Nerve Fibers in Overactive and Painful Bladder Disorders and Their Correlation With Symptoms. <i>Urology</i> , 2010, 75, 1514.e15-1514.e20.	1.0	55
16	Mechanisms Underlying Clinical Efficacy of Angiotensin II Type 2 Receptor (AT ₂ R) Antagonist EMA401 in Neuropathic Pain: Clinical Tissue and in Vitro Studies. <i>Molecular Pain</i> , 2015, 11, s12990-015-0038.	2.1	53
17	Sensory purinergic receptor P2X3 is elevated in burning mouth syndrome. <i>International Journal of Oral and Maxillofacial Surgery</i> , 2010, 39, 815-819.	1.5	50
18	Capsaicin receptor VR1 and ATP purinoceptor P2X3 in painful and nonpainful human tooth pulp. <i>Journal of Orofacial Pain</i> , 2003, 17, 245-50.	1.7	48

#	ARTICLE	IF	CITATIONS
19	Molecular forms of NGF in human and rat neuropathic tissues: decreased NGF precursor-like immunoreactivity in human diabetic skin. <i>Journal of the Peripheral Nervous System</i> , 2002, 7, 190-197.	3.1	40
20	Voltage-gated ion channel Nav1.7 innervation in patients with idiopathic rectal hypersensitivity and paroxysmal extreme pain disorder (familial rectal pain). <i>Neuroscience Letters</i> , 2007, 427, 77-82.	2.1	39
21	Sodium channel Nav1.7 immunoreactivity in painful human dental pulp and burning mouth syndrome. <i>BMC Neuroscience</i> , 2010, 11, 71.	1.9	28
22	Granulocyte-macrophage colony-stimulating factor receptor expression in clinical pain disorder tissues and role in neuronal sensitization. <i>Pain Reports</i> , 2018, 3, e676.	2.7	28
23	Increased levels of SV2A botulinum neurotoxin receptor in clinical sensory disorders and functional effects of botulinum toxins A and E in cultured human sensory neurons. <i>Journal of Pain Research</i> , 2011, 4, 347.	2.0	25
24	Characterization of a novel prepro VIP derived peptide. <i>Biochemical and Biophysical Research Communications</i> , 1986, 139, 1142-1149.	2.1	22
25	Trench Foot or Non-Freezing Cold Injury As a Painful Vaso-Neuropathy: Clinical and Skin Biopsy Assessments. <i>Frontiers in Neurology</i> , 2017, 8, 514.	2.4	22
26	Plasticity of gene expression in injured human dorsal root ganglia revealed by GeneChip oligonucleotide microarrays. <i>Journal of Clinical Neuroscience</i> , 2004, 11, 289-299.	1.5	19
27	Nociceptin/orphanin FQ receptor expression in clinical pain disorders and functional effects in cultured neurons. <i>Pain</i> , 2016, 157, 1960-1969.	4.2	19
28	Glucagon-like peptide 1 receptor (GLP-1R) expression by nerve fibres in inflammatory bowel disease and functional effects in cultured neurons. <i>PLoS ONE</i> , 2018, 13, e0198024.	2.5	18
29	Infusion of prepro-VIP Derived Peptides in Man: Effect on Secretion of Prolactin. <i>Neuroendocrinology</i> , 1988, 48, 615-618.	2.5	15
30	Nerve and Vascular Biomarkers in Skin Biopsies Differentiate Painful From Painless Peripheral Neuropathy in Type 2 Diabetes. <i>Frontiers in Pain Research</i> , 2021, 2, 731658.	2.0	6
31	Effects of preprovasoactive intestinal polypeptide-derived peptides on ileal output. <i>Gastroenterology</i> , 1990, 98, 505-508.	1.3	5
32	Comparison of neuromedin-N and neurotensin on net fluid flux across rat small intestine. <i>European Journal of Pharmacology</i> , 1990, 175, 43-47.	3.5	4