Cong Xue

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8884640/publications.pdf

Version: 2024-02-01

567281 330143 1,434 40 15 37 citations h-index g-index papers 40 40 40 2926 times ranked docs citations citing authors all docs

#	Article	IF	CITATIONS
1	EBV-driven LMP1 and IFN- \hat{I}^3 up-regulate PD-L1 in nasopharyngeal carcinoma: Implications for oncotargeted therapy. Oncotarget, 2014, 5, 12189-12202.	1.8	324
2	High PD-L1 expression was associated with poor prognosis in 870 Chinese patients with breast cancer. Oncotarget, 2015, 6, 33972-33981.	1.8	159
3	High infiltration of tumor-associated macrophages in triple-negative breast cancer is associated with a higher risk of distant metastasis. OncoTargets and Therapy, 2014, 7, 1475.	2.0	149
4	Network Meta-Analysis of Erlotinib, Gefitinib, Afatinib and Icotinib in Patients with Advanced Non-Small-Cell Lung Cancer Harboring EGFR Mutations. PLoS ONE, 2014, 9, e85245.	2.5	125
5	Effect of Capecitabine Maintenance Therapy Using Lower Dosage and Higher Frequency vs Observation on Disease-Free Survival Among Patients With Early-Stage Triple-Negative Breast Cancer Who Had Received Standard Treatment. JAMA - Journal of the American Medical Association, 2021, 325, 50.	7.4	113
6	National survey of the medical treatment status for non-small cell lung cancer (NSCLC) in China. Lung Cancer, 2012, 77, 371-375.	2.0	112
7	PD-L1 is remarkably over-expressed in EBV-associated pulmonary lymphoepithelioma-like carcinoma and related to poor disease-free survival. Oncotarget, 2015, 6, 33019-33032.	1.8	69
8	Effects of an oral allosteric AKT inhibitor (MK-2206) on human nasopharyngeal cancer in vitro and in vivo. Drug Design, Development and Therapy, 2014, 8, 1827.	4.3	40
9	Optimized selection of three major EGFR-TKIs in advanced EGFR-positive non-small cell lung cancer: a network metaanalysis. Oncotarget, 2016, 7, 20093-20108.	1.8	31
10	A Large-scale Cross-sectional Study of ALK Rearrangements and EGFR Mutations in Non-small-cell Lung Cancer in Chinese Han Population. Scientific Reports, 2014, 4, 7268.	3.3	28
11	Trastuzumab Plus Endocrine Therapy or Chemotherapy as First-line Treatment for Patients with Hormone Receptor–Positive and HER2-Positive Metastatic Breast Cancer (SYSUCC-002). Clinical Cancer Research, 2022, 28, 637-645.	7.0	27
12	Efficacy of the hypoxiaâ€activated prodrug evofosfamide (THâ€302) in nasopharyngeal carcinoma in vitro and in vivo. Cancer Communications, 2018, 38, 1-9.	9.2	23
13	Adjuvant chemotherapy for small, lymph node–negative, tripleâ€negative breast cancer: A singleâ€center study and a metaâ€analysis of the published literature. Cancer, 2020, 126, 3837-3846.	4.1	20
14	The Effects of Ganglioside-Monosialic Acid in Taxane-Induced Peripheral Neurotoxicity in Patients with Breast Cancer: A Randomized Trial. Journal of the National Cancer Institute, 2020, 112, 55-62.	6.3	19
15	A multicenter, retrospective epidemiologic survey of the clinical features and management of bone metastatic disease in China. Chinese Journal of Cancer, 2016, 35, 40.	4.9	18
16	Olanzapine-Based Triple Regimens Versus Neurokinin-1 Receptor Antagonist-Based Triple Regimens in Preventing Chemotherapy-Induced Nausea and Vomiting Associated with Highly Emetogenic Chemotherapy: A Network Meta-Analysis. Oncologist, 2018, 23, 603-616.	3.7	17
17	A Novel Ferroptosis-Related Gene Signature Predicts Overall Survival of Breast Cancer Patients. Biology, 2021, 10, 151.	2.8	17
18	Risk of treatment-related deaths with vascular endothelial growth factor receptor tyrosine kinase inhibitors: a meta-analysis of 41 randomized controlled trials. OncoTargets and Therapy, 2014, 7, 1851.	2.0	15

#	Article	IF	CITATIONS
19	Safety, tolerability, and pharmacokinetics of BAT8001 in patients with HER2â€positive breast cancer: An openâ€label, doseâ€escalation, phase I study. Cancer Communications, 2021, 41, 171-182.	9.2	15
20	Tumor-infiltrating lymphocytes predict prognosis of breast cancer patients treated with anti-Her-2 therapy. Oncotarget, 2017, 8, 5219-5232.	1.8	15
21	In vitro and in vivo efficacy of afatinib as a single agent or in combination with gemcitabine for the treatment of nasopharyngeal carcinoma. Drug Design, Development and Therapy, 2016, 10, 1299.	4.3	13
22	An investigation of symptom burden and quality of life in Chinese chemo-naÃ-ve advanced lung cancer patients by using the Instrument-Cloud QOL System. Lung Cancer, 2014, 84, 301-306.	2.0	12
23	Multiple oncogenic mutations related to targeted therapy in nasopharyngeal carcinoma. Chinese Journal of Cancer, 2015, 34, 177-83.	4.9	10
24	Trastuzumab plus endocrine therapy or chemotherapy as first-line treatment for metastatic breast cancer with hormone receptor-positive and HER2-positive: The sysucc-002 randomized clinical trial Journal of Clinical Oncology, 2021, 39, 1003-1003.	1.6	10
25	QoL analyses from INFORM study, a phase III study of gefitinib versus placebo as maintenance therapy in advanced NSCLC. Scientific Reports, 2015, 5, 11934.	3.3	9
26	Pretreatment anti-Mullerian hormone-based nomogram predicts menstruation status after chemotherapy for premenopausal women with hormone receptor-positive early breast cancer. Breast Cancer Research and Treatment, 2019, 173, 619-628.	2.5	9
27	Randomized, Multicenter Study of Gefitinib Dose-escalation in Advanced Non-small-cell Lung Cancer Patients Achieved Stable Disease after One-month Gefitinib Treatment. Scientific Reports, 2015, 5, 10648.	3.3	7
28	Treatment Outcome of Different Chemotherapy in Patients With Relapsed or Metastatic Malignant Urachal Tumor. Frontiers in Oncology, 2021, 11, 739134.	2.8	5
29	Clinical analysis of 50 Eastern Asian patients with primary pulmonary large-cell neuroendocrine carcinoma. OncoTargets and Therapy, 2015, 8, 1219.	2.0	4
30	Met tyrosine kinase inhibitor, PF-2341066, suppresses growth and invasion of nasopharyngeal carcinoma. Drug Design, Development and Therapy, 2015, 9, 4897.	4.3	4
31	Optimal first-line treatment for platinum-eligible metastatic urothelial carcinoma: Comparison of chemo-immunotherapy, immunotherapy, and chemotherapy— A systematic review and meta-analysis. Clinical Immunology, 2022, 236, 108927.	3.2	4
32	Effectiveness of capecitabine with or without docetaxel therapy for the treatment of patients with advanced urothelial carcinoma: a single-institution experience. Oncotarget, 2016, 7, 63722-63729.	1.8	2
33	Ovarian Function, Not Age, Predicts the Benefit from Ovarian Suppression or Ablation for Premenopausal Women with Breast Cancer. PLoS ONE, 2016, 11, e0148849.	2.5	2
34	Immune checkpoint inhibitor plus anti-EGFR target therapy plus chemotherapy in patients with locally advanced penile squamous cell carcinoma Journal of Clinical Oncology, 2021, 39, e17016-e17016.	1.6	2
35	Role of a Modified Urothelium Immune Prognostic Index in Patients With Metastatic Urothelial Carcinoma Treated With Anti–PD-1/PD-L1–Based Therapy. Frontiers in Molecular Biosciences, 2021, 8, 621883.	3.5	2
36	Operable Breast Cancer of the Inner Hemisphere Is Associated with Poor Survival. Journal of Breast Cancer, 2015, 18, 36.	1.9	1

#	Article	IF	CITATIONS
37	Efficacy of BIBF 1120 or BIBF 1120 plus chemotherapy on nasopharyngeal carcinoma in vitro and in vivo. Drug Design, Development and Therapy, 2016, 10, 1173.	4.3	1
38	Predict the benefit of metronomic capecitabine maintenance in early-stage triple-negative breast cancer: Results from the SYSUCC-001 study Journal of Clinical Oncology, 2021, 39, 521-521.	1.6	1
39	Endocrine treatment-related symptoms and outcomes in breast cancer: A systematic review and meta-analysis Journal of Clinical Oncology, 2016, 34, e12001-e12001.	1.6	O
40	Two-year follow-up of pretreatment anti-mullerian hormone (AMH) to predict for resumption of mense after chemotherapy in premenopausal hormone-receptor (HR+) positive early breast cancer (eBC) Journal of Clinical Oncology, 2018, 36, e12511-e12511.	1.6	0