
Carmen Quero

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8903989/publications.pdf

Version: 2024-02-01



#	Article	IF	CITATIONS
1	Inhibitory effect of thymol on pheromone-mediated attraction in two pest moth species. Scientific Reports, 2021, 11, 1223.	3.3	5
2	Development of Novel Management Tools for Phortica variegata (Diptera: Drosophilidae), Vector of the Oriental Eyeworm, Thelazia callipaeda (Spirurida: Thelaziidae), in Europe. Journal of Medical Entomology, 2021, , .	1.8	3
3	Olean (1,7-dioxaspiro[5.5]undecane): A Novel Intraspecific Chemical Cue in Coraebus undatus (F.) (Coleoptera: Buprestidae). Insects, 2021, 12, 1085.	2.2	0
4	Two chiral types of randomly rotated ommatidia are distributed across the retina of the flathead oak borer, <i>Coraebus undatus</i> (Coleoptera: Buprestidae). Journal of Experimental Biology, 2020, 223, .	1.7	8
5	A proteomic analysis of the statocyst endolymph in common cuttlefish (Sepia officinalis): an assessment of acoustic trauma after exposure to sound. Scientific Reports, 2019, 9, 9340.	3.3	12
6	Influence of Age, Host Plant and Mating Status in Pheromone Production and New Insights on Perception Plasticity in Tuta Absoluta. Insects, 2019, 10, 256.	2.2	9
7	Enantioselective Synthesis and Activity of All Diastereoisomers of (<i>E</i>)-Phytal, a Pheromone Component of the Moroccan Locust, <i>Dociostaurus maroccanus</i> . Journal of Agricultural and Food Chemistry, 2019, 67, 72-80.	5.2	8
8	Short-term peripheral sensitization by brief exposure to pheromone components in Spodoptera littoralis. Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology, 2017, 203, 973-982.	1.6	6
9	Sexual communication in castniid moths: Males mark their territories and appear to bear all chemical burden. PLoS ONE, 2017, 12, e0171166.	2.5	7
10	Sexual communication in day-flying Lepidoptera with special reference to castniids or â€`butterfly-moths'. Bulletin of Entomological Research, 2016, 106, 421-431.	1.0	19
11	Synthesis, Functional Assays, Electrophysiological Activity, and Field Tests of Pheromone Antagonists of the Tomato Leafminer, Tuta absoluta. Journal of Agricultural and Food Chemistry, 2016, 64, 3523-3532.	5.2	10
12	Using a polymer probe characterized by MALDI-TOF/MS to assess river ecosystem functioning: From polymer selection to field tests. Science of the Total Environment, 2016, 573, 532-540.	8.0	6
13	MALDI-TOF MS Imaging evidences spatial differences in the degradation of solid polycaprolactone diol in water under aerobic and denitrifying conditions. Science of the Total Environment, 2016, 566-567, 27-33.	8.0	41
14	Field trapping of the flathead oak borer <i>Coroebus undatus</i> (Coleoptera: Buprestidae) with different traps and volatile lures. Insect Science, 2015, 22, 139-149.	3.0	13
15	Aggressive mimicry coexists with mutualism in an aphid. Proceedings of the National Academy of Sciences of the United States of America, 2015, 112, 1101-1106.	7.1	28
16	EAG Responses Increase of <i>Spodoptera littoralis</i> Antennae after a Single Pheromone Pulse. Natural Product Communications, 2014, 9, 1934578X1400900.	0.5	5
17	Semiochemical and natural product-based approaches to control Spodoptera spp. (Lepidoptera:) Tj ETQq1 1 0.	784314 rgBT 3.7	Overlock 29
18	EAG responses increase of Spodoptera littoralis antennae after a single pheromone pulse. Natural Product Communications, 2014, 9, 1099-101.	0.5	3

CARMEN QUERO

#	Article	IF	CITATIONS
19	Electrophysiological and behavioural responses of <i>Pityophthorus pubescens</i> (Coleoptera:) Tj ETQq1 1 0.784	1314 rgBT 3.4	/Overlock 1 9
19	(<i>S</i>)â€(â^')â€verbenone in <i>Pinus radiata</i> (Pinaceae) stands in northern Spain. Pest Management Science, 2013, 69, 40-47.	0.4	9
20	Phytal: A Candidate Sex Pheromone Component of the Moroccan Locust <i>Dociostaurus maroccanus</i> . ChemBioChem, 2013, 14, 1450-1459.	2.6	4
21	Electrophysiological and Behavioral Responses of the Black-Banded Oak Borer, Coroebus florentinus, to Conspecific and Host-Plant Volatiles. Journal of Chemical Ecology, 2012, 38, 378-388.	1.8	30
22	New identification of proanthocyanidins in cinnamon (Cinnamomum zeylanicum L.) using MALDI-TOF/TOF mass spectrometry. Analytical and Bioanalytical Chemistry, 2012, 402, 1327-1336.	3.7	51
23	Moths Behaving like Butterflies. Evolutionary Loss of Long Range Attractant Pheromones in Castniid Moths: A Paysandisia archon Model. PLoS ONE, 2012, 7, e29282.	2.5	33
24	Lowâ€frequency sounds induce acoustic trauma in cephalopods. Frontiers in Ecology and the Environment, 2011, 9, 489-493.	4.0	110
25	Proteomics of toxic oil syndrome in humans: Phenotype distribution in a population of patients. Chemico-Biological Interactions, 2011, 192, 129-135.	4.0	4
26	Evidence for (E)-pityol as an aggregation pheromone of Pityophthorus pubescens (Coleoptera:) Tj ETQq0 0 0 rgBT	/Oyerlock	18 Tf 50 46
27	Sex Pheromone of the Spanish Population of the Beet Armyworm Spodoptera exigua. Journal of Chemical Ecology, 2010, 36, 778-786.	1.8	26
28	Improved resolution in the acidic and basic region of 2â€ÐE of insect antennae proteins using hydroxyethyl disulfide. Electrophoresis, 2009, 30, 2613-2616.	2.4	2
29	Expression of differential antennal proteins in males and females of an important crop pest, Sesamia nonagrioides. Insect Biochemistry and Molecular Biology, 2009, 39, 11-19.	2.7	18
30	Biosynthetic pathways of the pheromone of the Egyptian armyworm <i>Spodoptera littoralis</i> . Physiological Entomology, 2008, 33, 275-290.	1.5	30
31	Biorational insecticides in pest management. Journal of Pesticide Sciences, 2008, 33, 103-121.	1.4	178
32	Inheritance of Olfactory Preferences II. Olfactory Receptor Neuron Responses from <i>Heliothis subflexa</i> × <i>Heliothis virescens</i> Hybrid Male Moths. Brain, Behavior and Evolution, 2006, 68, 75-89.	1.7	22
33	Proteotyping of human haptoglobin by MALDI-TOF profiling: Phenotype distribution in a population of toxic oil syndrome patients. Proteomics, 2006, 6, S272-S281.	2.2	9
34	Electrophysiological and Behavioral Responses of a Cuban Population of the Sweet Potato Weevil to its Sex Pheromone. Journal of Chemical Ecology, 2006, 32, 2177-2190.	1.8	6
35	A comparison of responses from olfactory receptor neurons of Heliothis subflexa and Heliothis virescens to components of their sex pheromone. Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology, 2004, 190, 155-165.	1.6	109
36	Responses of the olfactory receptor neurons of the corn stalk borerSesamia nonagrioides to components of the pheromone blend and their inhibition by a trifluoromethyl ketone analogue of the	3.4	19

components of the pheromone blend and their inhibition by a trifluoromethyl ketone analogue of the main component. Pest Management Science, 2004, 60, 719-726. 36 3.4

CARMEN QUERO

#	Article	IF	CITATIONS
37	Determination of protein markers in human serum: Analysis of protein expression in toxic oil syndrome studies. Proteomics, 2004, 4, 303-315.	2.2	44
38	New fluorinated derivatives as esterase inhibitors. Synthesis, hydration and crossed specificity studies. Bioorganic and Medicinal Chemistry, 2003, 11, 1047-1055.	3.0	26
39	Sex Pheromone of the Oak Processionary MothThaumetopoeaprocessionea. Identification and Biological Activity. Journal of Agricultural and Food Chemistry, 2003, 51, 2987-2991.	5.2	14
40	Activity of Octylthiotrifluoropropan-2-one, a Potent Esterase Inhibitor, on Growth, Development, and Intraspecific Communication in Spodoptera littoralis and Sesamia nonagrioides. Journal of Agricultural and Food Chemistry, 2002, 50, 7062-7068.	5.2	6
41	Responses of maleHelicoverpa zeato single pulses of sex pheromone and behavioural antagonist. Physiological Entomology, 2001, 26, 106-115.	1.5	27
42	Control of the biosynthetic pathway of Sesamia nonagrioides sex pheromone by the pheromone biosynthesis activating neuropeptide. Insect Biochemistry and Molecular Biology, 2000, 30, 455-459.	2.7	20
43	Title is missing!. Journal of Insect Behavior, 1999, 12, 701-710.	0.7	32
44	Effects of Trifluoromethyl Ketones and Related Compounds on the EAG and Behavioural Responses to Pheromones in Male Moths. Chemical Senses, 1997, 22, 407-416.	2.0	32
45	Reinvestigation of Female Sex Pheromone of Processionary Moth (Thaumetopoea pityocampa): No Evidence for Minor Components. Journal of Chemical Ecology, 1997, 23, 713-726.	1.8	14
46	Behavioral responses ofSpodoptera littoralis males to sex pheromone components and virgin females in wind tunnel. Journal of Chemical Ecology, 1996, 22, 1087-1102.	1.8	34
47	Behavior of processionary males (Thaumetopoea pityocampa) induced by sex pheromone and analogs in a wind tunnel. Journal of Chemical Ecology, 1995, 21, 1957-1969.	1.8	21
48	Inhibition of pheromone action inSesamia nonagrioidesby Haloacetate analogues. Pest Management Science, 1994, 41, 97-103.	0.4	14