

Alisha Holtzhausen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8906582/publications.pdf>

Version: 2024-02-01

21
papers

1,161
citations

687363

13
h-index

940533

16
g-index

21
all docs

21
docs citations

21
times ranked

2072
citing authors

#	ARTICLE	IF	CITATIONS
1	Pharmacological Wnt ligand inhibition overcomes key tumor-mediated resistance pathways to anti-PD-1 immunotherapy. <i>Cell Reports</i> , 2021, 35, 109071.	6.4	35
2	Overcoming Immunotherapy Resistance by Targeting the Tumor-Intrinsic NLRP3-HSP70 Signaling Axis. <i>Cancers</i> , 2021, 13, 4753.	3.7	9
3	A tumor-intrinsic PD-L1/NLRP3 inflammasome signaling pathway drives resistance to anti-PD-1 immunotherapy. <i>Journal of Clinical Investigation</i> , 2020, 130, 2570-2586.	8.2	134
4	TAM Family Receptor Kinase Inhibition Reverses MDSC-Mediated Suppression and Augments Anti-PD-1 Therapy in Melanoma. <i>Cancer Immunology Research</i> , 2019, 7, 1672-1686.	3.4	85
5	MERTK mediated novel site Akt phosphorylation alleviates SAV1 suppression. <i>Nature Communications</i> , 2019, 10, 1515.	12.8	25
6	Paracrine Wnt5a- β -Catenin Signaling Triggers a Metabolic Program that Drives Dendritic Cell Tolerization. <i>Immunity</i> , 2018, 48, 147-160.e7.	14.3	185
7	Stromal Fibroblasts Mediate Anti-PD-1 Resistance via MMP-9 and Dictate TGF- β Inhibitor Sequencing in Melanoma. <i>Cancer Immunology Research</i> , 2018, 6, 1459-1471.	3.4	81
8	Factor XIIIa-expressing inflammatory monocytes promote lung squamous cancer through fibrin cross-linking. <i>Nature Communications</i> , 2018, 9, 1988.	12.8	69
9	Tumor-secreted Pros1 inhibits macrophage M1 polarization to reduce antitumor immune response. <i>Journal of Clinical Investigation</i> , 2018, 128, 2356-2369.	8.2	118
10	Melanoma-Derived Wnt5a Promotes Local Dendritic-Cell Expression of IDO and Immunotolerance: Opportunities for Pharmacologic Enhancement of Immunotherapy. <i>Cancer Immunology Research</i> , 2015, 3, 1082-1095.	3.4	147
11	Targeting the Wnt5a- β -catenin pathway in the melanoma microenvironment to augment checkpoint inhibitor immunotherapy. <i>Journal of Clinical Oncology</i> , 2015, 33, 3054-3054.	1.6	2
12	Early Carcinogenesis Involves the Establishment of Immune Privilege via Intrinsic and Extrinsic Regulation of Indoleamine 2,3-dioxygenase-1: Translational Implications in Cancer Immunotherapy. <i>Frontiers in Immunology</i> , 2014, 5, 438.	4.8	12
13	Novel bone morphogenetic protein signaling through Smad2 and Smad3 to regulate cancer progression and development. <i>FASEB Journal</i> , 2014, 28, 1248-1267.	0.5	80
14	Combinatorial TGF- β signaling blockade and anti-CTLA-4 antibody immunotherapy in a murine BRAF ^{V600E} -PTEN ^{-/-} transgenic model of melanoma. <i>Journal of Clinical Oncology</i> , 2014, 32, 3011-3011.	1.6	25
15	Role of the Wnt- β -catenin signaling pathway in melanoma-mediated dendritic cell tolerization. , 2013, 1, P153.		3
16	Type III TGF- β receptor downregulation generates an immunotolerant tumor microenvironment. <i>Journal of Clinical Investigation</i> , 2013, 123, 3925-3940.	8.2	94
17	Abstract 3548: Loss of the type III TGF- β receptor during cancer progression generates an immunotolerant tumor microenvironment: Translational implications for TGF- β inhibition and immunotherapy biomarker development. , 2012, , .		0
18	Abstract 3035: Bone morphogenetic proteins signal through Smad2 and Smad3 to regulate cell migration and proliferation. , 2012, , .		0

#	ARTICLE	IF	CITATIONS
19	Type III TGF- β 2 Receptor Enhances Colon Cancer Cell Migration and Anchorage-Independent Growth. <i>Neoplasia</i> , 2011, 13, 758-IN28.	5.3	56
20	Abstract 3972: The role of the TGF- β 2 type III receptor in colon carcinogenesis. , 2010, , .		0
21	Identification of a Germline Pylori Variant in a Metastatic Melanoma Patient With Multiple Spontaneous Regressions and Immune-related Adverse Events. <i>Journal of Immunotherapy</i> , 0, Publish Ahead of Print, .	2.4	1