

# Maryam Salimi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8913270/publications.pdf>

Version: 2024-02-01

17  
papers

2,851  
citations

567281

15  
h-index

839539

18  
g-index

18  
all docs

18  
docs citations

18  
times ranked

4204  
citing authors

#	ARTICLE	IF	CITATIONS
1	IL-6 effector function of group 2 innate lymphoid cells (ILC2) is NOD2 dependent. <i>Science Immunology</i> , 2021, 6, .	11.9	8
2	Re-evaluation of human BDCA-2+ DC during acute sterile skin inflammation. <i>Journal of Experimental Medicine</i> , 2020, 217, .	8.5	29
3	Activated innate lymphoid cell populations accumulate in human tumour tissues. <i>BMC Cancer</i> , 2018, 18, 341.	2.6	85
4	Synergistic activation of pro-inflammatory type-2 CD8+ T lymphocytes by lipid mediators in severe eosinophilic asthma. <i>Mucosal Immunology</i> , 2018, 11, 1408-1419.	6.0	46
5	Cysteinyl leukotriene E 4 activates human group 2 innate lymphoid cells and enhances the effect of prostaglandin D 2 and epithelial cytokines. <i>Journal of Allergy and Clinical Immunology</i> , 2017, 140, 1090-1100.e11.	2.9	130
6	CD1a presentation of endogenous antigens by group 2 innate lymphoid cells. <i>Science Immunology</i> , 2017, 2, .	11.9	57
7	Engineering chimeric human and mouse major histocompatibility complex (MHC) class I tetramers for the production of T-cell receptor (TCR) mimic antibodies. <i>PLoS ONE</i> , 2017, 12, e0176642.	2.5	7
8	Psoriatic T cells recognize neolipid antigens generated by mast cell phospholipase delivered by exosomes and presented by CD1a. <i>Journal of Experimental Medicine</i> , 2016, 213, 2399-2412.	8.5	194
9	Filaggrin inhibits generation of CD1a neolipid antigens by house dust mite-derived phospholipase. <i>Science Translational Medicine</i> , 2016, 8, 325ra18.	12.4	77
10	Group 2 Innate Lymphoid Cells Express Functional NKp30 Receptor Inducing Type 2 Cytokine Production. <i>Journal of Immunology</i> , 2016, 196, 45-54.	0.8	73
11	Spontaneous atopic dermatitis is mediated by innate immunity, with the secondary lung inflammation of the atopic march requiring adaptive immunity. <i>Journal of Allergy and Clinical Immunology</i> , 2016, 137, 482-491.	2.9	117
12	Bee venom processes human skin lipids for presentation by CD1a. <i>Journal of Experimental Medicine</i> , 2015, 212, 149-163.	8.5	98
13	Functionality of Dengue Virus Specific Memory T Cell Responses in Individuals Who Were Hospitalized or Who Had Mild or Subclinical Dengue Infection. <i>PLoS Neglected Tropical Diseases</i> , 2015, 9, e0003673.	3.0	41
14	Prostaglandin D2 and leukotriene E4 synergize to stimulate diverse TH2 functions and TH2 cell/neutrophil crosstalk. <i>Journal of Allergy and Clinical Immunology</i> , 2015, 135, 1358-1366.e11.	2.9	50
15	MHCII-Mediated Dialog between Group 2 Innate Lymphoid Cells and CD4+ T Cells Potentiates Type 2 Immunity and Promotes Parasitic Helminth Expulsion. <i>Immunity</i> , 2014, 41, 283-295.	14.3	601
16	Prostaglandin D2 activates group 2 innate lymphoid cells through chemoattractant receptor-homologous molecule expressed on TH2 cells. <i>Journal of Allergy and Clinical Immunology</i> , 2014, 133, 1184-1194.e7.	2.9	433
17	A role for IL-25 and IL-33-driven type-2 innate lymphoid cells in atopic dermatitis. <i>Journal of Experimental Medicine</i> , 2013, 210, 2939-2950.	8.5	803