

Gary Mitchell Crowfoot

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8935865/publications.pdf>

Version: 2024-02-01

16
papers

119
citations

1307594

7
h-index

1372567

10
g-index

16
all docs

16
docs citations

16
times ranked

198
citing authors

#	ARTICLE	IF	CITATIONS
1	Frequent, short bouts of light-intensity exercises while standing decreases systolic blood pressure: Breaking Up Sitting Time after Stroke (BUST-Stroke) trial. <i>International Journal of Stroke</i> , 2018, 13, 932-940.	5.9	37
2	Real-life experiences of people with transient ischaemic attack or minor stroke: A qualitative literature review. <i>Journal of Clinical Nursing</i> , 2018, 27, 1381-1398.	3.0	14
3	Breaking up sitting time after stroke (BUST-stroke). <i>International Journal of Stroke</i> , 2018, 13, 921-931.	5.9	14
4	What is the effect of interrupting prolonged sitting with frequent bouts of physical activity or standing on first or recurrent stroke risk factors? A scoping review. <i>PLoS ONE</i> , 2019, 14, e0217981.	2.5	14
5	Participation, Fear of Falling, and Upper Limb Impairment are Associated with High Sitting Time in People with Stroke. <i>Occupational Therapy in Health Care</i> , 2019, 33, 181-196.	0.3	10
6	Emergency healthcare workers' preparedness for disaster management: An integrative review. <i>Journal of Clinical Nursing</i> , 2021, , .	3.0	9
7	Living within stories: Exploring the experiences of people with transient ischemic attack. <i>Australian Journal of Cancer Nursing</i> , 2016, 18, 52-57.	1.6	7
8	Patient readiness for risk-reduction education and lifestyle change following transient ischemic attack. <i>Disability and Rehabilitation</i> , 2021, 43, 400-405.	1.8	3
9	What do stroke survivors value about participating in research and what are the most important research problems related to stroke or transient ischemic attack (TIA)? A survey. <i>BMC Medical Research Methodology</i> , 2021, 21, 209.	3.1	3
10	Breaking up sitting time after stroke – How much less sitting is needed to improve blood pressure after stroke (BUST-BP-Dose): Protocol for a dose-finding study. <i>Contemporary Clinical Trials Communications</i> , 2019, 13, 100310.	1.1	2
11	Investigating the rigour of research findings in experimental studies assessing the effects of breaking up prolonged sitting – extended scoping review. <i>Brazilian Journal of Physical Therapy</i> , 2021, 25, 4-16.	2.5	2
12	Acute Effects of Frequent Light-Intensity Standing-Based Exercises That Interrupt 8 Hours of Prolonged Sitting on Postprandial Glucose in Stroke Survivors: A Dose-Escalation Trial. <i>Journal of Physical Activity and Health</i> , 2021, 18, 644-652.	2.0	2
13	Participants' Perspective of Engaging in a Gym-Based Health Service Delivered Secondary Stroke Prevention Program after TIA or Mild Stroke. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 11448.	2.6	2
14	Sepsis 2016 Paris. <i>Critical Care</i> , 2016, 20, .	5.8	0
15	Comparing the physical activity of stroke survivors in high-income countries and low to middle-income countries. <i>Physiotherapy Research International</i> , 2021, 26, e1918.	1.5	0
16	The Effects of Interrupting Prolonged Sitting With Frequent Bouts of Light-Intensity Standing Exercises on Blood Pressure in Stroke Survivors: A Dose Escalation Trial. <i>Journal of Physical Activity and Health</i> , 2021, 18, 988-997.	2.0	0