

Man Fai Lam

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8945497/publications.pdf>

Version: 2024-02-01

14
papers

222
citations

1163117

8
h-index

1125743

13
g-index

14
all docs

14
docs citations

14
times ranked

284
citing authors

#	ARTICLE	IF	CITATIONS
1	A Longitudinal Study on the Prevalence and Risk Factors for Depression and Anxiety, Quality of Life, and Clinical Outcomes in Incident Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2019, 39, 74-82.	2.3	17
2	Risk factors and prognosis of late acute rejection in Chinese kidney transplant recipients. <i>Nephrology</i> , 2017, 22, 985-992.	1.6	0
3	Psychosocial Factors Predict Nonadherence to PD Treatment: A Hong Kong Survey. <i>Peritoneal Dialysis International</i> , 2017, 37, 331-337.	2.3	6
4	A formative research as the groundwork for designing evidence-based adherence promotion campaigns for patients on peritoneal dialysis. <i>Journal of Communication in Healthcare</i> , 2016, 9, 98-108.	1.5	2
5	Prevalence of Cognitive Impairment among Peritoneal Dialysis Patients, Impact on Peritonitis and Role of Assisted Dialysis. <i>Peritoneal Dialysis International</i> , 2016, 36, 284-290.	2.3	26
6	Novel genes and variants associated with IgA nephropathy by co-segregating with the disease phenotypes in 10 IgAN families. <i>Gene</i> , 2015, 571, 43-51.	2.2	14
7	Macromolecular IgA1 taken from patients with familial IgA Nephropathy or their asymptomatic relatives have higher reactivity to mesangial cells in vitro. <i>Kidney International</i> , 2009, 75, 1330-1339.	5.2	32
8	Procalcitonin Fails to Differentiate Inflammatory Status or Predict Long-Term Outcomes in Peritoneal Dialysis-Associated Peritonitis. <i>Peritoneal Dialysis International</i> , 2008, 28, 377-384.	2.3	19
9	Ampicillin-sulbactam and amikacin used as second-line antibiotics for patients with culture-negative peritonitis. <i>Peritoneal Dialysis International</i> , 2008, 28, 540-2.	2.3	3
10	Hyperleptinaemia and chronic inflammation after peritonitis predicts poor nutritional status and mortality in patients on peritoneal dialysis. <i>Nephrology Dialysis Transplantation</i> , 2007, 22, 1445-1450.	0.7	35
11	ASPD: A prospective study of adequacy in Asian patients on long term, small volume, continuous ambulatory peritoneal dialysis. <i>Peritoneal Dialysis International</i> , 2006, 26, 466-74.	2.3	7
12	Mannose binding lectin level and polymorphism in patients on long-term peritoneal dialysis. <i>Nephrology Dialysis Transplantation</i> , 2005, 20, 2489-2496.	0.7	14
13	Family-Based Association Study Showing that Immunoglobulin A Nephropathy Is Associated with the Polymorphisms 2093C and 2180T in the 3' Untranslated Region of the Megin Gene. <i>Journal of the American Society of Nephrology: JASN</i> , 2004, 15, 1739-1743.	6.1	45
14	PERINEPHRIC HEMATOMA AND MASSIVE GASTROINTESTINAL HEMORRHAGE (GIB) COMPLICATING POLYARTERITIS NODOSA (PAN). <i>Renal Failure</i> , 2000, 22, 379-384.	2.1	2