

Miguel Remacha Moreno

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8968365/publications.pdf>

Version: 2024-02-01

49
papers

3,103
citations

218677

26
h-index

214800

47
g-index

50
all docs

50
docs citations

50
times ranked

1507
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | The complete DNA sequence of yeast chromosome III. <i>Nature</i> , 1992, 357, 38-46. | 27.8 | 924 |
| 2 | Complete DNA sequence of yeast chromosome XI. <i>Nature</i> , 1994, 369, 371-378. | 27.8 | 382 |
| 3 | Ribosomal Acidic Phosphoproteins P1 and P2 Are Not Required for Cell Viability but Regulate the Pattern of Protein Expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 4754-4762. | 2.3 | 124 |
| 4 | The Large Ribosomal Subunit Stalk as a Regulatory Element of the Eukaryotic Translational Machinery. <i>Progress in Molecular Biology and Translational Science</i> , 1996, 55, 157-193. | 1.9 | 116 |
| 5 | The nucleotide sequence of <i>Saccharomyces cerevisiae</i> chromosome IV. <i>Nature</i> , 1997, 387, 75-78. | 27.8 | 95 |
| 6 | Independent genes coding for three acidic proteins of the large ribosomal subunit from <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1988, 263, 9094-9101. | 3.4 | 94 |
| 7 | Proteins P1, P2, and P0, components of the eukaryotic ribosome stalk. New structural and functional aspects. <i>Biochemistry and Cell Biology</i> , 1995, 73, 959-968. | 2.0 | 92 |
| 8 | Independent genes coding for three acidic proteins of the large ribosomal subunit from <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1988, 263, 9094-101. | 3.4 | 82 |
| 9 | The acidic ribosomal proteins as regulators of the eukaryotic ribosomal activity. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1990, 1050, 51-55. | 2.4 | 75 |
| 10 | Characterization of the yeast acidic ribosomal phosphoproteins using monoclonal antibodies. Proteins L44/L45 and L44' have different functional roles. <i>FEBS Journal</i> , 1991, 196, 407-414. | 0.2 | 72 |
| 11 | Stable binding of the eukaryotic acidic phosphoproteins to the ribosome is not an absolute requirement for in vivo protein synthesis.. <i>Journal of Biological Chemistry</i> , 1992, 267, 12061-12067. | 3.4 | 71 |
| 12 | Phosphorylation of the Acidic Ribosomal P Proteins in <i>Saccharomyces cerevisiae</i> : A Reappraisal. <i>Biochemistry</i> , 1997, 36, 14439-14446. | 2.5 | 65 |
| 13 | The nucleotide sequence of <i>Saccharomyces cerevisiae</i> chromosome XIV and its evolutionary implications. <i>Nature</i> , 1997, 387, 93-98. | 27.8 | 65 |
| 14 | Phosphorylation and N-terminal region of yeast ribosomal protein P1 mediate its degradation, which is prevented by protein P2. <i>EMBO Journal</i> , 2000, 19, 6075-6084. | 7.8 | 63 |
| 15 | Stable binding of the eukaryotic acidic phosphoproteins to the ribosome is not an absolute requirement for in vivo protein synthesis. <i>Journal of Biological Chemistry</i> , 1992, 267, 12061-7. | 3.4 | 55 |
| 16 | The GTPase Center Protein L12 Is Required for Correct Ribosomal Stalk Assembly but Not for <i>Saccharomyces cerevisiae</i> Viability. <i>Journal of Biological Chemistry</i> , 1998, 273, 31956-31961. | 3.4 | 51 |
| 17 | Assembly of <i>Saccharomyces cerevisiae</i> Ribosomal Stalk: Binding of P1 Proteins Is Required for the Interaction of P2 Proteins. <i>Biochemistry</i> , 2000, 39, 8929-8934. | 2.5 | 48 |
| 18 | The RNA Interacting Domain but Not the Protein Interacting Domain Is Highly Conserved in Ribosomal Protein P0. <i>Journal of Biological Chemistry</i> , 2000, 275, 2130-2136. | 3.4 | 47 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 19 | Asymmetric Interactions between the Acidic P1 and P2 Proteins in the <i>Saccharomyces cerevisiae</i> Ribosomal Stalk. <i>Journal of Biological Chemistry</i> , 2001, 276, 32474-32479. | 3.4 | 47 |
| 20 | Disruption of single-copy genes encoding acidic ribosomal proteins in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 2182-2190. | 2.3 | 45 |
| 21 | Phosphorylation of Ribosomal Protein P0 Is Not Essential for Ribosome Function but Can Affect Translation. <i>Biochemistry</i> , 1998, 37, 16620-16626. | 2.5 | 45 |
| 22 | Phosphorylation of the yeast ribosomal stalk. Functional effects and enzymes involved in the process. <i>FEMS Microbiology Reviews</i> , 1999, 23, 537-550. | 8.6 | 39 |
| 23 | Characterization of interaction sites in the <i>Saccharomyces cerevisiae</i> ribosomal stalk components. <i>Molecular Microbiology</i> , 2002, 46, 719-792. | 2.5 | 39 |
| 24 | Structure-activity relationships of sparsomycin and its analogs. Inhibition of peptide bond formation in cell-free systems and of L1210 and bacterial cell growth. <i>Journal of Medicinal Chemistry</i> , 1987, 30, 325-333. | 6.4 | 34 |
| 25 | Eukaryotic acidic phosphoproteins interact with the ribosome through their amino-terminal domain. <i>Biochemistry</i> , 1995, 34, 7941-7948. | 2.5 | 34 |
| 26 | The acidic phosphoproteins from <i>Saccharomyces cerevisiae</i> ribosomes. Amino terminal acetylation is a conserved difference between P1 and P2 proteins. <i>Biochemistry</i> , 1993, 32, 4231-4236. | 2.5 | 27 |
| 27 | Ribosomal protein interactions in yeast. Protein L15 forms a complex with the acidic proteins. <i>FEBS Journal</i> , 1988, 177, 531-537. | 0.2 | 26 |
| 28 | Disruption of six <i>Saccharomyces cerevisiae</i> novel genes and phenotypic analysis of the deletants. <i>Yeast</i> , 1999, 15, 945-953. | 1.7 | 24 |
| 29 | Structural Differences between <i>Saccharomyces cerevisiae</i> Ribosomal Stalk Proteins P1 and P2 Support Their Functional Diversity. <i>Biochemistry</i> , 2000, 39, 8935-8943. | 2.5 | 22 |
| 30 | Ribosomal P0 Protein Domain Involved in Selectivity of Antifungal Sordarin Derivatives. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 2930-2936. | 3.2 | 21 |
| 31 | <i>Chlamydia trachomatis</i> RNA polymerase alpha subunit: sequence and structural analysis. <i>Journal of Bacteriology</i> , 1995, 177, 2594-2601. | 2.2 | 19 |
| 32 | The 26S rRNA binding ribosomal protein equivalent to bacterial protein L11 is encoded by unspliced duplicated genes in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1990, 18, 4409-4416. | 14.5 | 17 |
| 33 | The activity-controlling phosphorylation site is not the same in the four acidic ribosomal proteins from <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1993, 268, 2451-2457. | 3.4 | 14 |
| 34 | Effect of acidic ribosomal phosphoprotein mRNA 5'-untranslated region on gene expression and protein accumulation.. <i>Journal of Biological Chemistry</i> , 1994, 269, 3968-3975. | 3.4 | 12 |
| 35 | Functional domains of chlamydial histone H1-like protein. <i>Biochemical Journal</i> , 1996, 315, 481-486. | 3.7 | 11 |
| 36 | Carboxy terminal modifications of the P0 protein reveal alternative mechanisms of nuclear ribosomal stalk assembly. <i>Nucleic Acids Research</i> , 2013, 41, 8628-8636. | 14.5 | 11 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Disruption of single-copy genes encoding acidic ribosomal proteins in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 2182-2190. | 2.3 | 10 |
| 38 | Effect of acidic ribosomal phosphoprotein mRNA 5'-untranslated region on gene expression and protein accumulation. <i>Journal of Biological Chemistry</i> , 1994, 269, 3968-75. | 3.4 | 10 |
| 39 | Ribosomal Stalk Protein Phosphorylating Activities in <i>Saccharomyces cerevisiae</i> . <i>Archives of Biochemistry and Biophysics</i> , 2000, 375, 83-89. | 3.0 | 9 |
| 40 | The Acidic Ribosomal Proteins and the Control of Protein Synthesis in Yeast. , 1993, , 67-80. | | 9 |
| 41 | The activity-controlling phosphorylation site is not the same in the four acidic ribosomal proteins from <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1993, 268, 2451-7. | 3.4 | 9 |
| 42 | The complete sequence of a 15 820 bp segment of <i>Saccharomyces cerevisiae</i> chromosome XI contains the <i>UBI2</i> and <i>MPL1</i> genes and three new open reading frames. <i>Yeast</i> , 1993, 9, 1349-1354. | 1.7 | 8 |
| 43 | Structure and function of the stalk, a putative regulatory element of the yeast ribosome. Role of stalk protein phosphorylation. <i>Folia Microbiologica</i> , 1999, 44, 153-163. | 2.3 | 8 |
| 44 | Deletion of 24 open reading frames from chromosome XI from <i>Saccharomyces cerevisiae</i> and phenotypic analysis of the deletants. <i>Gene</i> , 1999, 233, 141-150. | 2.2 | 7 |
| 45 | Chromosome location of a family of genes encoding different acidic ribosomal proteins in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1990, 17, 535-536. | 1.7 | 6 |
| 46 | Cloning and characterization of a <i>secY</i> homolog from <i>Chlamydia trachomatis</i> . <i>Molecular Genetics and Genomics</i> , 1994, 243, 482-487. | 2.4 | 6 |
| 47 | The sequence of a 17 933 bp segment of <i>Saccharomyces cerevisiae</i> chromosome XIV contains the <i>RHO2</i> , <i>TOP2</i> , <i>MKT1</i> and <i>END3</i> genes and five new open reading frames. , 1996, 12, 485-491. | | 6 |
| 48 | Heterologous expression of the highly conserved acidic ribosomal phosphoproteins from <i>Dictyostelium discoideum</i> in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1995, 1263, 45-52. | 2.4 | 5 |
| 49 | Characterization of Sparsomycin Resistance in <i>Streptomyces sparsogenes</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2002, 46, 2914-2919. | 3.2 | 2 |