

Muhammad Abdul-Ghani

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9019106/publications.pdf>

Version: 2024-02-01

199
papers

55,778
citations

2427

97
h-index

2509

196
g-index

202
all docs

202
docs citations

202
times ranked

35647
citing authors

#	ARTICLE	IF	CITATIONS
1	Insulin sensitivity indices obtained from oral glucose tolerance testing: comparison with the euglycemic insulin clamp.. Diabetes Care, 1999, 22, 1462-1470.	8.6	4,832
2	Glucose clamp technique: a method for quantifying insulin secretion and resistance.. American Journal of Physiology - Endocrinology and Metabolism, 1979, 237, E214.	3.5	4,022
3	Insulin Resistance: A Multifaceted Syndrome Responsible for NIDDM, Obesity, Hypertension, Dyslipidemia, and Atherosclerotic Cardiovascular Disease. Diabetes Care, 1991, 14, 173-194.	8.6	3,723
4	The Triumvirate: β -Cell, Muscle, Liver: A Collusion Responsible for NIDDM. Diabetes, 1988, 37, 667-687.	0.6	2,466
5	From the Triumvirate to the Ominous Octet: A New Paradigm for the Treatment of Type 2 Diabetes Mellitus. Diabetes, 2009, 58, 773-795.	0.6	2,251
6	Coordinated reduction of genes of oxidative metabolism in humans with insulin resistance and diabetes: Potential role of <i>PGC1α</i> and <i>NRF1</i> . Proceedings of the National Academy of Sciences of the United States of America, 2003, 100, 8466-8471.	7.1	1,800
7	A Placebo-Controlled Trial of Pioglitazone in Subjects with Nonalcoholic Steatohepatitis. New England Journal of Medicine, 2006, 355, 2297-2307.	27.0	1,584
8	Type 2 diabetes mellitus. Nature Reviews Disease Primers, 2015, 1, 15019.	30.5	1,308
9	Efficacy of Metformin in Patients with Non-Insulin-Dependent Diabetes Mellitus. New England Journal of Medicine, 1995, 333, 541-549.	27.0	1,213
10	Quantitation of Muscle Glycogen Synthesis in Normal Subjects and Subjects with Non-Insulin-Dependent Diabetes by ^{13}C Nuclear Magnetic Resonance Spectroscopy. New England Journal of Medicine, 1990, 322, 223-228.	27.0	1,181
11	Effects of insulin on peripheral and splanchnic glucose metabolism in noninsulin-dependent (type II) diabetes mellitus.. Journal of Clinical Investigation, 1985, 76, 149-155.	8.2	974
12	Insulin resistance differentially affects the PI 3-kinase α and MAP kinase α -mediated signaling in human muscle. Journal of Clinical Investigation, 2000, 105, 311-320.	8.2	953
13	Glucose Toxicity. Diabetes Care, 1990, 13, 610-630.	8.6	889
14	Correction of hyperglycemia with phlorizin normalizes tissue sensitivity to insulin in diabetic rats.. Journal of Clinical Investigation, 1987, 79, 1510-1515.	8.2	722
15	Glucose and free fatty acid metabolism in non-insulin-dependent diabetes mellitus. Evidence for multiple sites of insulin resistance.. Journal of Clinical Investigation, 1989, 84, 205-213.	8.2	710
16	Insulin resistance, lipotoxicity, type 2 diabetes and atherosclerosis: the missing links. The Claude Bernard Lecture 2009. Diabetologia, 2010, 53, 1270-1287.	6.3	678
17	Dapagliflozin improves muscle insulin sensitivity but enhances endogenous glucose production. Journal of Clinical Investigation, 2014, 124, 509-514.	8.2	661
18	Non-Alcoholic Fatty Liver Disease (NAFLD) and Its Connection with Insulin Resistance, Dyslipidemia, Atherosclerosis and Coronary Heart Disease. Nutrients, 2013, 5, 1544-1560.	4.1	648

#	ARTICLE	IF	CITATIONS
19	Pioglitazone for Diabetes Prevention in Impaired Glucose Tolerance. <i>New England Journal of Medicine</i> , 2011, 364, 1104-1115.	27.0	646
20	Effect of Pioglitazone on Abdominal Fat Distribution and Insulin Sensitivity in Type 2 Diabetic Patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2002, 87, 2784-2791.	3.6	629
21	Role of the Adipocyte, Free Fatty Acids, and Ectopic Fat in Pathogenesis of Type 2 Diabetes Mellitus: Peroxisomal Proliferator-Activated Receptor Agonists Provide a Rational Therapeutic Approach. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004, 89, 463-478.	3.6	570
22	Regulation of Splanchnic and Peripheral Glucose Uptake by Insulin and Hyperglycemia in Man. <i>Diabetes</i> , 1983, 32, 35-45.	0.6	548
23	Relationship Between Hepatic/Visceral Fat and Hepatic Insulin Resistance in Nondiabetic and Type 2 Diabetic Subjects. <i>Gastroenterology</i> , 2007, 133, 496-506.	1.3	500
24	Fasting hyperglycemia in non-insulin-dependent diabetes mellitus: Contributions of excessive hepatic glucose production and impaired tissue glucose uptake. <i>Metabolism: Clinical and Experimental</i> , 1989, 38, 387-395.	3.4	492
25	Î²-Cell Function in Subjects Spanning the Range from Normal Glucose Tolerance to Overt Diabetes: A New Analysis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005, 90, 493-500.	3.6	470
26	A Sustained Increase in Plasma Free Fatty Acids Impairs Insulin Secretion in Nondiabetic Subjects Genetically Predisposed to Develop Type 2 Diabetes. <i>Diabetes</i> , 2003, 52, 2461-2474.	0.6	447
27	Pathogenesis of Insulin Resistance in Skeletal Muscle. <i>Journal of Biomedicine and Biotechnology</i> , 2010, 2010, 1-19.	3.0	441
28	Insulin Secretion and Action in Subjects With Impaired Fasting Glucose and Impaired Glucose Tolerance: Results From the Veterans Administration Genetic Epidemiology Study. <i>Diabetes</i> , 2006, 55, 1430-1435.	0.6	429
29	The role of the kidneys in glucose homeostasis: a new path towards normalizing glycaemia. <i>Diabetes, Obesity and Metabolism</i> , 2012, 14, 5-14.	4.4	398
30	Renal, metabolic and cardiovascular considerations of SGLT2 inhibition. <i>Nature Reviews Nephrology</i> , 2017, 13, 11-26.	9.6	398
31	The effects of glucose and insulin on renal electrolyte transport.. <i>Journal of Clinical Investigation</i> , 1976, 58, 83-90.	8.2	398
32	Contributions of Î²-Cell Dysfunction and Insulin Resistance to the Pathogenesis of Impaired Glucose Tolerance and Impaired Fasting Glucose. <i>Diabetes Care</i> , 2006, 29, 1130-1139.	8.6	382
33	Insulin resistance and endothelial dysfunction: the road map to cardiovascular diseases. <i>Diabetes/Metabolism Research and Reviews</i> , 2006, 22, 423-436.	4.0	373
34	Improved Glycemic Control and Enhanced Insulin Sensitivity in Type 2 Diabetic Subjects Treated With Pioglitazone. <i>Diabetes Care</i> , 2001, 24, 710-719.	8.6	367
35	Mechanism of Metformin Action in Obese and Lean Noninsulin-Dependent Diabetic Subjects*. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1991, 73, 1294-1301.	3.6	363
36	Consensus Statement by the American Association of Clinical Endocrinologists and American College of Endocrinology on the Comprehensive type 2 Diabetes Management Algorithm â€” 2017 Executive Summary. <i>Endocrine Practice</i> , 2017, 23, 207-238.	2.1	362

#	ARTICLE	IF	CITATIONS
37	Metabolic effects of metformin on glucose and lactate metabolism in noninsulin-dependent diabetes mellitus.. Journal of Clinical Endocrinology and Metabolism, 1996, 81, 4059-4067.	3.6	349
38	Role of Sodium-Glucose Cotransporter 2 (SGLT 2) Inhibitors in the Treatment of Type 2 Diabetes. Endocrine Reviews, 2011, 32, 515-531.	20.1	344
39	Effects of exenatide versus sitagliptin on postprandial glucose, insulin and glucagon secretion, gastric emptying, and caloric intake: a randomized, cross-over study. Current Medical Research and Opinion, 2008, 24, 2943-2952.	1.9	341
40	Decreased Plasma Adiponectin Concentrations Are Closely Related to Hepatic Fat Content and Hepatic Insulin Resistance in Pioglitazone-Treated Type 2 Diabetic Patients. Journal of Clinical Endocrinology and Metabolism, 2004, 89, 200-206.	3.6	340
41	Obesity and insulin resistance in humans: A dose-response study. Metabolism: Clinical and Experimental, 1990, 39, 452-459.	3.4	333
42	Dose-Response Effect of Elevated Plasma Free Fatty Acid on Insulin Signaling. Diabetes, 2005, 54, 1640-1648.	0.6	333
43	Glucose Intolerance and Aging. Diabetes Care, 1981, 4, 493-501.	8.6	329
44	Elevated Toll-Like Receptor 4 Expression and Signaling in Muscle From Insulin-Resistant Subjects. Diabetes, 2008, 57, 2595-2602.	0.6	319
45	Effect of chronic hyperglycemia on in vivo insulin secretion in partially pancreatectomized rats.. Journal of Clinical Investigation, 1987, 80, 1037-1044.	8.2	312
46	Beta-cell dysfunction and glucose intolerance: results from the San Antonio metabolism (SAM) study. Diabetologia, 2004, 47, 31-39.	6.3	287
47	Effect of sustained physiologic hyperinsulinaemia and hyperglycaemia on insulin secretion and insulin sensitivity in man. Diabetologia, 1994, 37, 1025-1035.	6.3	280
48	Influence of hyperinsulinemia, hyperglycemia, and the route of glucose administration on splanchnic glucose exchange. Proceedings of the National Academy of Sciences of the United States of America, 1978, 75, 5173-5177.	7.1	279
49	The Metabolic Profile of NIDDM Is Fully Established in Glucose-Tolerant Offspring of Two Mexican-American NIDDM Parents. Diabetes, 1992, 41, 1575-1586.	0.6	277
50	The disposal of an oral glucose load in patients with non-insulin-dependent diabetes. Metabolism: Clinical and Experimental, 1988, 37, 79-85.	3.4	268
51	Pioglitazone Reduces Hepatic Fat Content and Augments Splanchnic Glucose Uptake in Patients With Type 2 Diabetes. Diabetes, 2003, 52, 1364-1370.	0.6	265
52	A Meta-analysis Comparing the Effect of Thiazolidinediones on Cardiovascular Risk Factors. Archives of Internal Medicine, 2004, 164, 2097.	3.8	264
53	Role of Lipid Oxidation in Pathogenesis of Insulin Resistance of Obesity and Type II Diabetes. Diabetes, 1987, 36, 1341-1350.	0.6	259
54	Effect of rosiglitazone on glucose and non-esterified fatty acid metabolism in Type II diabetic patients. Diabetologia, 2001, 44, 2210-2219.	6.3	253

#	ARTICLE	IF	CITATIONS
55	SGLT2 Inhibitors and Cardiovascular Risk: Lessons Learned From the EMPA-REG OUTCOME Study. <i>Diabetes Care</i> , 2016, 39, 717-725.	8.6	244
56	Pathophysiology of diabetic kidney disease: impact of SGLT2 inhibitors. <i>Nature Reviews Nephrology</i> , 2021, 17, 319-334.	9.6	244
57	Combination of Empagliflozin and Linagliptin as Second-Line Therapy in Subjects With Type 2 Diabetes Inadequately Controlled on Metformin. <i>Diabetes Care</i> , 2015, 38, 384-393.	8.6	241
58	The Primary Glucose-Lowering Effect of Metformin Resides in the Gut, Not the Circulation: Results From Short-term Pharmacokinetic and 12-Week Dose-Ranging Studies. <i>Diabetes Care</i> , 2016, 39, 198-205.	8.6	240
59	Insulin Binding to Monocytes and Insulin Action in Human Obesity, Starvation, and Refeeding. <i>Journal of Clinical Investigation</i> , 1978, 62, 204-213.	8.2	235
60	Role of Adipose Tissue Insulin Resistance in the Natural History of Type 2 Diabetes: Results From the San Antonio Metabolism Study. <i>Diabetes</i> , 2017, 66, 815-822.	0.6	234
61	Characterization of Renal Glucose Reabsorption in Response to Dapagliflozin in Healthy Subjects and Subjects With Type 2 Diabetes. <i>Diabetes Care</i> , 2013, 36, 3169-3176.	8.6	233
62	Insulin resistance: a multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidaemia and atherosclerosis. <i>Netherlands Journal of Medicine</i> , 1997, 50, 191-197.	0.5	232
63	Bromocriptine: a novel approach to the treatment of type 2 diabetes.. <i>Diabetes Care</i> , 2000, 23, 1154-1161.	8.6	224
64	Effect of insulin on the distribution and disposition of glucose in man.. <i>Journal of Clinical Investigation</i> , 1985, 76, 357-364.	8.2	223
65	Physical training and insulin sensitivity. <i>Diabetes/metabolism Reviews</i> , 1986, 1, 445-481.	0.3	222
66	Lipid Infusion Decreases the Expression of Nuclear Encoded Mitochondrial Genes and Increases the Expression of Extracellular Matrix Genes in Human Skeletal Muscle. <i>Journal of Biological Chemistry</i> , 2005, 280, 10290-10297.	3.4	217
67	Insulin Resistance and Atherosclerosis: Implications for Insulin-Sensitizing Agents. <i>Endocrine Reviews</i> , 2019, 40, 1447-1467.	20.1	210
68	Bromocriptine: A Sympatholytic, D2-Dopamine Agonist for the Treatment of Type 2 Diabetes. <i>Diabetes Care</i> , 2011, 34, 789-794.	8.6	209
69	Preservation of β -Cell Function: The Key to Diabetes Prevention. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011, 96, 2354-2366.	3.6	207
70	Indirect calorimetry: methodological and interpretative problems. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 1990, 258, E399-E412.	3.5	200
71	Novel Hypothesis to Explain Why SGLT2 Inhibitors Inhibit Only 30-50% of Filtered Glucose Load in Humans. <i>Diabetes</i> , 2013, 62, 3324-3328.	0.6	198
72	Insulin resistance and coronary artery disease. <i>Diabetologia</i> , 1996, 39, 1345-1350.	6.3	187

#	ARTICLE	IF	CITATIONS
73	Relationship of baseline HbA _{1c} and efficacy of current glucose-lowering therapies: a meta-analysis of randomized clinical trials. <i>Diabetic Medicine</i> , 2010, 27, 309-317.	2.3	183
74	Dysfunctional fat cells, lipotoxicity and type 2 diabetes. <i>International Journal of Clinical Practice</i> , 2004, 58, 9-21.	1.7	175
75	Thiazolidinediones improve β -cell function in type 2 diabetic patients. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2007, 292, E871-E883.	3.5	167
76	Effect of a Sustained Reduction in Plasma Free Fatty Acid Concentration on Intramuscular Long-Chain Fatty Acyl-CoAs and Insulin Action in Type 2 Diabetic Patients. <i>Diabetes</i> , 2005, 54, 3148-3153.	0.6	162
77	Initial combination therapy with metformin, pioglitazone and exenatide is more effective than sequential add-on therapy in subjects with new-onset diabetes. Results from the efficacy and durability of initial combination therapy for type 2 diabetes (EDICT): a randomized trial. <i>Diabetes, Obesity and Metabolism</i> , 2015, 17, 268-275.	4.4	160
78	Dapagliflozin Enhances Fat Oxidation and Ketone Production in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2016, 39, 2036-2041.	8.6	155
79	Pioglitazone: The forgotten, cost-effective cardioprotective drug for type 2 diabetes. <i>Diabetes and Vascular Disease Research</i> , 2019, 16, 133-143.	2.0	155
80	Normalization of blood glucose in diabetic rats with phlorizin treatment reverses insulin-resistant glucose transport in adipose cells without restoring glucose transporter gene expression.. <i>Journal of Clinical Investigation</i> , 1991, 87, 561-570.	8.2	155
81	Effect of Physical Training on Insulin Action in Obesity. <i>Diabetes</i> , 1987, 36, 1379-1385.	0.6	153
82	Total Body Fat Content and Fat Topography Are Associated Differently With In Vivo Glucose Metabolism in Nonobese and Obese Nondiabetic Women. <i>Diabetes</i> , 1992, 41, 1151-1159.	0.6	145
83	Mechanism of action of exenatide to reduce postprandial hyperglycemia in type 2 diabetes. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2008, 294, E846-E852.	3.5	144
84	Dapagliflozin Lowers Plasma Glucose Concentration and Improves β -Cell Function. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 1927-1932.	3.6	133
85	Rosiglitazone Improves Downstream Insulin Receptor Signaling in Type 2 Diabetic Patients. <i>Diabetes</i> , 2003, 52, 1943-1950.	0.6	128
86	Pioglitazone stimulates AMP-activated protein kinase signalling and increases the expression of genes involved in adiponectin signalling, mitochondrial function and fat oxidation in human skeletal muscle in vivo: a randomised trial. <i>Diabetologia</i> , 2009, 52, 723-732.	6.3	127
87	Insulin Resistance Is Associated with Impaired Nitric Oxide Synthase Activity in Skeletal Muscle of Type 2 Diabetic Subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005, 90, 1100-1105.	3.6	124
88	Reduced Skeletal Muscle Inhibitor of β -Tubulin Content Is Associated With Insulin Resistance in Subjects With Type 2 Diabetes: Reversal by Exercise Training. <i>Diabetes</i> , 2006, 55, 760-767.	0.6	124
89	Effects of peroxisome proliferator-activated receptor (PPAR)- α and PPAR- γ agonists on glucose and lipid metabolism in patients with type 2 diabetes mellitus. <i>Diabetologia</i> , 2007, 50, 1723-1731.	6.3	124
90	Combination therapy with GLP-1 receptor agonist and SGLT2 inhibitor. <i>Diabetes, Obesity and Metabolism</i> , 2017, 19, 1353-1362.	4.4	120

#	ARTICLE	IF	CITATIONS
91	The role of fractional glucose extraction in the regulation of splanchnic glucose metabolism in normal and diabetic man. <i>Metabolism: Clinical and Experimental</i> , 1980, 29, 28-35.	3.4	117
92	Increased Glutamine:Fructose-6-Phosphate Amidotransferase Activity in Skeletal Muscle of Patients With NIDDM. <i>Diabetes</i> , 1996, 45, 302-307.	0.6	113
93	Renal sodium-glucose cotransporter inhibition in the management of type 2 diabetes mellitus. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 309, F889-F900.	2.7	113
94	Cardiovascular Disease and Type 2 Diabetes: Has the Dawn of a New Era Arrived?. <i>Diabetes Care</i> , 2017, 40, 813-820.	8.6	109
95	Predominant role of reduced beta-cell sensitivity to glucose over insulin resistance in impaired glucose tolerance. <i>Diabetologia</i> , 2003, 46, 1211-1219.	6.3	103
96	Effect of loss of first-phase insulin secretion on hepatic glucose production and tissue glucose disposal in humans. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 1989, 257, E241-E246.	3.5	101
97	Skeletal Muscle Insulin Resistance in Normoglycemic Subjects With a Strong Family History of Type 2 Diabetes Is Associated With Decreased Insulin-Stimulated Insulin Receptor Substrate-1 Tyrosine Phosphorylation. <i>Diabetes</i> , 2001, 50, 2572-2578.	0.6	100
98	Effects of Exenatide Plus Rosiglitazone on β -Cell Function and Insulin Sensitivity in Subjects With Type 2 Diabetes on Metformin. <i>Diabetes Care</i> , 2010, 33, 951-957.	8.6	100
99	Insulin Reduces Plasma Arginase Activity in Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2008, 31, 134-139.	8.6	97
100	Pioglitazone Slows Progression of Atherosclerosis in Prediabetes Independent of Changes in Cardiovascular Risk Factors. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013, 33, 393-399.	2.4	97
101	Metabolic and molecular basis of insulin resistance. <i>Journal of Nuclear Cardiology</i> , 2003, 10, 311-323.	2.1	96
102	Distinct β -Cell Defects in Impaired Fasting Glucose and Impaired Glucose Tolerance. <i>Diabetes</i> , 2012, 61, 447-453.	0.6	96
103	Once-daily delayed-release metformin lowers plasma glucose and enhances fasting and postprandial GLP-1 and PYY: results from two randomised trials. <i>Diabetologia</i> , 2016, 59, 1645-1654.	6.3	95
104	In Vivo Actions of Peroxisome Proliferator-Activated Receptors. <i>Diabetes Care</i> , 2013, 36, S162-S174.	8.6	94
105	Discordant effects of a chronic physiological increase in plasma FFA on insulin signaling in healthy subjects with or without a family history of type 2 diabetes. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2004, 287, E537-E546.	3.5	89
106	Efficacy and Tolerability of the DPP-4 Inhibitor Alogliptin Combined with Pioglitazone, in Metformin-Treated Patients with Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, 1615-1622.	3.6	88
107	Prevention of Diabetes With Pioglitazone in ACT NOW. <i>Diabetes</i> , 2013, 62, 3920-3926.	0.6	83
108	A Test of the Hypothesis that the Rate of Fall in Glucose Concentration Triggers Counterregulatory Hormonal Responses in Man. <i>Diabetes</i> , 1977, 26, 445-452.	0.6	80

#	ARTICLE	IF	CITATIONS
109	Reduction in Hematocrit and Hemoglobin Following Pioglitazone Treatment is not Hemodilutional in Type II Diabetes Mellitus. <i>Clinical Pharmacology and Therapeutics</i> , 2007, 82, 275-281.	4.7	80
110	Glucagon dose-response curve for hepatic glucose production and glucose disposal in type 2 diabetic patients and normal individuals. <i>Metabolism: Clinical and Experimental</i> , 2002, 51, 1111-1119.	3.4	76
111	The Effect of Pioglitazone on the Liver: Role of adiponectin. <i>Diabetes Care</i> , 2006, 29, 2275-2281.	8.6	76
112	Rosiglitazone and pioglitazone similarly improve insulin sensitivity and secretion, glucose tolerance and adipocytokines in type 2 diabetic patients. <i>Diabetes, Obesity and Metabolism</i> , 2008, 10, 1204-1211.	4.4	76
113	Effect of acute physiological hyperinsulinemia on gene expression in human skeletal muscle in vivo. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2008, 294, E910-E917.	3.5	76
114	Sodium-glucose co-transporter (<scp>SGLT</scp>) and glucose transporter (<scp>GLUT</scp>) expression in the kidney of type 2 diabetic subjects. <i>Diabetes, Obesity and Metabolism</i> , 2017, 19, 1322-1326.	4.4	74
115	Effects of Pioglitazone on Intramyocellular Fat Metabolism in Patients with Type 2 Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010, 95, 1916-1923.	3.6	72
116	Petition to replace current OGTT criteria for diagnosing prediabetes with the 1-hour post-load plasma glucose ≥ 155 mg/dl (8.6 mmol/L). <i>Diabetes Research and Clinical Practice</i> , 2018, 146, 18-33.	2.8	71
117	The relationship between fasting hyperglycemia and insulin secretion in subjects with normal or impaired glucose tolerance. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2008, 295, E401-E406.	3.5	70
118	Empagliflozin and Kinetics of Renal Glucose Transport in Healthy Individuals and Individuals With Type 2 Diabetes. <i>Diabetes</i> , 2017, 66, 1999-2006.	0.6	67
119	Empagliflozin Treatment Is Associated With Improved β -Cell Function in Type 2 Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 1402-1407.	3.6	63
120	Determinants of the increase in ketone concentration during <scp>SGLT2</scp> inhibition in <scp>NGT</scp>, <scp>IFG</scp> and <scp>T2DM</scp> patients. <i>Diabetes, Obesity and Metabolism</i> , 2017, 19, 809-813.	4.4	61
121	Impaired early- but not late-phase insulin secretion in subjects with impaired fasting glucose. <i>Acta Diabetologica</i> , 2011, 48, 209-217.	2.5	55
122	The Relationship Between β -Cell Function and Glycated Hemoglobin: Results from the Veterans Administration Genetic Epidemiology Study. <i>Diabetes Care</i> , 2011, 34, 1006-1010.	8.6	53
123	Mechanisms of Glucose Lowering of Dipeptidyl Peptidase-4 Inhibitor Sitagliptin When Used Alone or With Metformin in Type 2 Diabetes. <i>Diabetes Care</i> , 2013, 36, 2756-2762.	8.6	52
124	Adaptation of Insulin Clearance to Metabolic Demand Is a Key Determinant of Glucose Tolerance. <i>Diabetes</i> , 2021, 70, 377-385.	0.6	47
125	UDP-N-acetylglucosamine transferase and glutamine: fructose 6-phosphate amidotransferase activities in insulin-sensitive tissues. <i>Diabetologia</i> , 1997, 40, 76-81.	6.3	45
126	Physiological hyperinsulinemia impairs insulin-stimulated glycogen synthase activity and glycogen synthesis. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2001, 280, E712-E719.	3.5	45

#	ARTICLE	IF	CITATIONS
127	Effect of Exenatide on Splanchnic and Peripheral Glucose Metabolism in Type 2 Diabetic Subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011, 96, 1763-1770.	3.6	45
128	Pioglitazone Improves Left Ventricular Diastolic Function in Subjects With Diabetes. <i>Diabetes Care</i> , 2017, 40, 1530-1536.	8.6	45
129	Free Fatty Acids Reduce Splanchnic and Peripheral Glucose Uptake in Patients With Type 2 Diabetes. <i>Diabetes</i> , 2002, 51, 3043-3048.	0.6	44
130	Revitalization of pioglitazone: the optimum agent to be combined with a sodium-glucose co-transporter ² inhibitor. <i>Diabetes, Obesity and Metabolism</i> , 2016, 18, 454-462.	4.4	44
131	Endogenous Glucose Production and Hormonal Changes in Response to Canagliflozin and Liraglutide Combination Therapy. <i>Diabetes</i> , 2018, 67, 1182-1189.	0.6	44
132	Is It Time to Change the Type 2 Diabetes Treatment Paradigm? Yes! GLP-1 RAs Should Replace Metformin in the Type 2 Diabetes Algorithm. <i>Diabetes Care</i> , 2017, 40, 1121-1127.	8.6	43
133	Chronic physiologic hyperinsulinemia impairs suppression of plasma free fatty acids and increases de novo lipogenesis but does not cause dyslipidemia in conscious normal rats. <i>Metabolism: Clinical and Experimental</i> , 1999, 48, 330-337.	3.4	41
134	Therapeutic Manipulation of Myocardial Metabolism. <i>Journal of the American College of Cardiology</i> , 2021, 77, 2022-2039.	2.8	40
135	Insulin Resistance the Link between T2DM and CVD: Basic Mechanisms and Clinical Implications. <i>Current Vascular Pharmacology</i> , 2019, 17, 153-163.	1.7	39
136	Reciprocal variations in insulin-stimulated glucose uptake and pancreatic insulin secretion in women with normal glucose tolerance. <i>Journal of the Society for Gynecologic Investigation</i> , 1995, 2, 708-715.	1.7	37
137	Prediabetes and risk of diabetes and associated complications. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2016, 19, 394-399.	2.5	35
138	The Disposition Index Does Not Reflect β -Cell Function in IGT Subjects Treated With Pioglitazone. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, 3774-3781.	3.6	34
139	Combination Therapy With Exenatide Plus Pioglitazone Versus Basal/Bolus Insulin in Patients With Poorly Controlled Type 2 Diabetes on Sulfonylurea Plus Metformin: The Qatar Study. <i>Diabetes Care</i> , 2017, 40, 325-331.	8.6	32
140	Reciprocal Variations in Insulin-Stimulated Glucose Uptake and Pancreatic Insulin Secretion in Women With Normal Glucose Tolerance. <i>Journal of the Society for Gynecologic Investigation</i> , 1995, 2, 708-715.	1.7	31
141	Decreased Non-Insulin-Dependent Glucose Clearance Contributes to the Rise in Fasting Plasma Glucose in the Nondiabetic Range. <i>Diabetes Care</i> , 2008, 31, 311-315.	8.6	30
142	Combination Therapy With Canagliflozin Plus Liraglutide Exerts Additive Effect on Weight Loss, but Not on HbA _{1c} , in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2020, 43, 1234-1241.	8.6	30
143	Durability of Triple Combination Therapy Versus Stepwise Addition Therapy in Patients With New-Onset T2DM: 3-Year Follow-up of EDICT. <i>Diabetes Care</i> , 2021, 44, 433-439.	8.6	29
144	Pioglitazone inhibits mitochondrial pyruvate metabolism and glucose production in hepatocytes. <i>FEBS Journal</i> , 2017, 284, 451-465.	4.7	27

#	ARTICLE	IF	CITATIONS
145	Effect of Chronic Hyperglycemia on Glucose Metabolism in Subjects With Normal Glucose Tolerance. <i>Diabetes</i> , 2018, 67, 2507-2517.	0.6	26
146	Accuracy of 1-Hour Plasma Glucose During the Oral Glucose Tolerance Test in Diagnosis of Type 2 Diabetes in Adults: A Meta-analysis. <i>Diabetes Care</i> , 2021, 44, 1062-1069.	8.6	25
147	Pleiotropic Effects of Thiazolidinediones: Implications for the Treatment of Patients With Type 2 Diabetes Mellitus. <i>Hospital Practice (1995)</i> , 2013, 41, 132-147.	1.0	24
148	Evidence Against an Important Role of Plasma Insulin and Glucagon Concentrations in the Increase in EGP Caused by SGLT2 Inhibitors. <i>Diabetes</i> , 2020, 69, 681-688.	0.6	23
149	Pioglitazone corrects dysregulation of skeletal muscle mitochondrial proteins involved in ATP synthesis in type 2 diabetes. <i>Metabolism: Clinical and Experimental</i> , 2021, 114, 154416.	3.4	23
150	Adiponectin Alleviates Diet-Induced Inflammation in the Liver by Suppressing MCP-1 Expression and Macrophage Infiltration. <i>Diabetes</i> , 2021, 70, 1303-1316.	0.6	22
151	What are the pharmacotherapy options for treating prediabetes?. <i>Expert Opinion on Pharmacotherapy</i> , 2014, 15, 2003-2018.	1.8	21
152	Inhibition of Renal Sodium-Glucose Cotransport With Empagliflozin Lowers Fasting Plasma Glucose and Improves β -Cell Function in Subjects With Impaired Fasting Glucose. <i>Diabetes</i> , 2017, 66, 2495-2502.	0.6	21
153	Mild Physiologic Hyperglycemia Induces Hepatic Insulin Resistance in Healthy Normal Glucose-Tolerant Participants. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 2842-2850.	3.6	18
154	Increase in endogenous glucose production with SGLT2 inhibition is attenuated in individuals who underwent kidney transplantation and bilateral native nephrectomy. <i>Diabetologia</i> , 2020, 63, 2423-2433.	6.3	17
155	Effect of sustained physiologic hyperinsulinaemia and hyperglycaemia on insulin secretion and insulin sensitivity in man. <i>Diabetologia</i> , 1994, 37, 1025-1035.	6.3	17
156	Impaired Suppression of Glucagon in Obese Subjects Parallels Decline in Insulin Sensitivity and Beta-Cell Function. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021, 106, 1398-1409.	3.6	16
157	Insulin Resistance and Hyperinsulinemia: the Egg and the Chicken. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021, 106, 1897-1899.	3.6	16
158	Discordance Between Central (Brain) and Pancreatic Action of Exenatide in Lean and Obese Subjects. <i>Diabetes Care</i> , 2016, 39, 1804-1810.	8.6	15
159	Reduced skeletal muscle phosphocreatine concentration in type 2 diabetic patients: a quantitative image-based phosphorus-31 MR spectroscopy study. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2018, 315, E229-E239.	3.5	15
160	Effect of Mild Physiologic Hyperglycemia on Insulin Secretion, Insulin Clearance, and Insulin Sensitivity in Healthy Glucose-Tolerant Subjects. <i>Diabetes</i> , 2021, 70, 204-213.	0.6	15
161	Combination therapy with pioglitazone/exenatide/metformin reduces the prevalence of hepatic fibrosis and steatosis: The efficacy and durability of initial combination therapy for type 2 diabetes (<sc>EDICT</sc>). <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 899-907.	4.4	15
162	Insulin secretion, insulin resistance, and obesity. , 1982, 6 Suppl 1, 73-82.		15

#	ARTICLE	IF	CITATIONS
163	The tumor suppressor TMEM127 regulates insulin sensitivity in a tissue-specific manner. <i>Nature Communications</i> , 2019, 10, 4720.	12.8	14
164	Glucose-Mediated Glucose Disposal at Baseline Insulin Is Impaired in IFG. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 163-171.	3.6	14
165	Comprehensive assessment of postischemic vascular reactivity in Hispanic children and adults with and without diabetes mellitus. <i>Pediatric Diabetes</i> , 2006, 7, 329-335.	2.9	13
166	Efficacy of lower doses of pioglitazone after stroke or transient ischaemic attack in patients with insulin resistance. <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 1150-1158.	4.4	13
167	Differential effect of hyperglycemia and hyperinsulinemia on pathways of hepatic glycogen repletion. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 1991, 260, E731-E735.	3.5	12
168	Efficacy of Exenatide Plus Pioglitazone Vs Basal/Bolus Insulin in T2DM Patients With Very High HbA1c. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, 2162-2170.	3.6	12
169	Slope of change in HbA _{1c} from baseline with empagliflozin compared with sitagliptin or glimepiride in patients with type 2 diabetes. <i>Endocrinology, Diabetes and Metabolism</i> , 2018, 1, e00016.	2.4	12
170	Improved Beta Cell Glucose Sensitivity Plays Predominant Role in the Decrease in HbA _{1c} with Cana and Lira in T2DM. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, 3226-3233.	3.6	10
171	Strong Association Between Insulin-Mediated Glucose Uptake and the 2-Hour, Not the Fasting Plasma Glucose Concentration, in the Normal Glucose Tolerance Range. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, 3444-3449.	3.6	9
172	Diverted sleeve gastrectomy with ileal transposition in overweight, obese, and morbidly obese patients with type 2 diabetes: results of 1-year follow-up. <i>Surgery for Obesity and Related Diseases</i> , 2016, 12, 541-549.	1.2	9
173	Glucose lowering and vascular protective effects of cycloset added to GLP-1 receptor agonists in patients with type 2 diabetes. <i>Endocrinology, Diabetes and Metabolism</i> , 2018, 1, e00034.	2.4	9
174	Modifying chronic kidney disease progression with the mineralocorticoid receptor antagonist finerenone in patients with type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 1197-1205.	4.4	9
175	Effect of vildagliptin add-on treatment to metformin on plasma asymmetric dimethylarginine in type 2 diabetes mellitus patients. <i>Drug Design, Development and Therapy</i> , 2014, 8, 239.	4.3	7
176	Insulin secretion predicts the response to therapy with exenatide plus pioglitazone, but not to basal/bolus insulin in poorly controlled T2DM patients: Results from the Qatar study. <i>Diabetes, Obesity and Metabolism</i> , 2018, 20, 1075-1079.	4.4	7
177	Combination therapy with pioglitazone/exenatide improves beta cell function and produces superior glycaemic control compared with basal/bolus insulin in poorly controlled type 2 diabetes: A 3-year follow-up of the Qatar study. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 2287-2294.	4.4	7
178	Impaired left ventricular diastolic function in T2DM patients is closely related to glycemic control. <i>Endocrinology, Diabetes and Metabolism</i> , 2018, 1, e00014.	2.4	6
179	Insulin resistance, hypertension and cellular ion transport systems. <i>Acta Diabetologica</i> , 1992, 29, 196-200.	2.5	5
180	Impact of ethnicity and obesity on insulin resistance in two ethnic groups at very high risk of type 2 diabetes. <i>Diabetes and Metabolism</i> , 2017, 43, 292-294.	2.9	5

#	ARTICLE	IF	CITATIONS
181	Hormoneâ€‘substrate changes with exenatide plus dapagliflozin versus each drug alone: The randomized, activeâ€‘controlled DURATIONâ€‘8 study. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 99-106.	4.4	5
182	Glycated hemoglobin versus oral glucose tolerance test in the identification of subjects with prediabetes in Qatari population. <i>BMC Endocrine Disorders</i> , 2019, 19, 87.	2.2	4
183	Pioglitazone and cardiovascular risk in T2DM patients: is it good for all?. <i>Annals of Translational Medicine</i> , 2018, 6, 192-192.	1.7	4
184	Dapagliflozin Impairs the Suppression of Endogenous Glucose Production in Type 2 Diabetes Following Oral Glucose. <i>Diabetes Care</i> , 2022, 45, 1372-1380.	8.6	4
185	Effects of Sustained Hyperglycemia on Skeletal Muscle Lipids in Healthy Subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, e3177-e3185.	3.6	4
186	FGF21 contributes to metabolic improvements elicited by combination therapy with exenatide and pioglitazone in patients with type 2 diabetes. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2022, 323, E123-E132.	3.5	4
187	Fiber orientation measurements by diffusion tensor imaging improve hydrogen-1 magnetic resonance spectroscopy of intramyocellular lipids in human leg muscles. <i>Journal of Medical Imaging</i> , 2015, 2, 026002.	1.5	3
188	Exenatide modulates visual cortex responses. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3167.	4.0	3
189	Pioglitazone prevents the increase in plasma ketone concentration associated with dapagliflozin in insulinâ€‘treated T2DM patients: Results from the Qatar Study. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 705-709.	4.4	3
190	Clinical Parameters, Fuel Oxidation, and Glucose Kinetics in Patients With Type 2 Diabetes Treated With Dapagliflozin Plus Saxagliptin. <i>Diabetes Care</i> , 2020, 43, 2519-2527.	8.6	3
191	Insulin Secretion Predicts the Response to Antidiabetic Therapy in Patients With New-onset Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021, 106, 3497-3504.	3.6	3
192	Therapeutic strategies for type 2 diabetes mellitus patients with very high HbA1c: is insulin the only option?. <i>Annals of Translational Medicine</i> , 2018, 6, S95-S95.	1.7	3
193	<sc>Type 2 diabetes</sc> subgroups and response to glucoseâ€‘lowering therapy: Results from the <sc>EDICT</sc> and Qatar studies. <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 1810-1818.	4.4	3
194	Insulin secretion is a strong predictor for need of insulin therapy in patients with newâ€‘onset diabetes and <sc>HbA1c of more than 10%: A</sc> post hoc analysis of the <sc>EDICT</sc> study. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 1631-1639.	4.4	2
195	Personalized approach for type 2 diabetes pharmacotherapy: where are we and where do we need to be?. <i>Expert Opinion on Pharmacotherapy</i> , 2021, 22, 1-13.	1.8	2
196	Sodiumâ€‘Glucose Cotransporter 2 Inhibitors and the Kidney. <i>Diabetes Spectrum</i> , 2021, 34, 225-234.	1.0	1
197	Preface: Cardiorenal Considerations for Type 2 Diabetesâ€‘Time to Exit the Dark Ages. <i>Diabetes Spectrum</i> , 2021, 34, 214-215.	1.0	0
198	Comprehensive analysis of circulating miRNA expression profiles in insulin resistance and type 2 diabetes in Qatari population. <i>International Journal of Transgender Health</i> , 2022, 15, 191-202.	2.3	0

#	ARTICLE	IF	CITATIONS
199	Comparison of the effects of human recombinant insulin-like growth factor I and insulin on plasma amino acid concentrations and leucine kinetics in humans. <i>Diabetologia</i> , 1995, 38, 732-738.	6.3	0