

# Tetsuji Sawada

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9035470/publications.pdf>

Version: 2024-02-01

17  
papers

1,648  
citations

840585  
11  
h-index

996849  
15  
g-index

18  
all docs

18  
docs citations

18  
times ranked

2001  
citing authors

#	ARTICLE	IF	CITATIONS
1	Detection of salivary citrullinated cytokeratin 13 in healthy individuals and patients with rheumatoid arthritis by proteomics analysis. PLoS ONE, 2022, 17, e0265687.	1.1	3
2	Distal Interphalangeal Joint Involvement May Be Associated with Disease Activity and Affected Joint Distribution in Rheumatoid Arthritis. Journal of Clinical Medicine, 2022, 11, 1405.	1.0	1
3	Influence of Concomitant Methotrexate use on the Clinical Effectiveness, Retention, and Safety of Abatacept in Biologic-Naïve Patients with Rheumatoid Arthritis: Post-hoc Subgroup Analysis of ORIGAMI Study. Modern Rheumatology, 2022, , .	0.9	0
4	Clinical variables, including novel joint index, associated with future patient-physician discordance in global assessment of rheumatoid arthritis (<sc>RA</sc>) disease activity based on a large <sc>RA</sc> database in Japan. International Journal of Rheumatic Diseases, 2022, 25, 1020-1028.	0.9	2
5	Recommendations for the Management of Neuro-Behçet's Disease by the Japanese National Research Committee for Behçet's Disease. Internal Medicine, 2020, 59, 2359-2367.	0.3	11
6	Serum phosphatidylserine-specific phospholipase A 1 as a novel biomarker for monitoring systemic lupus erythematosus disease activity. International Journal of Rheumatic Diseases, 2019, 22, 2059-2066.	0.9	19
7	Influence of seasonal changes on disease activity and distribution of affected joints in rheumatoid arthritis. BMC Musculoskeletal Disorders, 2019, 20, 30.	0.8	23
8	Systemic lupus erythematosus and immunodeficiency. Immunological Medicine, 2019, 42, 1-9.	1.4	45
9	Influence of large joint involvement on patient-physician discordance in global assessment of rheumatoid arthritis disease activity analyzed by a novel joint index. International Journal of Rheumatic Diseases, 2018, 21, 1237-1245.	0.9	8
10	The age at onset of rheumatoid arthritis is increasing in Japan: a nationwide database study. International Journal of Rheumatic Diseases, 2017, 20, 839-845.	0.9	87
11	Head-to-head comparison of the safety of tocilizumab and tumor necrosis factor inhibitors in rheumatoid arthritis patients (RA) in clinical practice: results from the registry of Japanese RA patients on biologics for long-term safety (REAL) registry. Arthritis Research and Therapy, 2015, 17, 74.	1.6	53
12	Retrospective analysis of long-term outcome of chronic progressive neurological manifestations in Behçet's disease. Journal of the Neurological Sciences, 2015, 349, 143-148.	0.3	32
13	Identification of citrullinated cellular fibronectin in synovial fluid from patients with rheumatoid arthritis. Modern Rheumatology, 2014, 24, 766-769.	0.9	16
14	Clinical characteristics of neuro-Behçet's disease in Japan: a multicenter retrospective analysis. Modern Rheumatology, 2012, 22, 405-413.	0.9	73
15	Incidence and Risk Factors for Serious Infection in Patients with Rheumatoid Arthritis Treated with Tumor Necrosis Factor Inhibitors: A Report from the Registry of Japanese Rheumatoid Arthritis Patients for Longterm Safety. Journal of Rheumatology, 2011, 38, 1258-1264.	1.0	141
16	A novel enzyme immunoassay for the determination of phosphatidylserine-specific phospholipase A1 in human serum samples. Clinica Chimica Acta, 2010, 411, 1090-1094.	0.5	23
17	Functional haplotypes of PADI4, encoding citrullinating enzyme peptidylarginine deiminase 4, are associated with rheumatoid arthritis. Nature Genetics, 2003, 34, 395-402.	9.4	1,111