

Mark P Simmons

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9046529/publications.pdf>

Version: 2024-02-01

114
papers

7,997
citations

81900

39
h-index

49909

87
g-index

114
all docs

114
docs citations

114
times ranked

7896
citing authors

#	ARTICLE	IF	CITATIONS
1	Gene-tree misrooting drives conflicts in phylogenomic coalescent analyses of palaeognath birds. <i>Molecular Phylogenetics and Evolution</i> , 2022, 167, 107344.	2.7	16
2	Benefits of alignment quality control processing steps and an Angiosperms353 phylogenomics pipeline applied to the Celastrales. <i>Cladistics</i> , 2022, 38, 595-611.	3.3	1
3	Collapsing dubiously resolved gene-tree branches in phylogenomic coalescent analyses. <i>Molecular Phylogenetics and Evolution</i> , 2021, 158, 107092.	2.7	24
4	Divergence and support among slightly suboptimal likelihood gene trees. <i>Cladistics</i> , 2020, 36, 322-340.	3.3	20
5	ILS-Aware Analysis of Low-Homoplasmy Retroelement Insertions: Inference of Species Trees and Introgression Using Quartets. <i>Journal of Heredity</i> , 2020, 111, 147-168.	2.4	30
6	Adenine-cytosine substitutions are an alternative pathway of compensatory mutation in angiosperm ITS2. <i>Rna</i> , 2020, 26, 209-217.	3.5	7
7	Thistle be a mess: Untangling the taxonomy of <i>Cirsium</i> (Cardueae: Compositae) in North America. <i>Journal of Systematics and Evolution</i> , 2020, 58, 881-912.	3.1	10
8	New Guinea has the world's richest island flora. <i>Nature</i> , 2020, 584, 579-583.	27.8	108
9	Clonal Diversity, Cultivar Traits, Geographic Dispersal, and the Ethnotaxonomy of Cultivated Qat (<i>Catha edulis</i> , Celastraceae). <i>Economic Botany</i> , 2020, 74, 273-291.	1.7	0
10	Mitochondrial fatty acid oxidation is required for storage lipid catabolism in a marine diatom. <i>New Phytologist</i> , 2020, 228, 946-958.	7.3	25
11	Partitioned coalescence support reveals biases in species-tree methods and detects gene trees that determine phylogenomic conflicts. <i>Molecular Phylogenetics and Evolution</i> , 2019, 139, 106539.	2.7	31
12	Alternative analyses of compensatory base changes in an ITS2 phylogeny of <i>Corydalis</i> (Papaveraceae). <i>Annals of Botany</i> , 2019, 124, 233-243.	2.9	9
13	Gene-wise resampling outperforms site-wise resampling in phylogenetic coalescence analyses. <i>Molecular Phylogenetics and Evolution</i> , 2019, 131, 80-92.	2.7	33
14	Transcriptome-wide comparison of selenium hyperaccumulator and nonaccumulator <i>Stanleya</i> species provides new insight into key processes mediating the hyperaccumulation syndrome. <i>Plant Biotechnology Journal</i> , 2018, 16, 1582-1594.	8.3	67
15	Narrow thermal tolerance and low dispersal drive higher speciation in tropical mountains. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, 12471-12476.	7.1	161
16	Ten Rules for Associate Editors. <i>Botanical Review</i> , The, 2018, 84, 99-107.	3.9	0
17	Relative benefits of amino acid, codon, degeneracy, DNA, and purine-pyrimidine character coding for phylogenetic analyses of exons. <i>Journal of Systematics and Evolution</i> , 2017, 55, 85-109.	3.1	24
18	Employing Two-stage Derivatisation and GC-MS to Assay for Cathine and Related Stimulant Alkaloids across the Celastraceae. <i>Phytochemical Analysis</i> , 2017, 28, 257-266.	2.4	7

#	ARTICLE	IF	CITATIONS
19	Phylogeography of the wild and cultivated stimulant plant qat (<i>Catha edulis</i> , Celastraceae) in areas of historical cultivation. <i>American Journal of Botany</i> , 2017, 104, 538-549.	1.7	9
20	Mutually exclusive phylogenomic inferences at the root of the angiosperms: <i>Amborella</i> is supported as sister and Observed Variability is biased. <i>Cladistics</i> , 2017, 33, 488-512.	3.3	16
21	Resolution of a concatenation/coalescence keruffle: partitioned coalescence support and a robust family-level tree for Mammalia. <i>Cladistics</i> , 2017, 33, 295-332.	3.3	70
22	Systematics of New World <i>Maytenus</i> (Celastraceae) and a New Delimitation of the Genus. <i>Systematic Botany</i> , 2017, 42, 680-693.	0.5	23
23	The Complete Plastid Genome of <i>Lagerstroemia fauriei</i> and Loss of rpl2 Intron from <i>Lagerstroemia</i> (Lythraceae). <i>PLoS ONE</i> , 2016, 11, e0150752.	2.5	52
24	<i>Pleurostylia serrulata</i> and Two Allied New Species from Africa are Actually Members of the New World <i>Crossopetalum</i> (Celastraceae). <i>Systematic Botany</i> , 2016, 41, 851-864.	0.5	2
25	Biases of tree-independent-character-subsampling methods. <i>Molecular Phylogenetics and Evolution</i> , 2016, 100, 424-443.	2.7	11
26	Co-variation in methanotroph community composition and activity in three temperate grassland soils. <i>Soil Biology and Biochemistry</i> , 2016, 95, 78-86.	8.8	22
27	Linear Plasmids and the Rate of Sequence Evolution in Plant Mitochondrial Genomes. <i>Genome Biology and Evolution</i> , 2016, 8, 364-374.	2.5	29
28	The effects of subsampling gene trees on coalescent methods applied to ancient divergences. <i>Molecular Phylogenetics and Evolution</i> , 2016, 97, 76-89.	2.7	45
29	Biogeography of the Malagasy Celastraceae: Multiple independent origins followed by widespread dispersal of genera from Madagascar. <i>Molecular Phylogenetics and Evolution</i> , 2016, 94, 365-382.	2.7	27
30	Ten Simple Rules for Writing a Reply Paper. <i>PLoS Computational Biology</i> , 2015, 11, e1004536.	3.2	1
31	Coalescence vs. concatenation: Sophisticated analyses vs. first principles applied to rooting the angiosperms. <i>Molecular Phylogenetics and Evolution</i> , 2015, 91, 98-122.	2.7	77
32	Evolution of selenium hyperaccumulation in <i>Stanleya</i> (Brassicaceae) as inferred from phylogeny, physiology and X-ray microprobe analysis. <i>New Phytologist</i> , 2015, 205, 583-595.	7.3	37
33	Characterization of selenium and sulfur accumulation across the genus <i>Stanleya</i> (Brassicaceae): A field survey and common-garden experiment. <i>American Journal of Botany</i> , 2014, 101, 830-839.	1.7	55
34	Divergent maximum-likelihood-branch-support values for polytomies. <i>Molecular Phylogenetics and Evolution</i> , 2014, 73, 87-96.	2.7	49
35	Limitations of locally sampled characters in phylogenetic analyses of sparse supermatrices. <i>Molecular Phylogenetics and Evolution</i> , 2014, 74, 1-14.	2.7	15
36	Molecular phylogeny and revised classification of <i>Eucosma</i> and related genera (Lepidoptera: Tortricidae: Eucosmini). <i>Systematic Entomology</i> , 2014, 39, 49-67.	3.9	14

#	ARTICLE	IF	CITATIONS
37	Haydenoxylon, a Replacement Name for Haydenia (Celastraceae). <i>Novon</i> , 2014, 23, 224-225.	0.3	2
38	A confounding effect of missing data on character conflict in maximum likelihood and Bayesian MCMC phylogenetic analyses. <i>Molecular Phylogenetics and Evolution</i> , 2014, 80, 267-280.	2.7	53
39	Bias in Tree Searches and its Consequences for Measuring Group Supports. <i>Systematic Biology</i> , 2014, 63, 851-861.	5.6	9
40	Disparate parametric branch-support values from ambiguous characters. <i>Molecular Phylogenetics and Evolution</i> , 2014, 78, 66-86.	2.7	18
41	A New Species of <i>Maytenus</i> (Celastraceae) with Fleshy Fruits from Eastern Brazil, with Notes on the Delimitation of <i>Maytenus</i> . <i>Systematic Botany</i> , 2014, 39, 478-484.	0.5	10
42	Dubious resolution and support from published sparse supermatrices: The importance of thorough tree searches. <i>Molecular Phylogenetics and Evolution</i> , 2014, 78, 334-348.	2.7	26
43	Quantification and relative severity of inflated branch-support values generated by alternative methods: An empirical example. <i>Molecular Phylogenetics and Evolution</i> , 2013, 67, 277-296.	2.7	27
44	Evaluation of microsatellites of <i>Catha edulis</i> (Celastraceae) identified using pyrosequencing. <i>Biochemical Systematics and Ecology</i> , 2013, 49, 1-9.	1.3	17
45	Phylogeny of the Asian <i>Hedyotis</i> - <i>Oldenlandia</i> complex (Spermacoceae, Rubiaceae): Evidence for high levels of polyphyly and the parallel evolution of diplophragmous capsules. <i>Molecular Phylogenetics and Evolution</i> , 2013, 67, 110-122.	2.7	42
46	<i>Wilczekra</i> , a New Genus of Celastraceae for a Disjunct Lineage of <i>Euonymus</i> . <i>Systematic Botany</i> , 2013, 38, 148-153.	0.5	6
47	GEOGRAPHIC AND TAXONOMIC DISPARITIES IN SPECIES DIVERSITY: DISPERSAL AND DIVERSIFICATION RATES ACROSS WALLACE'S LINE. <i>Evolution; International Journal of Organic Evolution</i> , 2013, 67, 2058-2071.	2.3	42
48	An artifact caused by undersampling optimal trees in supermatrix analyses of locally sampled characters. <i>Molecular Phylogenetics and Evolution</i> , 2013, 69, 265-275.	2.7	26
49	Miocene Dispersal Drives Island Radiations in the Palm Tribe Trachycarpeae (Arecaceae). <i>Systematic Biology</i> , 2012, 61, 426-442.	5.6	77
50	Phylogeny of Celastraceae tribe Euomyneae inferred from morphological characters and nuclear and plastid genes. <i>Molecular Phylogenetics and Evolution</i> , 2012, 62, 9-20.	2.7	33
51	Radical instability and spurious branch support by likelihood when applied to matrices with non-random distributions of missing data. <i>Molecular Phylogenetics and Evolution</i> , 2012, 62, 472-484.	2.7	75
52	Evaluating multiple criteria for species delimitation: an empirical example using Hawaiian palms (Arecaceae: Pritchardia). <i>BMC Evolutionary Biology</i> , 2012, 12, 23.	3.2	42
53	Rapid Intraspecific Evolution of miRNA and siRNA Genes in the Mosquito <i>Aedes aegypti</i> . <i>PLoS ONE</i> , 2012, 7, e44198.	2.5	52
54	Phylogeny of Celastraceae Subfamilies Cassinoideae and Tripterygioideae Inferred from Morphological Characters and Nuclear and Plastid Loci. <i>Systematic Botany</i> , 2012, 37, .	0.5	21

#	ARTICLE	IF	CITATIONS
55	Misleading results of likelihood-based phylogenetic analyses in the presence of missing data. <i>Cladistics</i> , 2012, 28, 208-222.	3.3	90
56	Delimitation of the Segregate Genera of <i>Maytenus</i> s. l. (Celastraceae) Based on Morphological and Molecular Characters. <i>Systematic Botany</i> , 2011, 36, 922-932.	0.5	38
57	Phylogenetic inference using non-redundant coding of dependent characters versus alternative approaches for protein-coding genes. <i>Cladistics</i> , 2011, 27, 186-196.	3.3	2
58	The deterministic effects of alignment bias in phylogenetic inference. <i>Cladistics</i> , 2011, 27, 402-416.	3.3	10
59	Application of DNA barcodes in <i>Hedyotis</i> L. (Spermacoceae, Rubiaceae). <i>Journal of Systematics and Evolution</i> , 2011, 49, 203-212.	3.1	31
60	Molecular analyses of the Chinese herb Leigongteng (<i>Tripterygium wilfordii</i> Hook.f.). <i>Phytochemistry</i> , 2011, 72, 21-26.	2.9	74
61	Phylogeny of Celastraceae subfamily Hippocrateoideae inferred from morphological characters and nuclear and plastid loci. <i>Molecular Phylogenetics and Evolution</i> , 2011, 59, 320-330.	2.7	23
62	Spurious 99% bootstrap and jackknife support for unsupported clades. <i>Molecular Phylogenetics and Evolution</i> , 2011, 61, 177-191.	2.7	76
63	Development of microsatellites in the Hawaiian endemic palm <i>Pritchardia martii</i> (Arecaceae) and their utility in congeners. <i>American Journal of Botany</i> , 2011, 98, e139-e140.	1.7	3
64	Alignment of, and phylogenetic inference from, random sequences: The susceptibility of alternative alignment methods to creating artifactual resolution and support. <i>Molecular Phylogenetics and Evolution</i> , 2010, 57, 1004-1016.	2.7	21
65	Phylogeny of Celastraceae Subfamily Salacioideae and Tribe Lophopetaleae Inferred from Morphological Characters and Nuclear and Plastid Genes. <i>Systematic Botany</i> , 2010, 35, 358-367.	0.5	25
66	Contribution of molecular cladistics to the taxonomy of Rutaceae in China*. <i>Journal of Systematics and Evolution</i> , 2009, 47, 144-150.	3.1	5
67	Phylogeny of the Celastreae (Celastraceae) and the relationships of <i>Catha edulis</i> (qat) inferred from morphological characters and nuclear and plastid genes. <i>Molecular Phylogenetics and Evolution</i> , 2008, 48, 745-757.	2.7	59
68	Potential use of host-derived genome signatures to root virus phylogenies. <i>Molecular Phylogenetics and Evolution</i> , 2008, 49, 969-978.	2.7	8
69	Incorporation of gap characters and lineage-specific regions into phylogenetic analyses of gene families from divergent clades: an example from the kinesin superfamily across eukaryotes. <i>Cladistics</i> , 2008, 24, 372-384.	3.3	7
70	The relative sensitivity of different alignment methods and character codings in sensitivity analysis. <i>Cladistics</i> , 2008, 24, 1039-1050.	3.3	7
71	Congruence of Molecular, Morphological, and Biochemical Profiles in Rutaceae: a Cladistic Analysis of the Subfamilies Rutoideae and Toddalioidae. <i>Systematic Botany</i> , 2007, 32, 837-846.	0.5	43
72	A penalty of using anonymous dominant markers (AFLPs, ISSRs, and RAPDs) for phylogenetic inference. <i>Molecular Phylogenetics and Evolution</i> , 2007, 42, 528-542.	2.7	51

#	ARTICLE	IF	CITATIONS
73	A phylogeny of Anisophylleaceae based on six nuclear and plastid loci: Ancient disjunctions and recent dispersal between South America, Africa, and Asia. <i>Molecular Phylogenetics and Evolution</i> , 2007, 44, 1057-1067.	2.7	21
74	The relative performance of indel-coding methods in simulations. <i>Molecular Phylogenetics and Evolution</i> , 2007, 44, 724-740.	2.7	88
75	Are mitochondrial genes useful for the analysis of monocot relationships?. <i>Taxon</i> , 2006, 55, 857-870.	0.7	37
76	Functional trait niches of North American lotic insects: traits-based ecological applications in light of phylogenetic relationships. <i>Journal of the North American Benthological Society</i> , 2006, 25, 730-755.	3.1	704
77	A Thorny Dilemma: Testing Alternative Intrageneric Classifications within <i>Ziziphus</i> (Rhamnaceae). <i>Systematic Botany</i> , 2006, 31, 826-842.	0.5	55
78	Phylogeny of the <i>Elaeodendron</i> Group (Celastraceae) Inferred from Morphological Characters and Nuclear and Plastid Genes. <i>Systematic Botany</i> , 2006, 31, 512-524.	0.5	20
79	The relative performance of Bayesian and parsimony approaches when sampling characters evolving under homogeneous and heterogeneous sets of parameters. <i>Cladistics</i> , 2006, 22, 171-185.	3.3	17
80	Quantification of the success of phylogenetic inference in simulations. <i>Cladistics</i> , 2006, 22, 249-255.	3.3	26
81	Phylogeny of the Cucurbitales based on DNA sequences of nine loci from three genomes: Implications for morphological and sexual system evolution. <i>Molecular Phylogenetics and Evolution</i> , 2006, 39, 305-322.	2.7	121
82	Comprehensive comparative analysis of kinesins in photosynthetic eukaryotes. <i>BMC Genomics</i> , 2006, 7, 18.	2.8	102
83	Phylogeny and Delimitation of the Celastrales Inferred from Nuclear and Plastid Genes. <i>Systematic Botany</i> , 2006, 31, 122-137.	0.5	99
84	How Can Third Codon Positions Outperform First and Second Codon Positions in Phylogenetic Inference? An Empirical Example from the Seed Plants. <i>Systematic Biology</i> , 2006, 55, 245-258.	5.6	32
85	Origin and Evolution of Kinesin-Like Calmodulin-Binding Protein. <i>Plant Physiology</i> , 2005, 138, 1711-1722.	4.8	34
86	Character-state space versus rate of evolution in phylogenetic inference. <i>Cladistics</i> , 2004, 20, 191-204.	3.3	33
87	Efficiently resolving the basal clades of a phylogenetic tree using Bayesian and parsimony approaches: a case study using mitogenomic data from 100 higher teleost fishes. <i>Molecular Phylogenetics and Evolution</i> , 2004, 31, 351-362.	2.7	49
88	Independence of alignment and tree search. <i>Molecular Phylogenetics and Evolution</i> , 2004, 31, 874-879.	2.7	272
89	Relative character-state space, amount of potential phylogenetic information, and heterogeneity of nucleotide and amino acid characters*1. <i>Molecular Phylogenetics and Evolution</i> , 2004, 32, 913-913.	2.7	0
90	How Meaningful Are Bayesian Support Values?. <i>Molecular Biology and Evolution</i> , 2004, 21, 188-199.	8.9	326

#	ARTICLE	IF	CITATIONS
91	A Phylogeny of the Monocots, as Inferred from rbcL and atpA Sequence Variation, and a Comparison of Methods for Calculating Jackknife and Bootstrap Values. <i>Systematic Botany</i> , 2004, 29, 467-510.	0.5	173
92	Relative character-state space, amount of potential phylogenetic information, and heterogeneity of nucleotide and amino acid characters. <i>Molecular Phylogenetics and Evolution</i> , 2004, 32, 913-926.	2.7	39
93	The effects of increasing genetic distance on alignment of, and tree construction from, rDNA internal transcribed spacer sequences. <i>Molecular Phylogenetics and Evolution</i> , 2003, 26, 444-451.	2.7	42
94	From basepairs to birdsongs: phylogenetic data in the age of genomics. <i>Cladistics</i> , 2003, 19, 333-347.	3.3	38
95	Whole-Genome Comparison of Leucine-Rich Repeat Extensins in Arabidopsis and Rice. A Conserved Family of Cell Wall Proteins Form a Vegetative and a Reproductive Clade. <i>Plant Physiology</i> , 2003, 131, 1313-1326.	4.8	128
96	From basepairs to birdsongs: phylogenetic data in the age of genomics. <i>Cladistics</i> , 2003, 19, 333-347.	3.3	3
97	Limitations of Relative Apparent Synapomorphy Analysis (RASA) for Measuring Phylogenetic Signal. <i>Molecular Biology and Evolution</i> , 2002, 19, 14-23.	8.9	20
98	Uninode coding vs gene tree parsimony for phylogenetic reconstruction using duplicate genes. <i>Molecular Phylogenetics and Evolution</i> , 2002, 23, 481-498.	2.7	39
99	Amino acid vs. nucleotide characters: challenging preconceived notions. <i>Molecular Phylogenetics and Evolution</i> , 2002, 24, 78-90.	2.7	55
100	Conflict between Amino Acid and Nucleotide Characters. <i>Cladistics</i> , 2002, 18, 200-206.	3.3	27
101	Artifacts of Coding Amino Acids and Other Composite Characters for Phylogenetic Analysis. <i>Cladistics</i> , 2002, 18, 354-365.	3.3	40
102	Artifacts of Coding Amino Acids and Other Composite Characters for Phylogenetic Analysis. <i>Cladistics</i> , 2002, 18, 354-365.	3.3	11
103	Phylogeny of the Celastraceae Inferred from 26S Nuclear Ribosomal DNA, Phytochrome B, rbcL, atpB, and Morphology. <i>Molecular Phylogenetics and Evolution</i> , 2001, 19, 353-366.	2.7	89
104	Phylogeny of the Celastraceae inferred from phytochrome B gene sequence and morphology. <i>American Journal of Botany</i> , 2001, 88, 313-325.	1.7	75
105	Incorporation, Relative Homoplasy, and Effect of Gap Characters in Sequence-Based Phylogenetic Analyses. <i>Systematic Biology</i> , 2001, 50, 454-462.	5.6	138
106	Incorporation, Relative Homoplasy, and Effect of Gap Characters in Sequence-Based Phylogenetic Analyses. <i>Systematic Biology</i> , 2001, 50, 454-462.	5.6	166
107	Phylogeny Reconstruction Using Duplicate Genes. <i>Molecular Biology and Evolution</i> , 2000, 17, 469-473.	8.9	30
108	A Fundamental Problem with Amino-Acid-Sequence Characters for Phylogenetic Analyses. <i>Cladistics</i> , 2000, 16, 274-282.	3.3	33

#	ARTICLE	IF	CITATIONS
109	Gaps as Characters in Sequence-Based Phylogenetic Analyses. <i>Systematic Biology</i> , 2000, 49, 369-381.	5.6	2,312
110	A Fundamental Problem with Amino-Acid-Sequence Characters for Phylogenetic Analyses. <i>Cladistics</i> , 2000, 16, 274-282.	3.3	9
111	Relationships and Morphological Character Change among Genera of Celastraceae Sensu Lato (Including Hippocrateaceae). <i>Annals of the Missouri Botanical Garden</i> , 1999, 86, 723.	1.3	50
112	Data Decisiveness, Data Quality, and Incongruence in Phylogenetic Analysis: An Example From the Monocotyledons Using Mitochondrial atp A Sequences. <i>Systematic Biology</i> , 1998, 47, 282-310.	5.6	157
113	Revision of the Cerrado Hemicryptophytic Chamaesyce of Boissier's "Pleiadeniae" (Euphorbiaceae). <i>Brittonia</i> , 1997, 49, 155.	0.2	6
114	Wood Anatomy of Amanoa (Euphorbiaceae). <i>IAWA Journal</i> , 1993, 14, 205-213.	2.7	3