

Yiguo Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9053129/publications.pdf>

Version: 2024-02-01

59
papers

2,262
citations

201674

27
h-index

243625

44
g-index

78
all docs

78
docs citations

78
times ranked

2018
citing authors

#	ARTICLE	IF	CITATIONS
1	Involvement of the Acid Sphingomyelinase Pathway in UVA-induced Apoptosis. <i>Journal of Biological Chemistry</i> , 2001, 276, 11775-11782.	3.4	134
2	Molecular and cellular basis for the unique functioning of Nrf1, an indispensable transcription factor for maintaining cell homeostasis and organ integrity. <i>Biochemical Journal</i> , 2016, 473, 961-1000.	3.7	117
3	Negative regulation of the Nrf1 transcription factor by its N-terminal domain is independent of Keap1: Nrf1, but not Nrf2, is targeted to the endoplasmic reticulum. <i>Biochemical Journal</i> , 2006, 399, 373-385.	3.7	112
4	The NHB1 (N-terminal homology box 1) sequence in transcription factor Nrf1 is required to anchor it to the endoplasmic reticulum and also to enable its asparagine-glycosylation. <i>Biochemical Journal</i> , 2007, 408, 161-172.	3.7	94
5	Involvement of c-jun NH2-terminal kinases in resveratrol-induced activation of p53 and apoptosis. <i>Molecular Carcinogenesis</i> , 2002, 33, 244-250.	2.7	91
6	Signal Transduction Pathways Involved in Phosphorylation and Activation of p70S6K Following Exposure to UVA Irradiation. <i>Journal of Biological Chemistry</i> , 2001, 276, 20913-20923.	3.4	86
7	Requirement of ATM in UVA-induced Signaling and Apoptosis. <i>Journal of Biological Chemistry</i> , 2002, 277, 3124-3131.	3.4	79
8	UVA Induces Ser381 Phosphorylation of p90RSK/MAPKAP-K1 via ERK and JNK Pathways. <i>Journal of Biological Chemistry</i> , 2001, 276, 14572-14580.	3.4	77
9	MSK1 and JNKs Mediate Phosphorylation of STAT3 in UVA-irradiated Mouse Epidermal JB6 Cells. <i>Journal of Biological Chemistry</i> , 2001, 276, 42534-42542.	3.4	72
10	The Nrf1 CNC/bZIP protein is a nuclear envelope-bound transcription factor that is activated by t-butyl hydroquinone but not by endoplasmic reticulum stressors. <i>Biochemical Journal</i> , 2009, 418, 293-310.	3.7	69
11	Phosphorylation of 4E-BP1 Is Mediated by the p38/MSK1 Pathway in Response to UVB Irradiation. <i>Journal of Biological Chemistry</i> , 2002, 277, 8810-8816.	3.4	68
12	The Nrf3 Transcription Factor Is a Membrane-bound Glycoprotein Targeted to the Endoplasmic Reticulum through Its N-terminal Homology Box 1 Sequence. <i>Journal of Biological Chemistry</i> , 2009, 284, 3195-3210.	3.4	65
13	Nrf1 and Nrf2 Transcription Factors Regulate Androgen Receptor Transactivation in Prostate Cancer Cells. <i>PLoS ONE</i> , 2014, 9, e87204.	2.5	59
14	MAP Kinases Mediate UVB-induced Phosphorylation of Histone H3 at Serine 28. <i>Journal of Biological Chemistry</i> , 2001, 276, 12932-12937.	3.4	57
15	Identification of topological determinants in the N-terminal domain of transcription factor Nrf1 that control its orientation in the endoplasmic reticulum membrane. <i>Biochemical Journal</i> , 2010, 430, 497-510.	3.7	52
16	Transcription Factor Nrf1 Is Topologically Repartitioned across Membranes to Enable Target Gene Transactivation through Its Acidic Glucose-Responsive Domains. <i>PLoS ONE</i> , 2014, 9, e93458.	2.5	49
17	Arsenite-induced Phosphorylation of Histone H3 at Serine 10 Is Mediated by Akt1, Extracellular Signal-regulated Kinase 2, and p90 Ribosomal S6 Kinase 2 but Not Mitogen- and Stress-activated Protein Kinase 1. <i>Journal of Biological Chemistry</i> , 2003, 278, 10588-10593.	3.4	48
18	The selective post-translational processing of transcription factor Nrf1 yields distinct isoforms that dictate its ability to differentially regulate gene expression. <i>Scientific Reports</i> , 2015, 5, 12983.	3.3	48

#	ARTICLE	IF	CITATIONS
19	Ataxia Telangiectasia Mutated Proteins, MAPKs, and RSK2 Are Involved in the Phosphorylation of STAT3. <i>Journal of Biological Chemistry</i> , 2003, 278, 12650-12659.	3.4	47
20	Nrf1 is paved as a new strategic avenue to prevent and treat cancer, neurodegenerative and other diseases. <i>Toxicology and Applied Pharmacology</i> , 2018, 360, 273-283.	2.8	46
21	TALENs-directed knockout of the full-length transcription factor Nrf1 [±] that represses malignant behaviour of human hepatocellular carcinoma (HepG2) cells. <i>Scientific Reports</i> , 2016, 6, 23775.	3.3	44
22	Oncogenic Activation of Nrf2, Though as a Master Antioxidant Transcription Factor, Liberated by Specific Knockout of the Full-Length Nrf1 [±] that Acts as a Dominant Tumor Repressor. <i>Cancers</i> , 2018, 10, 520.	3.7	42
23	The signal transduction networks required for phosphorylation of STAT1 at Ser727 in mouse epidermal JB6 cells in the UVB response and inhibitory mechanisms of tea polyphenols. <i>Carcinogenesis</i> , 2004, 26, 331-342.	2.8	39
24	The membrane-topogenic vectorial behaviour of Nrf1 controls its post-translational modification and transactivation activity. <i>Scientific Reports</i> , 2013, 3, 2006.	3.3	39
25	Mechanisms controlling the multistage post-translational processing of endogenous Nrf1 [±] /TCF11 proteins to yield distinct isoforms within the coupled positive and negative feedback circuits. <i>Toxicology and Applied Pharmacology</i> , 2018, 360, 212-235.	2.8	39
26	Unification of Opposites between Two Antioxidant Transcription Factors Nrf1 and Nrf2 in Mediating Distinct Cellular Responses to the Endoplasmic Reticulum Stressor Tunicamycin. <i>Antioxidants</i> , 2020, 9, 4.	5.1	39
27	Evidence of STAT1 phosphorylation modulated by MAPKs, MEK1 and MSK1. <i>Carcinogenesis</i> , 2004, 25, 1165-1175.	2.8	35
28	Adipocyte-specific deficiency of Nfe2l1 disrupts plasticity of white adipose tissues and metabolic homeostasis in mice. <i>Biochemical and Biophysical Research Communications</i> , 2018, 503, 264-270.	2.1	35
29	UVA, UVB and UVC Induce Differential Response Signaling Pathways Converged on the eIF2 [±] Phosphorylation. <i>Photochemistry and Photobiology</i> , 2011, 87, 1092-1104.	2.5	33
30	Mechanisms Underlying Chemopreventive Effects of Flavonoids via Multiple Signaling Nodes within Nrf2-ARE and AhR-XRE Gene Regulatory Networks. <i>Current Chemical Biology</i> , 2013, 7, 151-176.	0.5	29
31	An optimal tumor marker group-coupled artificial neural network for diagnosis of lung cancer. <i>Expert Systems With Applications</i> , 2011, 38, 11329-11334.	7.6	27
32	Transcription factor Nrf1 is negatively regulated by its <i>ε</i> -GlcNAcylation status. <i>FEBS Letters</i> , 2015, 589, 2347-2358.	2.8	26
33	Distinct isoforms of Nrf1 diversely regulate different subsets of its cognate target genes. <i>Scientific Reports</i> , 2019, 9, 2960.	3.3	26
34	Induction of EGFR-Dependent and EGFR-Independent Signaling Pathways by Ultraviolet A Irradiation. <i>DNA and Cell Biology</i> , 2001, 20, 769-779.	1.9	25
35	NS-398 and Piroxicam Suppress UVB-induced Activator Protein 1 Activity by Mechanisms Independent of Cyclooxygenase-2. <i>Journal of Biological Chemistry</i> , 2003, 278, 2124-2130.	3.4	23
36	Involvement of ERKs, RSK2 and PKR in UVA-induced signal transduction toward phosphorylation of eIF2 [±] (Ser51). <i>Carcinogenesis</i> , 2007, 28, 1543-1551.	2.8	23

#	ARTICLE	IF	CITATIONS
37	Novel flavagline-like compounds with potent Flt3 inhibitory activity suppress diverse types of leukemia. <i>FEBS Journal</i> , 2018, 285, 4631-4645.	4.7	22
38	The C-Terminal Domain of Nrf1 Negatively Regulates the Full-Length CNC-bZIP Factor and Its Shorter Isoform LCR-F1/Nrf1 ^{Δ2} ; Both Are Also Inhibited by the Small Dominant-Negative Nrf1 ^{Δ3} /Nrf1 ^{Δ4} Isoforms that Down-Regulate ARE-Battery Gene Expression. <i>PLoS ONE</i> , 2014, 9, e109159.	2.5	21
39	Topovectorial mechanisms control the juxtamembrane proteolytic processing of Nrf1 to remove its N-terminal polypeptides during maturation of the CNC-bZIP factor. <i>Toxicology and Applied Pharmacology</i> , 2018, 360, 160-184.	2.8	21
40	Involvement of ERKs and mitogen- and stress-activated protein kinase in UVC-induced phosphorylation of ATF2 in JB6 cells. <i>Carcinogenesis</i> , 2004, 25, 1847-1852.	2.8	17
41	Nrf1 Is Endowed with a Dominant Tumor-Repressing Effect onto the Wnt/β-Catenin-Dependent and Wnt/β-Catenin-Independent Signaling Networks in the Human Liver Cancer. <i>Oxidative Medicine and Cellular Longevity</i> , 2020, 2020, 1-28.	4.0	15
42	Nach Is a Novel Subgroup at an Early Evolutionary Stage of the CNC-bZIP Subfamily Transcription Factors from the Marine Bacteria to Humans. <i>International Journal of Molecular Sciences</i> , 2018, 19, 2927.	4.1	14
43	Differential Yet Integral Contributions of Nrf1 and Nrf2 in the Human HepG2 Cells on Antioxidant Cytoprotective Response against Tert-Butylhydroquinone as a Pro-Oxidative Stressor. <i>Antioxidants</i> , 2021, 10, 1610.	5.1	14
44	The Role of MicroRNA in the Regulation of Tumor Epithelial-Mesenchymal Transition. <i>Cells</i> , 2022, 11, 1981.	4.1	14
45	Dysfunction of the energy sensor NFE2L1 triggers uncontrollable AMPK signaling and glucose metabolism reprogramming. <i>Cell Death and Disease</i> , 2022, 13, .	6.3	13
46	Glucose Starvation-Induced Rapid Death of Nrf1-Deficient, but Not Nrf2-Deficient, Hepatoma Cells Results from Its Fatal Defects in the Redox Metabolism Reprogramming. <i>Oxidative Medicine and Cellular Longevity</i> , 2020, 2020, 1-20.	4.0	12
47	Metformin leads to accumulation of reactive oxygen species by inhibiting the NFE2L1 expression in human hepatocellular carcinoma cells. <i>Toxicology and Applied Pharmacology</i> , 2021, 420, 115523.	2.8	11
48	Organ-specific distribution of AP-1 in AP-1 luciferase transgenic mice during the maturation process. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2001, 280, R376-R381.	1.8	10
49	A microwell pattern for C17.2 cell aggregate formation with concave cylindrical surface induced cell peeling. <i>Biomaterials</i> , 2014, 35, 9423-9437.	11.4	10
50	TCF11 Has a Potent Tumor-Repressing Effect Than Its Prototypic Nrf1 by Definition of Both Similar Yet Different Regulatory Profiles, With a Striking Disparity From Nrf2. <i>Frontiers in Oncology</i> , 2021, 11, 707032.	2.8	10
51	Activation of the membrane-bound Nrf1 transcription factor by USP19, a ubiquitin-specific protease C-terminally anchored in the endoplasmic reticulum. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2022, 1869, 119299.	4.1	9
52	A Naturally-Occurring Dominant-Negative Inhibitor of Keap1 Competitively against Its Negative Regulation of Nrf2. <i>International Journal of Molecular Sciences</i> , 2018, 19, 2150.	4.1	8
53	eIF2 alpha phosphorylation alleviates UVA-induced HO-1 expression in mouse epidermal cells. <i>Free Radical Research</i> , 2018, 52, 1359-1370.	3.3	6
54	NF-E2-related factor 2 serves a key function in resistance to malignant transformation of BEAS-2B cells induced by coal tar pitch. <i>Oncology Letters</i> , 2018, 15, 5143-5148.	1.8	5

#	ARTICLE	IF	CITATIONS
55	Synergism and Antagonism of Two Distinct, but Confused, Nrf1 Factors in Integral Regulation of the Nuclear-to-Mitochondrial Respiratory and Antioxidant Transcription Networks. <i>Oxidative Medicine and Cellular Longevity</i> , 2020, 2020, 1-33.	4.0	4
56	Commentary on Distinct, but Previously Confused, Nrf1 Transcription Factors and Their Functions in Redox Regulation. <i>Developmental Cell</i> , 2020, 53, 377-378.	7.0	4
57	Nrf1D Is the First Candidate Secretory Transcription Factor in the Blood Plasma, Its Precursor Existing as a Unique Redox-Sensitive Transmembrane CNC-bZIP Protein in Hemopoietic and Somatic Tissues. <i>International Journal of Molecular Sciences</i> , 2018, 19, 2940.	4.1	2
58	Oxidative stress and the Nrf1 and Nrf2 transcription factors. <i>Toxicology Letters</i> , 2007, 172, S10.	0.8	0
59	Abstract 4556: Activation of sonic hedgehog signaling is essential for non-alcoholic steatohepatitis induced by liver-specific disruption of Nrf1. , 2016, , .		0