

Fumiaki Katagiri

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/906073/publications.pdf>

Version: 2024-02-01

70
papers

12,209
citations

70961

41
h-index

91712

69
g-index

73
all docs

73
docs citations

73
times ranked

12568
citing authors

#	ARTICLE	IF	CITATIONS
1	Pathogen-driven coevolution across the CBP60 plant immune regulator subfamilies confers resilience on the regulator module. <i>New Phytologist</i> , 2022, 233, 479-495.	3.5	14
2	Letter to the Editor: DNA Purification-Free PCR from Plant Tissues. <i>Plant and Cell Physiology</i> , 2021, 62, 1503-1505.	1.5	9
3	Environmental Association Identifies Candidates for Tolerance to Low Temperature and Drought. G3: Genes, Genomes, Genetics, 2019, 9, 3423-3438.	0.8	18
4	Different Modes of Negative Regulation of Plant Immunity by Calmodulin-Related Genes. <i>Plant Physiology</i> , 2018, 176, 3046-3061.	2.3	31
5	WRKY70 prevents axenic activation of plant immunity by direct repression of <i>SARD1</i> . <i>New Phytologist</i> , 2018, 217, 700-712.	3.5	60
6	Involvement of Adapter Protein Complex 4 in Hypersensitive Cell Death Induced by Avirulent Bacteria. <i>Plant Physiology</i> , 2018, 176, 1824-1834.	2.3	25
7	Review: Plant immune signaling from a network perspective. <i>Plant Science</i> , 2018, 276, 14-21.	1.7	17
8	Quantification of Plant Cell Death by Electrolyte Leakage Assay. <i>Bio-protocol</i> , 2018, 8, e2758.	0.2	50
9	Nup82 functions redundantly with Nup136 in a salicylic acid-dependent defense response of <i>Arabidopsis thaliana</i> . <i>Nucleus</i> , 2017, 8, 301-311.	0.6	16
10	A plant effector-triggered immunity signaling sector is inhibited by pattern-triggered immunity. <i>EMBO Journal</i> , 2017, 36, 2758-2769.	3.5	69
11	The highly buffered <i>Arabidopsis</i> immune signaling network conceals the functions of its components. <i>PLoS Genetics</i> , 2017, 13, e1006639.	1.5	138
12	Network Reconstitution for Quantitative Subnetwork Interaction Analysis. <i>Methods in Molecular Biology</i> , 2017, 1578, 223-231.	0.4	3
13	The $\frac{1}{4}$ Subunit of <i>Arabidopsis</i> Adaptor Protein-2 Is Involved in Effector-Triggered Immunity Mediated by Membrane-Localized Resistance Proteins. <i>Molecular Plant-Microbe Interactions</i> , 2016, 29, 345-351.	1.4	24
14	Pectin Biosynthesis Is Critical for Cell Wall Integrity and Immunity in <i>Arabidopsis thaliana</i> . <i>Plant Cell</i> , 2016, 28, 537-556.	3.1	144
15	Toward predictive modeling of large and complex biological signaling networks. <i>Physiological and Molecular Plant Pathology</i> , 2016, 95, 77-83.	1.3	3
16	The receptor-like cytoplasmic kinase <i>PCRK1</i> contributes to pattern-triggered immunity against <i>Pseudomonas syringae</i> in <i>Arabidopsis thaliana</i> . <i>New Phytologist</i> , 2015, 207, 78-90.	3.5	50
17	Putative Serine Protease Effectors of <i>Clavibacter michiganensis</i> Induce a Hypersensitive Response in the Apoplast of <i>Nicotiana</i> Species. <i>Molecular Plant-Microbe Interactions</i> , 2015, 28, 1216-1226.	1.4	32
18	Identification of differentially expressed genes between developing seeds of different soybean cultivars. <i>Genomics Data</i> , 2015, 6, 92-98.	1.3	3

#	ARTICLE	IF	CITATIONS
19	Design and Construction of an Inexpensive Homemade Plant Growth Chamber. <i>PLoS ONE</i> , 2015, 10, e0126826.	1.1	16
20	Mechanisms Underlying Robustness and Tunability in a Plant Immune Signaling Network. <i>Cell Host and Microbe</i> , 2014, 15, 84-94.	5.1	117
21	<i>Arabidopsis</i> <i>PECTIN METHYLESTERASEs</i> Contribute to Immunity against <i>Pseudomonas syringae</i> . <i>Plant Physiology</i> , 2014, 164, 1093-1107.	2.3	166
22	The CALMODULIN-BINDING PROTEIN60 Family Includes Both Negative and Positive Regulators of Plant Immunity. <i>Plant Physiology</i> , 2013, 163, 1741-1751.	2.3	91
23	Dual Regulation of Gene Expression Mediated by Extended MAPK Activation and Salicylic Acid Contributes to Robust Innate Immunity in <i>Arabidopsis thaliana</i> . <i>PLoS Genetics</i> , 2013, 9, e1004015.	1.5	208
24	BR-SIGNALING KINASE1 Physically Associates with FLAGELLIN SENSING2 and Regulates Plant Innate Immunity in <i>Arabidopsis</i> A. <i>Plant Cell</i> , 2013, 25, 1143-1157.	3.1	212
25	Spatio-Temporal Expression Patterns of <i>Arabidopsis thaliana</i> and <i>Medicago truncatula</i> Defensin-Like Genes. <i>PLoS ONE</i> , 2013, 8, e58992.	1.1	54
26	Pattern-Triggered Immunity Suppresses Programmed Cell Death Triggered by Fumonisin B1. <i>PLoS ONE</i> , 2013, 8, e60769.	1.1	30
27	Membrane microdomain may be a platform for immune signaling. <i>Plant Signaling and Behavior</i> , 2012, 7, 454-456.	1.2	15
28	Activation of the <i>Arabidopsis thaliana</i> Mitogen-Activated Protein Kinase MPK11 by the Flagellin-Derived Elicitor Peptide, flg22. <i>Molecular Plant-Microbe Interactions</i> , 2012, 25, 471-480.	1.4	123
29	An efficient <i>Agrobacterium</i> -mediated transient transformation of <i>Arabidopsis</i> . <i>Plant Journal</i> , 2012, 69, 713-719.	2.8	95
30	The peptide growth factor, phytoalexin, attenuates pattern-triggered immunity. <i>Plant Journal</i> , 2012, 71, 194-204.	2.8	128
31	<i>Arabidopsis</i> lysin-motif proteins LYM1 LYM3 CERK1 mediate bacterial peptidoglycan sensing and immunity to bacterial infection. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 19824-19829.	3.3	442
32	CO ₂ -mediated changes of plant traits and their effects on herbivores are determined by leaf age. <i>Ecological Entomology</i> , 2011, 36, 1-13.	1.1	17
33	CBP60g and SARD1 play partially redundant critical roles in salicylic acid signaling. <i>Plant Journal</i> , 2011, 67, 1029-1041.	2.8	244
34	Physical association of pattern-triggered immunity (PTI) and effector-triggered immunity (ETI) immune receptors in <i>Arabidopsis</i> . <i>Molecular Plant Pathology</i> , 2011, 12, 702-708.	2.0	91
35	Identification and utilization of a sow thistle powdery mildew as a poorly adapted pathogen to dissect post-invasion non-host resistance mechanisms in <i>Arabidopsis</i> . <i>Journal of Experimental Botany</i> , 2011, 62, 2117-2129.	2.4	39
36	Physical Association of <i>Arabidopsis</i> Hypersensitive Induced Reaction Proteins (HIRs) with the Immune Receptor RPS2. <i>Journal of Biological Chemistry</i> , 2011, 286, 31297-31307.	1.6	94

#	ARTICLE	IF	CITATIONS
37	Purification of Resistance Protein Complexes Using a Biotinylated Affinity (HPB) Tag. <i>Methods in Molecular Biology</i> , 2011, 712, 21-30.	0.4	3
38	A Putative RNA-Binding Protein Positively Regulates Salicylic Acid-Mediated Immunity in <i>Arabidopsis</i> . <i>Molecular Plant-Microbe Interactions</i> , 2010, 23, 1573-1583.	1.4	45
39	Comparing signaling mechanisms engaged in pattern-triggered and effector-triggered immunity. <i>Current Opinion in Plant Biology</i> , 2010, 13, 459-465.	3.5	705
40	Network Modeling Reveals Prevalent Negative Regulatory Relationships between Signaling Sectors in <i>Arabidopsis</i> Immune Signaling. <i>PLoS Pathogens</i> , 2010, 6, e1001011.	2.1	110
41	Endosome-Associated CRT1 Functions Early in Resistance Gene-Mediated Defense Signaling in <i>Arabidopsis</i> and Tobacco. <i>Plant Cell</i> , 2010, 22, 918-936.	3.1	55
42	Understanding the Plant Immune System. <i>Molecular Plant-Microbe Interactions</i> , 2010, 23, 1531-1536.	1.4	212
43	<i>Arabidopsis</i> CaM Binding Protein CBP60g Contributes to MAMP-Induced SA Accumulation and Is Involved in Disease Resistance against <i>Pseudomonas syringae</i> . <i>PLoS Pathogens</i> , 2009, 5, e1000301.	2.1	242
44	Network Properties of Robust Immunity in Plants. <i>PLoS Genetics</i> , 2009, 5, e1000772.	1.5	489
45	Purification of low-abundance <i>Arabidopsis</i> plasma-membrane protein complexes and identification of candidate components. <i>Plant Journal</i> , 2009, 57, 932-944.	2.8	85
46	Overview of mRNA Expression Profiling Using DNA Microarrays. <i>Current Protocols in Molecular Biology</i> , 2009, 85, Unit 22.4.	2.9	14
47	Pattern Discovery in Expression Profiling Data. <i>Current Protocols in Molecular Biology</i> , 2009, 85, Unit 22.5.	2.9	8
48	<i>Arabidopsis</i> defense response against <i>Pseudomonas syringae</i> - Effects of major regulatory genes and the impact of coronatine. , 2009, , .		0
49	Interplay between MAMP-triggered and SA-mediated defense responses. <i>Plant Journal</i> , 2008, 53, 763-775.	2.8	318
50	Unsupervised reduction of random noise in complex data by a row-specific, sorted principal component-guided method. <i>BMC Bioinformatics</i> , 2008, 9, 508.	1.2	3
51	The interplay between MAMP and SA signaling. <i>Plant Signaling and Behavior</i> , 2008, 3, 359-361.	1.2	33
52	Natural Variation in RPS2-Mediated Resistance among <i>Arabidopsis</i> Accessions: Correlation between Gene Expression Profiles and Phenotypic Responses. <i>Plant Cell</i> , 2008, 19, 4046-4060.	3.1	37
53	The Genetic Network Controlling the <i>Arabidopsis</i> Transcriptional Response to <i>Pseudomonas syringae</i> pv. <i>maculicola</i> : Roles of Major Regulators and the Phytotoxin Coronatine. <i>Molecular Plant-Microbe Interactions</i> , 2008, 21, 1408-1420.	1.4	64
54	Natural Variation among <i>Arabidopsis thaliana</i> Accessions for Transcriptome Response to Exogenous Salicylic Acid. <i>Plant Cell</i> , 2007, 19, 2099-2110.	3.1	101

#	ARTICLE	IF	CITATIONS
55	A high-performance, small-scale microarray for expression profiling of many samples in Arabidopsis-pathogen studies. <i>Plant Journal</i> , 2007, 49, 565-577.	2.8	51
56	Expression profiles as detailed snapshots of biological states. <i>Transgenic Research</i> , 2007, 16, 399-403.	1.3	2
57	Pattern Discovery in Expression Profiling Data. <i>Current Protocols in Molecular Biology</i> , 2005, 69, Unit 22.5.	2.9	1
58	A global view of defense gene expression regulation â€” a highly interconnected signaling network. <i>Current Opinion in Plant Biology</i> , 2004, 7, 506-511.	3.5	133
59	Overview of mRNA Expression Profiling Using Microarrays. <i>Current Protocols in Molecular Biology</i> , 2004, 67, Unit 22.4.	2.9	3
60	Topology of the network integrating salicylate and jasmonate signal transduction derived from global expression phenotyping. <i>Plant Journal</i> , 2003, 34, 217-228.	2.8	466
61	Direct delivery of bacterial avirulence proteins into resistant Arabidopsis protoplasts leads to hypersensitive cell death. <i>Plant Journal</i> , 2003, 33, 131-137.	2.8	35
62	Local Context Finder (LCF) reveals multidimensional relationships among mRNA expression profiles of Arabidopsis responding to pathogen infection. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 10842-10847.	3.3	19
63	Quantitative Nature of Arabidopsis Responses during Compatible and Incompatible Interactions with the Bacterial Pathogen <i>Pseudomonas syringae</i> [W]. <i>Plant Cell</i> , 2003, 15, 317-330.	3.1	641
64	The Arabidopsis <i>Thaliana</i> - <i>Pseudomonas Syringae</i> Interaction. <i>The Arabidopsis Book</i> , 2002, 1, e0039.	0.5	421
65	A High-Throughput Arabidopsis Reverse Genetics System. <i>Plant Cell</i> , 2002, 14, 2985-2994.	3.1	873
66	A Draft Sequence of the Rice Genome (<i>Oryza sativa</i> L. ssp. japonica). <i>Science</i> , 2002, 296, 92-100.	6.0	2,866
67	The <i>Pseudomonas syringae</i> avrRpt2 Gene Product Promotes Pathogen Virulence from Inside Plant Cells. <i>Molecular Plant-Microbe Interactions</i> , 2000, 13, 1312-1321.	1.4	122
68	Mutational Analysis of the Arabidopsis Nucleotide Binding Siteâ€”Leucine-Rich Repeat Resistance Gene RPS2. <i>Plant Cell</i> , 2000, 12, 2541-2554.	3.1	166
69	Eukaryotic Fatty Acylation Drives Plasma Membrane Targeting and Enhances Function of Several Type III Effector Proteins from <i>Pseudomonas syringae</i> . <i>Cell</i> , 2000, 101, 353-363.	13.5	308
70	The <i>A. thaliana</i> disease resistance gene RPS2 encodes a protein containing a nucleotide-binding site and leucine-rich repeats. <i>Cell</i> , 1994, 78, 1089-1099.	13.5	689