

El ChÃ©rif Ibrahim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9083588/publications.pdf>

Version: 2024-02-01

42
papers

1,958
citations

279701

23
h-index

276775

41
g-index

47
all docs

47
docs citations

47
times ranked

2967
citing authors

#	ARTICLE	IF	CITATIONS
1	Identification of HLA-G7 as a new splice variant of the HLA-G mRNA and expression of soluble HLA-G5, -G6, and -G7 transcripts in human transfected cells. <i>Human Immunology</i> , 2000, 61, 1138-1149.	1.2	292
2	Responder and nonresponder patients exhibit different peripheral transcriptional signatures during major depressive episode. <i>Translational Psychiatry</i> , 2012, 2, e185-e185.	2.4	192
3	Tumor-specific up-regulation of the nonclassical class I HLA-G antigen expression in renal carcinoma. <i>Cancer Research</i> , 2001, 61, 6838-45.	0.4	124
4	Serine/arginine-rich protein-dependent suppression of exon skipping by exonic splicing enhancers. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 5002-5007.	3.3	109
5	Dissociating Bottom-Up and Top-Down Mechanisms in the Cortico-Limbic System during Emotion Processing. <i>Cerebral Cortex</i> , 2016, 26, 144-155.	1.6	105
6	Analysis of HLA antigen expression in benign and malignant melanocytic lesions reveals that upregulation of HLA-G expression correlates with malignant transformation, high inflammatory infiltration and HLA-A1 genotype. <i>International Journal of Cancer</i> , 2004, 108, 243-250.	2.3	89
7	Heat shock and arsenite induce expression of the nonclassical class I histocompatibility HLA-G gene in tumor cell lines. <i>Cell Stress and Chaperones</i> , 2000, 5, 207.	1.2	79
8	Clinical variations modulate patterns of gene expression and define blood biomarkers in major depression. <i>Journal of Psychiatric Research</i> , 2010, 44, 1205-1213.	1.5	77
9	Glucocorticoid hormones upregulate levels of HLA-G transcripts in trophoblasts. <i>Transplantation Proceedings</i> , 2001, 33, 2277-2280.	0.3	72
10	Therapeutic potential and mechanism of kinetin as a treatment for the human splicing disease familial dysautonomia. <i>Journal of Molecular Medicine</i> , 2007, 85, 149-161.	1.7	58
11	Molecular mechanisms controlling constitutive and IFN- γ -inducible HLA-G expression in various cell types. <i>Journal of Reproductive Immunology</i> , 1999, 43, 213-224.	0.8	56
12	HLA-G expression in human melanoma cells: protection from NK cytotoxicity. <i>Journal of Reproductive Immunology</i> , 1999, 43, 183-193.	0.8	54
13	CX3CR1 is dysregulated in blood and brain from schizophrenia patients. <i>Schizophrenia Research</i> , 2015, 168, 434-443.	1.1	49
14	Severity of experimental autoimmune encephalomyelitis is unexpectedly reduced in mice born to vitamin D-deficient mothers. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2010, 121, 250-253.	1.2	47
15	miR-323a regulates ERBB4 and is involved in depression. <i>Molecular Psychiatry</i> , 2021, 26, 4191-4204.	4.1	47
16	Olfactory Stem Cells, a New Cellular Model for Studying Molecular Mechanisms Underlying Familial Dysautonomia. <i>PLoS ONE</i> , 2010, 5, e15590.	1.1	46
17	Altered Pattern of Major Histocompatibility Complex Expression in Renal Carcinoma. <i>American Journal of Pathology</i> , 2003, 162, 501-508.	1.9	44
18	Weak definition of KKBAPexon 20 leads to aberrant splicing in familial dysautonomia. <i>Human Mutation</i> , 2007, 28, 41-53.	1.1	44

#	ARTICLE	IF	CITATIONS
19	How to: Measuring blood cytokines in biological psychiatry using commercially available multiplex immunoassays. <i>Psychoneuroendocrinology</i> , 2017, 75, 72-82.	1.3	38
20	GPR56/ADGRG1 is associated with response to antidepressant treatment. <i>Nature Communications</i> , 2020, 11, 1635.	5.8	38
21	Antidepressive effects of targeting ELK-1 signal transduction. <i>Nature Medicine</i> , 2018, 24, 591-597.	15.2	33
22	Mood disorders are associated with a more severe hypovitaminosis D than schizophrenia. <i>Psychiatry Research</i> , 2015, 229, 613-616.	1.7	30
23	Predisposition to treatment response in major depressive episode: A peripheral blood gene coexpression network analysis. <i>Journal of Psychiatric Research</i> , 2016, 81, 119-126.	1.5	29
24	Translational Identification of Transcriptional Signatures of Major Depression and Antidepressant Response. <i>Frontiers in Molecular Neuroscience</i> , 2017, 10, 248.	1.4	29
25	Deficits in Social Behavior Precede Cognitive Decline in Middle-Aged Mice. <i>Frontiers in Behavioral Neuroscience</i> , 2019, 13, 55.	1.0	26
26	Genome-wide analysis of familial dysautonomia and kinetin target genes with patient olfactory ecto-mesenchymal stem cells. <i>Human Mutation</i> , 2012, 33, 530-540.	1.1	25
27	Serotonin transporter gene expression predicts the worsening of suicidal ideation and suicide attempts along a long-term follow-up of a Major Depressive Episode. <i>European Neuropsychopharmacology</i> , 2018, 28, 401-414.	0.3	18
28	Impaired cortico-limbic functional connectivity in schizophrenia patients during emotion processing. <i>Social Cognitive and Affective Neuroscience</i> , 2018, 13, 381-390.	1.5	17
29	MicroRNA screening identifies a link between NOVA1 expression and low level of IKAP/ELP1 in Familial dysautonomia. <i>DMM Disease Models and Mechanisms</i> , 2016, 9, 899-909.	1.2	16
30	Suicide and suicide behaviors: A review of transcriptomics and multiomics studies in psychiatric disorders. <i>Journal of Neuroscience Research</i> , 2020, 98, 601-615.	1.3	14
31	The NKG2D Ligands RAE-1 γ and RAE-1 μ Differ with Respect to Their Receptor Affinity, Expression Profiles and Transcriptional Regulation. <i>PLoS ONE</i> , 2010, 5, e13466.	1.1	10
32	Monitoring candidate gene expression variations before, during and after a first major depressive episode in a 51-year-old man. <i>BMC Psychiatry</i> , 2014, 14, 73.	1.1	8
33	Modeling a linkage between blood transcriptional expression and activity in brain regions to infer the phenotype of schizophrenia patients. <i>NPJ Schizophrenia</i> , 2017, 3, 25.	2.0	8
34	Proteasome inhibitors to alleviate aberrant IKBKAP mRNA splicing and low IKAP/hELP1 synthesis in familial dysautonomia. <i>Neurobiology of Disease</i> , 2017, 103, 113-122.	2.1	7
35	Longitudinal Monitoring of the Serotonin Transporter Gene Expression to Assess Major Depressive Episode Evolution. <i>Neuropsychobiology</i> , 2014, 70, 220-227.	0.9	6
36	Blood cytokines differentiate bipolar disorder and major depressive disorder during a major depressive episode: Initial discovery and independent sample replication. <i>Brain, Behavior, & Immunity - Health</i> , 2021, 13, 100232.	1.3	6

#	ARTICLE	IF	CITATIONS
37	Immune-Related Genetic Overlap Between Regional Gray Matter Reductions and Psychiatric Symptoms in Adolescents, and Gene-Set Validation in a Translational Model. <i>Frontiers in Systems Neuroscience</i> , 2021, 15, 725413.	1.2	4
38	Treatment of Comorbid Bipolar Disorder Improves Disabilities and Neuropsychological Functioning in DiGeorge Syndrome. <i>Journal of Clinical Psychopharmacology</i> , 2017, 37, 736-738.	0.7	3
39	Baseline levels of C-reactive protein and proinflammatory cytokines are not associated with early response to amisulpride in patients with First Episode Psychosis: the OPTiMiSE cohort study. <i>Schizophrenia Bulletin Open</i> , 0, , .	0.9	2
40	RNA therapeutics for mood disorders: current evidence toward clinical trials. <i>Expert Opinion on Investigational Drugs</i> , 2021, 30, 721-736.	1.9	2
41	Biomarker development: Current issues and perspectives. <i>Psychoneuroendocrinology</i> , 2017, 78, 253-254.	1.3	0
42	203. ELK-1, Trauma and Negative Life Event Exposure. <i>Biological Psychiatry</i> , 2019, 85, S84.	0.7	0